

Simultaneous statistical inference for epidemic trends: the case of COVID-19

Marina Khismatullina Michael Vogt

01/11/2020


Table of contents

1. Introduction
2. Model
3. Testing procedure
4. Theoretical properties
5. Application

Introduction


Motivation

Research question: How do outbreak patterns of COVID-19 compare across countries?




Motivation

Research question: How do outbreak patterns of COVID-19 compare across countries?




Motivation

Research question: How do outbreak patterns of COVID-19 compare across countries?



Motivation

Research question: How do outbreak patterns of COVID-19 compare across countries?



Aim of the paper

To develop new inference methods that allow to *identify* and *locate* differences between epidemic time trends.

Research goal

Suppose we have a set of intervals of different lengths, and we want to be able to say if two underlying trend functions are equal on each of these intervals.

Research goal

Suppose we have a set of intervals of different lengths, and we want to be able to say if two underlying trend functions are equal on each of these intervals.

Then we would like a procedure that:

- allows us to test multiple hypotheses simultaneously;

Research goal

Suppose we have a set of intervals of different lengths, and we want to be able to say if two underlying trend functions are equal on each of these intervals.

Then we would like a procedure that:

- allows us to test multiple hypotheses simultaneously;
- taking the differences in lengths of the intervals into account;

Research goal

Suppose we have a set of intervals of different lengths, and we want to be able to say if two underlying trend functions are equal on each of these intervals.

Then we would like a procedure that:

- allows us to test multiple hypotheses simultaneously;
- taking the differences in lengths of the intervals into account;
- controls the family-wise error rate (probability of making even one type I error);

Research goal

Suppose we have a set of intervals of different lengths, and we want to be able to say if two underlying trend functions are equal on each of these intervals.

Then we would like a procedure that:

- allows us to test multiple hypotheses simultaneously;
- taking the differences in lengths of the intervals into account;
- controls the family-wise error rate (probability of making even one type I error);
- while having an asymptotic power of 1 against local alternatives;

Suppose we have a set of intervals of different lengths, and we want to be able to say if two underlying trend functions are equal on each of these intervals.

Then we would like a procedure that:

- allows us to test multiple hypotheses simultaneously;
- taking the differences in lengths of the intervals into account;
- controls the family-wise error rate (probability of making even one type I error);
- while having an asymptotic power of 1 against local alternatives;
- is easy to use and ideally already has an implementation as an R-package.

Suppose we have a set of intervals of different lengths, and we want to be able to say if two underlying trend functions are equal on each of these intervals.

Then we would like a procedure that:

- allows us to test multiple hypotheses simultaneously;
- taking the differences in lengths of the intervals into account;
- controls the family-wise error rate (probability of making even one type I error);
- while having an asymptotic power of 1 against local alternatives;
- is easy to use and ideally already has an implementation as an R-package.

Comparison of deterministic trends:

- Park et al. (2009), Degras et al. (2012), Zhang et al. (2012), Hidalgo and Lee (2014), Chen and Wu (2019).

Comparison of deterministic trends:

- Park et al. (2009), Degras et al. (2012), Zhang et al. (2012), Hidalgo and Lee (2014), Chen and Wu (2019).

Studies of COVID-19:

- Time series analysis: Gu et al. (2020), Li and Linton (2020).

Model

Motivation for the model

We observe n time series $\mathcal{X}_i = \{X_{it} : 1 \leq t \leq T\}$ of length T .

Motivation for the model

We observe n time series $\mathcal{X}_i = \{X_{it} : 1 \leq t \leq T\}$ of length T .

X_{it} are non-negative integers \Rightarrow can be modelled by a Poisson distribution with time-varying parameter $\lambda_i(t/T)$: $X_{it} \sim P_{\lambda_i(t/T)}$.

Motivation for the model

We observe n time series $\mathcal{X}_i = \{X_{it} : 1 \leq t \leq T\}$ of length T .

X_{it} are non-negative integers \Rightarrow can be modelled by a Poisson distribution with time-varying parameter $\lambda_i(t/T)$: $X_{it} \sim P_{\lambda_i(t/T)}$.

Since $\lambda_i(t/T) = \mathbb{E}[X_{it}] = \text{Var}(X_{it})$, we can rewrite X_{it} as

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + u_{it} \quad \text{with} \quad u_{it} = \sqrt{\lambda_i\left(\frac{t}{T}\right)} \eta_{it}$$

where η_{it} has zero mean and unit variance.

Motivation for the model

We observe n time series $\mathcal{X}_i = \{X_{it} : 1 \leq t \leq T\}$ of length T .

X_{it} are non-negative integers \Rightarrow can be modelled by a Poisson distribution with time-varying parameter $\lambda_i(t/T)$: $X_{it} \sim P_{\lambda_i(t/T)}$.

Since $\lambda_i(t/T) = \mathbb{E}[X_{it}] = \text{Var}(X_{it})$, we can rewrite X_{it} as

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + u_{it} \quad \text{with} \quad u_{it} = \sqrt{\lambda_i\left(\frac{t}{T}\right)} \eta_{it}$$

where η_{it} has zero mean and unit variance.

In applications the variance can be larger than the mean \Rightarrow quasi-Poisson models.

Quasi-Poisson model:

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + \sigma\sqrt{\lambda_i\left(\frac{t}{T}\right)}\eta_{it},$$

Quasi-Poisson model:

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + \sigma \sqrt{\lambda_i\left(\frac{t}{T}\right)} \eta_{it},$$

where

- λ_i are unknown trend functions on $[0, 1]$;

Quasi-Poisson model:

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + \sigma \sqrt{\lambda_i\left(\frac{t}{T}\right)} \eta_{it},$$

where

- λ_i are unknown trend functions on $[0, 1]$;
- σ is the overdispersion parameter;

Quasi-Poisson model:

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + \sigma \sqrt{\lambda_i\left(\frac{t}{T}\right)} \eta_{it},$$

where

- λ_i are unknown trend functions on $[0, 1]$;
- σ is the overdispersion parameter;
- η_{it} are error terms that are independent across i and t and have zero mean and unit variance.

Testing procedure

Testing problem

Let $\mathcal{F} := \{\mathcal{I}_k \subseteq [0, 1] : 1 \leq k \leq K\}$ be a family of rescaled time intervals on $[0, 1]$, and let $H_0^{(ijk)}$ be the hypothesis that the functions λ_i and λ_j are equal on an interval \mathcal{I}_k ,

Testing problem

Let $\mathcal{F} := \{\mathcal{I}_k \subseteq [0, 1] : 1 \leq k \leq K\}$ be a family of rescaled time intervals on $[0, 1]$, and let $H_0^{(ijk)}$ be the hypothesis that the functions λ_i and λ_j are equal on an interval \mathcal{I}_k , i.e.

$$H_0^{(ijk)} : \lambda_i(w) = \lambda_j(w) \text{ for all } w \in \mathcal{I}_k$$

Testing problem

Let $\mathcal{F} := \{\mathcal{I}_k \subseteq [0, 1] : 1 \leq k \leq K\}$ be a family of rescaled time intervals on $[0, 1]$, and let $H_0^{(ijk)}$ be the hypothesis that the functions λ_i and λ_j are equal on an interval \mathcal{I}_k , i.e.

$$H_0^{(ijk)} : \lambda_i(w) = \lambda_j(w) \text{ for all } w \in \mathcal{I}_k$$

We want to test $H_0^{(ijk)}$ simultaneously for all pairs of countries i and j and all intervals \mathcal{I}_k in the family \mathcal{F} and we want to control the familywise error rate (FWER) at level α .

Testing problem

Let $\mathcal{F} := \{\mathcal{I}_k \subseteq [0, 1] : 1 \leq k \leq K\}$ be a family of rescaled time intervals on $[0, 1]$, and let $H_0^{(ijk)}$ be the hypothesis that the functions λ_i and λ_j are equal on an interval \mathcal{I}_k , i.e.

$$H_0^{(ijk)} : \lambda_i(w) = \lambda_j(w) \text{ for all } w \in \mathcal{I}_k$$

We want to test $H_0^{(ijk)}$ simultaneously for all pairs of countries i and j and all intervals \mathcal{I}_k in the family \mathcal{F} and we want to control the familywise error rate (FWER) at level α .

Let \mathcal{M}_0 be the set of triplets (i, j, k) for which $H_0^{(ijk)}$ holds true. Then, FWER is

$$\text{FWER}(\alpha) = \mathbb{P}\left(\exists (i, j, k) \in \mathcal{M}_0 : \text{we reject } H_0^{(ijk)}\right)$$

Test statistic

For the given interval \mathcal{I}_k and a pair of time series i and j we calculate

$$\hat{s}_{ijk} = \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} - X_{jt}),$$

where h_k is the length of \mathcal{I}_k .

Test statistic

For the given interval \mathcal{I}_k and a pair of time series i and j we calculate

$$\hat{s}_{ijk} = \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} - X_{jt}),$$

where h_k is the length of \mathcal{I}_k . \hat{s}_{ijk} estimates the average distance between λ_i and λ_j on \mathcal{I}_k :

Test statistic

For the given interval \mathcal{I}_k and a pair of time series i and j we calculate

$$\hat{s}_{ijk} = \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) (X_{it} - X_{jt}),$$

where h_k is the length of \mathcal{I}_k . \hat{s}_{ijk} estimates the average distance between λ_i and λ_j on \mathcal{I}_k :

$$\begin{aligned} \hat{s}_{ijk} = & \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \left(\lambda_i\left(\frac{t}{T}\right) - \lambda_j\left(\frac{t}{T}\right) \right) \\ & + \frac{\sigma}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \left(\sqrt{\lambda_i\left(\frac{t}{T}\right)} \eta_{it} - \sqrt{\lambda_j\left(\frac{t}{T}\right)} \eta_{jt} \right) \end{aligned}$$

Test statistic

For the given interval \mathcal{I}_k and a pair of time series i and j we calculate

$$\hat{s}_{ijk} = \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) (X_{it} - X_{jt}),$$

where h_k is the length of \mathcal{I}_k . \hat{s}_{ijk} estimates the average distance between λ_i and λ_j on \mathcal{I}_k :

$$\begin{aligned} \hat{s}_{ijk} = & \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \left(\lambda_i\left(\frac{t}{T}\right) - \lambda_j\left(\frac{t}{T}\right) \right) \\ & + \frac{\sigma}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \left(\sqrt{\lambda_i\left(\frac{t}{T}\right)} \eta_{it} - \sqrt{\lambda_j\left(\frac{t}{T}\right)} \eta_{jt} \right) \end{aligned}$$

Applying a law of large numbers, we get that

$$\hat{s}_{ijk} = \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \left(\lambda_i\left(\frac{t}{T}\right) - \lambda_j\left(\frac{t}{T}\right) \right) + o_P(1)$$

Under certain assumptions,

$$\text{Var}(\hat{s}_{ijk}) = \frac{\sigma^2}{T^2 h_k^2} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \left\{ \lambda_i\left(\frac{t}{T}\right) + \lambda_j\left(\frac{t}{T}\right) \right\}$$

Test statistic, part 2

Under certain assumptions,

$$\text{Var}(\hat{s}_{ijk}) = \frac{\sigma^2}{T^2 h_k^2} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \left\{ \lambda_i\left(\frac{t}{T}\right) + \lambda_j\left(\frac{t}{T}\right) \right\}$$

In order to normalize the variance of the statistic \hat{s}_{ijk} , we scale it by an estimator of its variance:

$$\widehat{\text{Var}}(\hat{s}_{ijk}) = \frac{\hat{\sigma}^2}{T^2 h_k^2} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) (X_{it} + X_{jt}),$$

with $\hat{\sigma}^2$ being an appropriate estimator of σ^2 . [Details](#)

Test statistic for the hypothesis $H_0^{(ijk)}$ is defined as

$$\hat{\psi}_{ijk} = \frac{\sum_{t=1}^T \mathbf{1}\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} - X_{jt})}{\hat{\sigma}\left\{\sum_{t=1}^T \mathbf{1}\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} + X_{jt})\right\}^{1/2}}$$

Test statistic, part 2

Test statistic for the hypothesis $H_0^{(ijk)}$ is defined as

$$\hat{\psi}_{ijk} = \frac{\sum_{t=1}^T \mathbf{1}\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} - X_{jt})}{\hat{\sigma}\left\{\sum_{t=1}^T \mathbf{1}\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} + X_{jt})\right\}^{1/2}}$$

Under certain conditions and under the null, $\hat{\psi}_{ijk}$ can be approximated by a Gaussian version of the test statistic:

$$\phi_{ijk} = \frac{1}{\sqrt{2Th_k}} \sum_{t=1}^T \mathbf{1}\left(\frac{t}{T} \in \mathcal{I}_k\right)(Z_{it} - Z_{jt}),$$

where Z_{it} are independent standard normal random variables.

How to construct critical values $c_{ijk}(\alpha)$?

How to construct critical values $c_{ijk}(\alpha)$?

- Traditional approach: $c_{ijk}(\alpha) = c(\alpha)$ for all (i, j, k) .

How to construct critical values $c_{ijk}(\alpha)$?

- Traditional approach: $c_{ijk}(\alpha) = c(\alpha)$ for all (i, j, k) .
- More modern approach: $c_{ijk}(\alpha)$ depend on the length h_k of the time interval (Dümbgen and Spokoiny (2001))

How to construct critical values $c_{ijk}(\alpha)$?

- Traditional approach: $c_{ijk}(\alpha) = c(\alpha)$ for all (i, j, k) .
- More modern approach: $c_{ijk}(\alpha)$ depend on the length h_k of the time interval (Dümbgen and Spokoiny (2001)):

$$c_{ijk}(\alpha) = c(\alpha, h_k) := b_k + q(\alpha)/a_k,$$

where a_k and b_k are scale-dependent constants and $q(\alpha)$ is chosen such that we control FWER. [Details](#)

Critical values, part 2

We want to control FWER:

$$\begin{aligned}\text{FWER}(\alpha) &= \mathbb{P}\left(\exists(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| > c_{ijk}(\alpha)\right) \\&= 1 - \mathbb{P}\left(\forall(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{ijk}(\alpha)\right) \\&= 1 - \mathbb{P}\left(\forall(i, j, k) \in \mathcal{M}_0 : a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&= 1 - \mathbb{P}\left(\max_{(i, j, k) \in \mathcal{M}_0} a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&\leq \alpha\end{aligned}$$

Critical values, part 2

We want to control FWER:

$$\begin{aligned}\text{FWER}(\alpha) &= \mathbb{P}\left(\exists(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| > c_{ijk}(\alpha)\right) \\&= 1 - \mathbb{P}\left(\forall(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{ijk}(\alpha)\right) \\&= 1 - \mathbb{P}\left(\forall(i, j, k) \in \mathcal{M}_0 : a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&= 1 - \mathbb{P}\left(\max_{(i, j, k) \in \mathcal{M}_0} a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&\leq \alpha\end{aligned}$$

Hence, we choose $q(\alpha)$ as the $(1 - \alpha)$ -quantile of the statistic

$$\hat{\Psi} = \max_{(i, j, k)} a_k(|\hat{\psi}_{ijk}^0| - b_k),$$

where $\hat{\psi}_{ijk}^0$ is equal to $\hat{\psi}_{ijk}$ under the null.

Test procedure

1. Consider the Gaussian test statistic

$$\Phi = \max_{(i,j,k)} a_k (|\phi_{ijk}| - b_k),$$

where a_k and b_k are scale-dependent constants and ϕ_{ijk} are weighted averages of the differences of standard normal random variables.

Test procedure

1. Consider the Gaussian test statistic

$$\Phi = \max_{(i,j,k)} a_k (|\phi_{ijk}| - b_k),$$

where a_k and b_k are scale-dependent constants and ϕ_{ijk} are weighted averages of the differences of standard normal random variables.

2. Compute a $(1 - \alpha)$ -quantile $q_{\text{Gauss}}(\alpha)$ of Φ by Monte Carlo simulations.

Test procedure

1. Consider the Gaussian test statistic

$$\Phi = \max_{(i,j,k)} a_k (|\phi_{ijk}| - b_k),$$

where a_k and b_k are scale-dependent constants and ϕ_{ijk} are weighted averages of the differences of standard normal random variables.

2. Compute a $(1 - \alpha)$ -quantile $q_{\text{Gauss}}(\alpha)$ of Φ by Monte Carlo simulations.
3. Adjust $q_{\text{Gauss}}(\alpha)$ by the scale-dependent constants

$$c_{\text{Gauss}}(\alpha, h_k) = b_k + q_{\text{Gauss}}(\alpha)/a_k$$

Test procedure

1. Consider the Gaussian test statistic

$$\Phi = \max_{(i,j,k)} a_k (|\phi_{ijk}| - b_k),$$

where a_k and b_k are scale-dependent constants and ϕ_{ijk} are weighted averages of the differences of standard normal random variables.

2. Compute a $(1 - \alpha)$ -quantile $q_{\text{Gauss}}(\alpha)$ of Φ by Monte Carlo simulations.
3. Adjust $q_{\text{Gauss}}(\alpha)$ by the scale-dependent constants

$$c_{\text{Gauss}}(\alpha, h_k) = b_k + q_{\text{Gauss}}(\alpha)/a_k$$

Test procedure

For the given significance level $\alpha \in (0, 1)$ and for each (i, j, k) , reject $H_0^{(ijk)}$ if $|\hat{\psi}_{ijk}| > c_{\text{Gauss}}(\alpha, h_k)$.

Theoretical properties

$\mathcal{C}1$ The functions λ_i are uniformly Lipschitz continuous:

$$|\lambda_i(u) - \lambda_i(v)| \leq L|u - v| \text{ for all } u, v \in [0, 1].$$

$\mathcal{C}1$ The functions λ_i are uniformly Lipschitz continuous:

$$|\lambda_i(u) - \lambda_i(v)| \leq L|u - v| \text{ for all } u, v \in [0, 1].$$

$\mathcal{C}2$ $0 < \lambda_{\min} \leq \lambda_i(w) \leq \lambda_{\max} < \infty$ for all $w \in [0, 1]$ and all i .

Assumptions

$\mathcal{C}1$ The functions λ_i are uniformly Lipschitz continuous:

$$|\lambda_i(u) - \lambda_i(v)| \leq L|u - v| \text{ for all } u, v \in [0, 1].$$

$\mathcal{C}2$ $0 < \lambda_{\min} \leq \lambda_i(w) \leq \lambda_{\max} < \infty$ for all $w \in [0, 1]$ and all i .

$\mathcal{C}3$ η_{it} are independent both across i and t .

Assumptions

$\mathcal{C}1$ The functions λ_i are uniformly Lipschitz continuous:

$$|\lambda_i(u) - \lambda_i(v)| \leq L|u - v| \text{ for all } u, v \in [0, 1].$$

$\mathcal{C}2$ $0 < \lambda_{\min} \leq \lambda_i(w) \leq \lambda_{\max} < \infty$ for all $w \in [0, 1]$ and all i .

$\mathcal{C}3$ η_{it} are independent both across i and t .

$\mathcal{C}4$ $\mathbb{E}[\eta_{it}] = 0$, $\mathbb{E}[\eta_{it}^2] = 1$ and $\mathbb{E}[|\eta_{it}|^\theta] \leq C_\theta < \infty$ for some $\theta > 4$.

Assumptions

$\mathcal{C}1$ The functions λ_i are uniformly Lipschitz continuous:

$$|\lambda_i(u) - \lambda_i(v)| \leq L|u - v| \text{ for all } u, v \in [0, 1].$$

$\mathcal{C}2$ $0 < \lambda_{\min} \leq \lambda_i(w) \leq \lambda_{\max} < \infty$ for all $w \in [0, 1]$ and all i .

$\mathcal{C}3$ η_{it} are independent both across i and t .

$\mathcal{C}4$ $\mathbb{E}[\eta_{it}] = 0$, $\mathbb{E}[\eta_{it}^2] = 1$ and $\mathbb{E}[|\eta_{it}|^\theta] \leq C_\theta < \infty$ for some $\theta > 4$.

$\mathcal{C}5$ $h_{\max} = o(1/\log T)$ and $h_{\min} \geq CT^{-b}$ for some $b \in (0, 1)$.

Assumptions

$\mathcal{C}1$ The functions λ_i are uniformly Lipschitz continuous:

$$|\lambda_i(u) - \lambda_i(v)| \leq L|u - v| \text{ for all } u, v \in [0, 1].$$

$\mathcal{C}2$ $0 < \lambda_{\min} \leq \lambda_i(w) \leq \lambda_{\max} < \infty$ for all $w \in [0, 1]$ and all i .

$\mathcal{C}3$ η_{it} are independent both across i and t .

$\mathcal{C}4$ $\mathbb{E}[\eta_{it}] = 0$, $\mathbb{E}[\eta_{it}^2] = 1$ and $\mathbb{E}[|\eta_{it}|^\theta] \leq C_\theta < \infty$ for some $\theta > 4$.

$\mathcal{C}5$ $h_{\max} = o(1/\log T)$ and $h_{\min} \geq CT^{-b}$ for some $b \in (0, 1)$.

$\mathcal{C}6$ $p := \{\#(i, j, k)\} = O(T^{(\theta/2)(1-b)-(1+\delta)})$ for some small $\delta > 0$.

Proposition

Let \mathcal{M}_0 be the set of triplets (i, j, k) for which $H_0^{(ijk)}$ holds true. Then under $\mathcal{C}1 - \mathcal{C}6$, it holds that

$$P\left(\forall (i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{\text{Gauss}}(\alpha, h_k)\right) \geq 1 - \alpha + o(1)$$

Theoretical properties

Proposition

Let \mathcal{M}_0 be the set of triplets (i, j, k) for which $H_0^{(ijk)}$ holds true. Then under C1 – C6, it holds that

$$\mathbb{P}\left(\forall (i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{\text{Gauss}}(\alpha, h_k)\right) \geq 1 - \alpha + o(1)$$

Proposition

Consider a sequence of functions $\lambda_i = \lambda_{i,T}$, $\lambda_j = \lambda_{j,T}$ such that

$$\exists \mathcal{I}_k : \lambda_i(w) - \lambda_j(w) \geq c_T \sqrt{\log T / (Th_k)} \quad \forall w \in \mathcal{I}_k, \quad (1)$$

and $c_T \rightarrow \infty$ faster than $\frac{\sqrt{\log T} \sqrt{\log \log T}}{\log \log \log T}$.

Theoretical properties

Proposition

Let \mathcal{M}_0 be the set of triplets (i, j, k) for which $H_0^{(ijk)}$ holds true. Then under $\mathcal{C}1 - \mathcal{C}6$, it holds that

$$\mathbb{P}\left(\forall (i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{\text{Gauss}}(\alpha, h_k)\right) \geq 1 - \alpha + o(1)$$

Proposition

Consider a sequence of functions $\lambda_i = \lambda_{i,T}$, $\lambda_j = \lambda_{j,T}$ such that

$$\exists \mathcal{I}_k : \lambda_i(w) - \lambda_j(w) \geq c_T \sqrt{\log T / (Th_k)} \quad \forall w \in \mathcal{I}_k, \quad (1)$$

and $c_T \rightarrow \infty$ faster than $\frac{\sqrt{\log T} \sqrt{\log \log T}}{\log \log \log T}$. Let \mathcal{M}_1 be the set of triplets (i, j, k) for which (1) holds true. Then under $\mathcal{C}1 - \mathcal{C}6$, it holds that

$$\mathbb{P}\left(\forall (i, j, k) \in \mathcal{M}_1 : |\hat{\psi}_{ijk}| > c_{\text{Gauss}}(\alpha, h_k)\right) = 1 - o(1)$$

Application

How to represent the results of the test? Plot the results of pairwise comparison $\mathcal{F}_{\text{reject}}(i, j)$.

How to represent the results of the test? Plot the results of pairwise comparison $\mathcal{F}_{\text{reject}}(i, j)$.

Minimal intervals

An interval $\mathcal{I}_k \in \mathcal{F}_{\text{reject}}(i, j)$ is called **minimal** if there is no other interval $\mathcal{I}_{k'} \in \mathcal{F}_{\text{reject}}(i, j)$ with $\mathcal{I}_{k'} \subset \mathcal{I}_k$. The set of minimal intervals is denoted $\mathcal{F}_{\text{reject}}^{\min}(i, j)$.

How to represent the results of the test? Plot the results of pairwise comparison $\mathcal{F}_{\text{reject}}(i, j)$.

Minimal intervals

An interval $\mathcal{I}_k \in \mathcal{F}_{\text{reject}}(i, j)$ is called **minimal** if there is no other interval $\mathcal{I}_{k'} \in \mathcal{F}_{\text{reject}}(i, j)$ with $\mathcal{I}_{k'} \subset \mathcal{I}_k$. The set of minimal intervals is denoted $\mathcal{F}_{\text{reject}}^{\min}(i, j)$.

We can make very similar confidence statement about the set of minimal intervals as well:

$$P\left(\forall (i, j, k) \in \mathcal{M}_0 : \mathcal{I}_k \notin \mathcal{F}_{\text{reject}}^{\min}(i, j)\right) \geq 1 - \alpha + o(1)$$

Application setting

- Five countries: Germany, Italy, Spain, France and UK; $T = 120$.

Application setting

- Five countries: Germany, Italy, Spain, France and UK; $T = 120$.
- $\alpha = 0.05$.


Application setting

- Five countries: Germany, Italy, Spain, France and UK; $T = 120$.
- $\alpha = 0.05$.
- Lengths of the time intervals 7, 14, 21, 28 days. The intervals start at days 1, 8, 15, ... and 4, 11, 19, ...


Application setting

- Five countries: Germany, Italy, Spain, France and UK; $T = 120$.
- $\alpha = 0.05$.
- Lengths of the time intervals 7, 14, 21, 28 days. The intervals start at days 1, 8, 15, ... and 4, 11, 19, ...



The family of intervals F




Application results







(b) minimal intervals produced by our test



(b) minimal intervals produced by our test



Application results, part 2



We can claim, with confidence of about 95%, that the null hypothesis is violated for all intervals (and all pairs of countries) for which our test rejects the null.

We can claim, with confidence of about 95%, that the null hypothesis is violated for all intervals (and all pairs of countries) for which our test rejects the null.

However, we can not say anything about the causes of such differences. This question requires further (probably not purely statistical) analysis.

We can claim, with confidence of about 95%, that the null hypothesis is violated for all intervals (and all pairs of countries) for which our test rejects the null.

However, we can not say anything about the causes of such differences. This question requires further (probably not purely statistical) analysis.

Further possible extensions:

- introduce scaling factor in the trend function, that allow for adjusting for the size of the country (population, density, testing regimes, etc.);

We can claim, with confidence of about 95%, that the null hypothesis is violated for all intervals (and all pairs of countries) for which our test rejects the null.

However, we can not say anything about the causes of such differences. This question requires further (probably not purely statistical) analysis.

Further possible extensions:

- introduce scaling factor in the trend function, that allow for adjusting for the size of the country (population, density, testing regimes, etc.);
- connect with data-driven techniques such as machine learning;

We can claim, with confidence of about 95%, that the null hypothesis is violated for all intervals (and all pairs of countries) for which our test rejects the null.

However, we can not say anything about the causes of such differences. This question requires further (probably not purely statistical) analysis.

Further possible extensions:

- introduce scaling factor in the trend function, that allow for adjusting for the size of the country (population, density, testing regimes, etc.);
- connect with data-driven techniques such as machine learning;
- cluster the countries based on the trends they exhibit.

Thank you!

Simulation results for the size of the test




Table 1: Size of the multiscale test

	$n = 5$			$n = 10$			$n = 50$		
	significance level α			significance level α			significance level α		
	0.01	0.05	0.1	0.01	0.05	0.1	0.01	0.05	0.1
$T = 100$	0.011	0.047	0.093	0.010	0.044	0.087	0.008	0.037	0.075
$T = 250$	0.009	0.047	0.091	0.009	0.046	0.087	0.008	0.035	0.069
$T = 500$	0.010	0.044	0.083	0.008	0.048	0.093	0.007	0.035	0.077

Simulation results for the power of the test




Table 2: Power of the multiscale test for scenario A

	$n = 5$			$n = 10$			$n = 50$		
	significance level α			significance level α			significance level α		
	0.01	0.05	0.1	0.01	0.05	0.1	0.01	0.05	0.1
$T = 100$	0.335	0.518	0.597	0.306	0.474	0.545	0.212	0.352	0.418
$T = 250$	0.615	0.790	0.836	0.580	0.764	0.800	0.470	0.648	0.705
$T = 500$	0.736	0.905	0.917	0.738	0.884	0.890	0.636	0.799	0.830

Simulation results for the power of the test





Table 3: Power of the multiscale test for scenario B

	$n = 5$			$n = 10$			$n = 50$		
	significance level α			significance level α			significance level α		
	0.01	0.05	0.1	0.01	0.05	0.1	0.01	0.05	0.1
$T = 100$	0.824	0.910	0.903	0.812	0.893	0.890	0.738	0.847	0.857
$T = 250$	0.991	0.972	0.941	0.991	0.960	0.920	0.991	0.965	0.933
$T = 500$	0.997	0.973	0.949	0.995	0.961	0.923	0.996	0.969	0.932

Estimator of σ^2

We estimate the overdispersion parameter σ^2 by

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n \hat{\sigma}_i^2 \text{ and } \hat{\sigma}_i^2 = \frac{\sum_{t=2}^T (X_{it} - X_{it-1})^2}{2 \sum_{t=1}^T X_{it}}$$

Estimator of σ^2

We estimate the overdispersion parameter σ^2 by

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n \hat{\sigma}_i^2 \text{ and } \hat{\sigma}_i^2 = \frac{\sum_{t=2}^T (X_{it} - X_{it-1})^2}{2 \sum_{t=1}^T X_{it}}$$

We assume that λ_i is Lipschitz continuous. Then

$$X_{it} - X_{it-1} = \sigma \sqrt{\lambda_i \left(\frac{t}{T} \right)} (\eta_{it} - \eta_{it-1}) + r_{it},$$

where $|r_{it}| \leq C(1 + |\eta_{it-1}|)/T$ with a sufficiently large C .

Estimator of σ^2

We estimate the overdispersion parameter σ^2 by

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n \hat{\sigma}_i^2 \text{ and } \hat{\sigma}_i^2 = \frac{\sum_{t=2}^T (X_{it} - X_{it-1})^2}{2 \sum_{t=1}^T X_{it}}$$

We assume that λ_i is Lipschitz continuous. Then

$$X_{it} - X_{it-1} = \sigma \sqrt{\lambda_i \left(\frac{t}{T} \right)} (\eta_{it} - \eta_{it-1}) + r_{it},$$

where $|r_{it}| \leq C(1 + |\eta_{it-1}|)/T$ with a sufficiently large C . Hence,

$$\frac{1}{T} \sum_{t=2}^T (X_{it} - X_{it-1})^2 = 2\sigma^2 \left\{ \frac{1}{T} \sum_{t=2}^T \lambda_i(t/T) \right\} + o_p(1)$$

Estimator of σ^2

We estimate the overdispersion parameter σ^2 by

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n \hat{\sigma}_i^2 \text{ and } \hat{\sigma}_i^2 = \frac{\sum_{t=2}^T (X_{it} - X_{it-1})^2}{2 \sum_{t=1}^T X_{it}}$$

We assume that λ_i is Lipschitz continuous. Then

$$X_{it} - X_{it-1} = \sigma \sqrt{\lambda_i \left(\frac{t}{T} \right)} (\eta_{it} - \eta_{it-1}) + r_{it},$$

where $|r_{it}| \leq C(1 + |\eta_{it-1}|)/T$ with a sufficiently large C . Hence,

$$\frac{1}{T} \sum_{t=2}^T (X_{it} - X_{it-1})^2 = 2\sigma^2 \left\{ \frac{1}{T} \sum_{t=2}^T \lambda_i(t/T) \right\} + o_p(1)$$

Together with

$$\frac{1}{T} \sum_{t=1}^T X_{it} = \frac{1}{T} \sum_{t=1}^T \lambda_i(t/T) + o_p(1),$$

we get that $\hat{\sigma}_i^2 = \sigma^2 + o_p(1)$ for any i and thus $\hat{\sigma}^2 = \sigma^2 + o_p(1)$.

[Go back](#)

Notation

In order to proceed with the proof, we will need the following notation:

$$\begin{aligned}\hat{\psi}_{ijk,T} &= \frac{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} - X_{jt})}{\hat{\sigma}\left\{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} + X_{jt})\right\}^{1/2}} \\ \hat{\psi}_{ijk,T}^0 &= \frac{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \sigma \bar{\lambda}_{ij}^{1/2}\left(\frac{t}{T}\right)(\eta_{it} - \eta_{jt})}{\hat{\sigma}\left\{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} + X_{jt})\right\}^{1/2}} & \hat{\Psi}_T &= \max_{(i,j,k)} a_k(|\hat{\psi}_{ijk,T}^0| - b_k) \\ \psi_{ijk,T}^0 &= \frac{1}{\sqrt{2Th_k}} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(\eta_{it} - \eta_{jt}) & \Psi_T &= \max_{(i,j,k)} a_k(|\psi_{ijk,T}^0| - b_k) \\ \phi_{ijk,T} &= \frac{1}{\sqrt{2Th_k}} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(Z_{it} - Z_{jt}) & \Phi_T &= \max_{(i,j,k)} a_k(|\phi_{ijk,T}| - b_k)\end{aligned}$$

Strategy of the proof

1. We prove that $|\hat{\psi}_T^0 - \psi_T| = o_p(r_T)$, where $\{r_T\}$ is some null sequence.

Strategy of the proof

1. We prove that $|\hat{\Psi}_T^0 - \Psi_T| = o_p(r_T)$, where $\{r_T\}$ is some null sequence.
2. With the help of results from Chernozhukov et al. (2017), we prove

$$\sup_{q \in \mathbb{R}} \left| P(\Psi_T \leq q) - P(\Phi_T \leq q) \right| = o(1)$$

Strategy of the proof

1. We prove that $|\hat{\Psi}_T^0 - \Psi_T| = o_p(r_T)$, where $\{r_T\}$ is some null sequence.
2. With the help of results from Chernozhukov et al. (2017), we prove

$$\sup_{q \in \mathbb{R}} \left| P(\Psi_T \leq q) - P(\Phi_T \leq q) \right| = o(1)$$

3. By using these two results, we now show that

$$\sup_{q \in \mathbb{R}} \left| P(\hat{\Psi}_T^0 \leq q) - P(\Phi_T \leq q) \right| = o(1) \quad (2)$$

Strategy of the proof

1. We prove that $|\hat{\Psi}_T^0 - \Psi_T| = o_p(r_T)$, where $\{r_T\}$ is some null sequence.
2. With the help of results from Chernozhukov et al. (2017), we prove

$$\sup_{q \in \mathbb{R}} \left| P(\Psi_T \leq q) - P(\Phi_T \leq q) \right| = o(1)$$

3. By using these two results, we now show that

$$\sup_{q \in \mathbb{R}} \left| P(\hat{\Psi}_T^0 \leq q) - P(\Phi_T \leq q) \right| = o(1) \quad (2)$$

4. It can be shown that $P(\Phi_T \leq q_{T,\text{Gauss}}(\alpha)) = 1 - \alpha$. From this and (2), it immediately follows that

$$P(\hat{\Psi}_T^0 \leq q_{T,\text{Gauss}}(\alpha)) = 1 - \alpha + o(1),$$

which in turn implies the desired statement.

Idea behind a_k and b_k

A more modern approach of constructing the individual critical values $c_{ijk}(\alpha)$ (Dümbgen and Spokoiny (2001)): let them depend on the scale of the testing problem, i.e. the length h_k of the time interval.

Idea behind a_k and b_k

A more modern approach of constructing the individual critical values $c_{ijk}(\alpha)$ (Dümbgen and Spokoiny (2001)): let them depend on the scale of the testing problem, i.e. the length h_k of the time interval. Specifically,

$$c_{ijk}(\alpha) = c(\alpha, h_k) := b_k + q(\alpha)/a_k,$$

where $a_k = \{\log(e/h_k)\}^{1/2} / \log \log(e^e/h_k)$ and $b_k = \sqrt{2 \log(1/h_k)}$ are scale-dependent constants and $q_T(\alpha)$ is chosen such that we control FWER. [Go back](#)

Idea behind the additive correction

Consider the uncorrected Gaussian statistic

$$\Phi^{\text{uncor}} = \max_{(i,j,k)} |\phi_{ijk}|$$

Idea behind the additive correction

Consider the uncorrected Gaussian statistic

$$\Phi^{\text{uncor}} = \max_{(i,j,k)} |\phi_{ijk}|$$

and let the family of intervals be

$$\mathcal{F} = \{[(m-1)h_l, mh_l] \text{ for } 1 \leq m \leq 1/h_l, 1 \leq l \leq L\}$$

Idea behind the additive correction

Consider the uncorrected Gaussian statistic

$$\Phi^{\text{uncor}} = \max_{(i,j,k)} |\phi_{ijk}|$$

and let the family of intervals be

$$\mathcal{F} = \{[(m-1)h_l, mh_l] \text{ for } 1 \leq m \leq 1/h_l, 1 \leq l \leq L\}$$

Then we can rewrite the uncorrected test statistic as

$$\Phi^{\text{uncor}} = \max_{i,j} \max_{\substack{1 \leq l \leq L, \\ 1 \leq m \leq 1/h_l}} \left| \frac{1}{\sqrt{2Th_l}} \sum_{t=1}^T 1\left(\frac{t}{T} \in [(m-1)h_l, mh_l]\right) (Z_{it} - Z_{jt}) \right|$$

Idea behind the additive correction

Consider the uncorrected Gaussian statistic

$$\Phi^{\text{uncor}} = \max_{(i,j,k)} |\phi_{ijk}|$$

and let the family of intervals be

$$\mathcal{F} = \{[(m-1)h_l, mh_l] \text{ for } 1 \leq m \leq 1/h_l, 1 \leq l \leq L\}$$

Then we can rewrite the uncorrected test statistic as

$$\Phi^{\text{uncor}} = \max_{i,j} \max_{\substack{1 \leq l \leq L, \\ 1 \leq m \leq 1/h_l}} \left| \frac{1}{\sqrt{2Th_l}} \sum_{t=1}^T 1\left(\frac{t}{T} \in [(m-1)h_l, mh_l]\right) (Z_{it} - Z_{jt}) \right|$$

$\Rightarrow \max_m \dots = \sqrt{2 \log(1/h_l)} + o_P(1) \rightarrow \infty$ as $h \rightarrow 0$ and the stochastic behavior of Φ^{uncor} is dominated by the elements with small bandwidths h_l . [Go back](#)