

Snail ID Guide

Jonathan Farr, Gabrielle Languedoc, Lucy Wall, Meredith Miller

10/18/2021

***Calliostoma ligatum* (Blue top snail)**

Description

Calliostoma ligatum's shells are cone-shaped with three distinct convex whorls, and are generally 2.5 to 3 cm in diameter and 1.7 to 3 cm tall (Tuskes 2019; Crowles 2004; Meschkat et al., 2013). The colour of their unbeaded spiral ridges generally alternates between light golden or tan and chocolate brown, although spirals can be white, grey, black or red. If the shell is worn or damaged a distinctive blue inner layer is visible, which can also be seen in the inner edge of empty shells. The shell has a closed umbilicus, and its opening is rounded with a pearly interior. The snail's foot has a distinctive bright orange sole.

Calliostoma ligatum is present from central Alaska to southern California (Metschkat et al., 2013; Harbo 2011). It can be found intertidally but also at depths of up to 30 meters in eel grass and kelp beds, and rocky areas. Blue top snails are omnivorous, and although they mainly consume *Macrocystis* and other brown algae they also eat diatoms, ascidians, sponges, bryozoans, and detritus (Crowles 2004; Meschkat et al., 2013). Their main predators are sea stars and sunflower stars, but, interestingly, they do not demonstrate antipredator behaviour towards predatory snails (Hoffman, 1980). As broadcast spawners, male *Calliostoma ligatum* release sperm into the water column, while eggs are released by females sheathed in mucous strands that await fertilization (Holyvoak 1988). After fertilization, they undergo several changes over 12 days to become planktonic larvae, veligers, and then juvenile snails. They reach sexual maturity after 1 year. Blue top snails may be confused for Purple-ringed top snails (*Calliostoma annulatum*), but the beaded spirals on Purple-ringed top snails are easily differentiated from the smooth spirals of Blue top snails. Other snails in the area either do not have a similar top-like shell morphology, or lack the distinguishable spirals and orange foot of the top snails.

Identification Questions

- 1) Is the shell of the snail top shaped (not cone shaped) with three convex whorls?
- 2) Does the snail have a bright orange foot?
- 3) Are the spiral ridges on the snail's shell smoothed (not beaded)?

Figures




Figure 1: *Calliostoma ligatum* positioned in ventral (A) lateral (B) views with important identifying characteristics labelled.



Figure 2: *Calliostoma ligatum* shells (containing Hermit crabs) collected at Scott's Bay.

***Nucella lamellosa* (Frilled/Wrinkled Dogwinkle)**

Description

Nucella lamellosa's shell can have very variable characteristics, such as color and texture. Typically in solid colours, they can be light brown, grey, to white, but can also have banding patterns (Harbo, 2011). Some have even been observed in orange and purple colouring with coloured bands. Their 'frilled' name comes from their lamella, which look like wrinkles or ruffles in the whelk's shell. A shell can have up to 12 lamellae (Proudfoot & Fretwell, 2015). Presence or absence of frills in this species has been observed to vary with both wave and predator (*Cancer productus*) exposure. Typically, wave exposed snails have smooth shells. Their shells are not 'polished' looking and are spirally coiled with 5-7 whorls (Bering et al. 2017). Overall, shell shape is elongated into a point and can reach up to 80mm in height (Proudfoot & Fretwell, 2015). The shell mouth opening is ovate and the lip of the shell is smooth and rounded with white coloring (or outside shell colour) on the inside (Bering et al. 2017). Its operculum is strongly spiraled and usually big enough to fully close the snail's shell mouth opening, and it has a closed umbilicus (Bering et al. 2017). Color, texture, thickness, coloured banding, and sometimes shape can vary widely for this species (Proudfoot & Fretwell, 2015). As there are 4 *Nucella* species in BC, this snail is commonly mistaken for its relatives. Greyish or white *N. lamellosa* can look a lot like *Nucella canaliculata*, however *N. canaliculata* is typically more streamlined in shape and doesn't have any frills on its shell (Proudfoot & Fretwell, 2015). *N. lamellosa*'s range extends from Alaska (Aleutian Islands) to California (central) (Proudfoot & Fretwell, 2015; Harbo, 2011). This range suggests Frilled Dogwinkles can live in a breadth of conditions, but distributions suggest preference for cold temperate waters (Bering et al. 2017). In terms of habitat preferences, these snails inhabit the rocky intertidal, specifically mid to low zones but can be in shallow subtidal locations as well (Proudfoot & Fretwell, 2015; Harbo, 2011). The Frilled Dogwinkle inhabits rocky crevices, rock faces, as well as barnacle and mussel beds. Frilled Dogwinkles are predatory and feed on mussels and acorn barnacles (among other mollusks) by drilling into them with their radula and using a siphon that penetrates the prey's shell to feed on internal tissues (Carefoot, 2021). Mating occurs in winter and spring. Sexually mature (>4yrs old) snails aggregate to breed with a group at their original hatching site (Bering et al. 2017). Females spawn eggs after 20 months and baby snails hatch from them after 140 days. Eggs are contained in capsules that protect them from factors such as salinity stress. Eggs ("sea oats") are pale yellow and shaped like ~10mm vases, they can be observed on rocks in clusters (Bering et al. 2017).

Questions

- 1) Does the snail have 5-7 whorls, with the last whorl being the largest by far?
- 2) Does it have an oval aperture that is around half the length of the shell?
- 3) Is the lip thick, rounded, and smooth, with white or the shell's outer color showing through?

Figures



Figure 3: A drawn diagram of **Nucella lamellosa** (L.Wall). These drawings show the morphological variety in shell colours, textures, and banding. 1) shows underside of the snail, highlighting the oval aperture. 2) shows an upright view of the snail, showing frills, five distinct whorls, and brown colouring. 3) shows a banding colour pattern and less frilly surface.



Figure 4: Photos showing two varieties of **N. lamellosa** snails collected from Scotts Bay in Bamfield, BC on October 15th 2021 (L.Wall). This highlights the variation in snail color and size, with smooth shell morphology suggesting plastic suitability to their wave-exposed habitat.

***Tegula funebralis* (Black Tegula/Black Turban Snail)**

Description

The *Tegula funebralis* has a thick and strong cone-shaped shell with a closed umbilicus (Fretwell & Starzomski, 2013). The rounded shell consists of four whorls, and it can grow up to three centimeters in diameter (Harbo, 2011). When wet, the shell appears non-shiny and black, but when dry can have a more dark purple or grey tone (Fretwell & Starzomski, 2013). The tip of the shell gets worn down over time, revealing a pearly white surface. The white interior contains a black-bodied snail, and the bottom of the foot appears tan. Males have a paler foot than females. Other similar species include the brown turban snail (*Chlorostoma brunnea*), and the dusky turban snail (*Tegula pulligo*). Both of these species have a lighter brown shell than the black turban snail, and *C. brunnea* is not found north of Oregon state.

The black turban snail is most commonly found along exposed or semi-protected rocky shorelines (Fretwell & Starzomski, 2013). It ranges along the west coast of North America from northern Vancouver Island down to central Mexico. These snails can be found along the entire intertidal, but mainly live in the mid/low zones. Black turban snails live in large groups, usually aggregated under rocks. If shells are found isolated from other individuals, there is most likely a hermit crab or other invertebrate living in its shell. These grazing herbivores will eat almost any common algae (Hiebert et al., 1979). They are also the prey to many intertidal invertebrates, including carnivorous snails, crabs, or sea stars. Humans are also a major predator of black turban snails, and there is evidence of human collection of these snails as far back as 12,000 years ago (Erlandson et al., 2015). These snails reach sexual maturity at about fourteen millimeters in diameter, and female snails will lay several hundred eggs approximately once per year (Hiebert et al., 1979). Juveniles live under small rocks or in the sand, then as young adults live in the upper intertidal for five to six years before migrating down to the mid/low intertidal.

Identification Questions

- 1) Is the snail surrounded by other similar individuals?
- 2) Does the snail have four whorls/spirals on its shell?
- 3) If the snail is wet, does it look black?

Figures




Figure 5: A drawn diagram of **Tegula funebralis**. The left drawing shows the underside of the snail, highlighting the aperture. The right drawing shows an upright view of the snail, including the location of the foot, highlighting the worn tip of the shell.



Figure 6: Photos showing three varieties of **Tegula funebralis** snails collected from Scotts Bay in Bamfield, BC on October 15th 2021. This highlights the similarities, such as the variations in tip morphology caused by wear and tear from rough intertidal conditions.

***Nucella ostrina* (Northern Striped Dogwinkle)**

Description

This whelk is distinguished by its alternating lines of thick and thin bands on its shell. Additionally, it can be identified by the three indistinct whorls on its shell. Colors can range from dark browns, blacks, and browns to light greys and oranges always with a purple interior (Meschkat et al., 2014). This snail typically lives in the mid to high intertidal zones of exposed rocky shorelines from the Bering Sea to Southern California. The end of its shell known as the umbilicus is closed. It is most commonly found along barnacle and mussel beds as this is their preferred diet. This species feeds by drilling a hole with its radula mouthpart into its prey and extracting digestive enzymes which are then extracted by its long proboscis (*E-Fauna BC Atlas Page*, 2021). *N. ostrina* is commonly mistaken for the northern channeled dog winkle (*Nucella canaliculata*) as well as the proposed northern lined dog winkle (*Nucella analoga*). Both of these species can be distinguished from *N. ostrina* because of their slender distinctive whorls and smaller body sizes. Mating occurs in Winter and Spring months with an annual spawn. To reproduce, females deposit leathery embryonic capsules attached by stalks to the rock from which the eggs develop and hatch as juveniles after about 2.5-4 months (Lloyd & Gosselin, 2007). Therefore, there is no larval feeding stage and maternal care is critical to the survival of the offspring. Each capsule will produce about 20 juveniles.

Identification Questions

- 1) Does this snail have a banding pattern that alternates between thick and thin stripes?
- 2) Does this snail have three indistinct whorls?
- 3) Is the inside of the shell purple?

Figures




Figure 7: Showing **Nucella ostrina** distinctive features such as closed umbilicus, indistinct whorls, purple interior shell and operculum.



Figure 8: Showing **Nucella ostrina** collected from Scotts Bay of Bamfield, BC, Canada in October of 2021.

Supplemental Information

Table 1: Shell morphometric measurements of four intertidal snails

Species	Length (mm)	Height (mm)
Nucella_ostrina	20.00	10.00
Nucella_lamellosa	16.00	6.30
Nucella_lamellosa	19.00	9.00
Calliostoma_ligatum	5.00	0.45
Calliostoma_ligatum	10.00	12.00
Tegula_funebralis	13.10	16.80
Tegula_funebralis	17.15	19.25
Tegula_funebralis	18.65	19.45

Figures




Figure 9: Snail shell length and height for four intertidal snails. Snails were collected from Scott's Bay, and measured in lab using calipers

Species	Size	Morphology	Trophic Role and Diet	Reproductive Mode
<i>Nucella ostrina</i>	10 mm in width and 20 mm in length	3 whorls, alternating lines of thick and thin bands. Colour range: black, browns, light grey, orange, always with a purple interior	Preys on barnacles and mussels. Main predator is <i>Pisaster ochraceus</i>	Spawns annually with mating occurring typically in Winter and Spring. Females deposit leathery embryonic capsules from which the eggs develop and hatch as juveniles.
<i>Nucella lamellosa</i>	Up to 50-80mm full grown in length	5-7 whorls, non-polished shells with elongated point Colour range: (highly varied) solid colours, light brown, grey, white, orange, banding patterns absent or present, inside is white (or colour shows through)	Carnivore, eats molluscs including barnacles, drills with radula and eats using siphon Main predators: crabs, mainly Cancer	Seasonal breeding site-return to hatching grounds Communal spawning ~20 month gestation 'Sea oats' cases hold eggs ~140 days between spawning and hatching of juveniles
<i>Tegula funebralis</i>	Can grow up to 3cm in diameter.	4 whorls, closed umbilicus Colour: black with worn down tips appearing pearly white, can appear dark grey if dried out, inside is pearly white	Herbivorous, eats diatoms and other common algae. Main predators: intertidal invertebrates, carnivorous snails, humans.	Reach sexual maturity at a diameter of ~14mm. The female snails lay several hundred eggs once per year. Small adults live together in the upper intertidal for 5-6 years before migrating towards the middle and lower.
<i>Calliostoma ligatum</i>	Up to 2.5 - 3 cm in diameter when fully grown	3 whorls, cone-shaped, range from 2.5-3cm in diameter, closed umbilicus Colour: (highly variable) Smooth spiral ridges generally alternate between light gold/tan and chocolate brown, spirals can be white, grey, black or red, worn shell appears blue, pearly interior	Omnivorous, eats kelp and small animals (bryozoans & sponges) Main predators: starfish	Broadcast spawners Rapid 12 day metamorphosis from larvae → juvenile snails → adult Reach sexual maturity after 1 year

Figure 10: Table 2. Summary of characteristics for four intertidal snails.

Bibliography

- Bering, N., Hext, T., & Parker, E. (2017). *Nucella lamellosa*. In: Oregon Estuarine Invertebrates: Rudys' Illustrated Guide to Common Species, 3rd ed. University of Oregon Libraries and Oregon Institute of Marine Biology, Charleston, OR. https://scholarsbank.uoregon.edu/xmlui/bitstream/handle/1794/12913/N_lamellosa_2017_final.pdf?sequence=3
- Carefoot, T. (2021) *Nucella lamellosa* (Gmelin, 1791) Dog Whelk; Frilled Dogwinkle; Wrinkled Dog Wheld. E-Fauna BC: Electronic Atlas of the Wildlife of British Columbia. <http://linnet.geog.ubc.ca/efauna/Atlas/Atlas.aspx?sciname=Nucella%20lamellosa&lifeform=184>
- Crowles, D. (2004). *Calliostoma ligatum*. Invertebrates of the Salish Sea. Walla Walla University. https://inverts.wallawalla.edu/Mollusca/Gastropoda/Prosobranchia/Order_Archaegastropoda/Suborder_Trochidae/Calliostoma_ligatum.html
- E-Fauna BC Atlas Page. (2021). *E-Fauna BC. Nucella ostrina* (Gould, 1852) Dog Whelk; Dog Winkle. <http://linnet.geog.ubc.ca/efauna/Atlas/Atlas.aspx?sciname=Nucella%20ostrina>
- Erlandson, J. M., Ainis, A. F., Braje, T. J., Jew, N. P., McVey, M., Rick, T. C., Vellanoweth, R. L., & Watts, J. (2015). 12,000 Years of Human Predation on Black Turban Snails (*Chlorostoma funebralis*) on Alta California's Northern Channel Islands. *California Archaeology*, 7(1), 59–91. <https://doi.org/10.1179/1947461X15Z.00000000056>
- Fretwell, K., & Starzomski, B. (2013). *Black turban snail, Tegula funebralis*. *Biodiversity of the Central Coast*. <https://www.centralcoastbiodiversity.org/black-turban-snail-bull-tegula-funebralis.html>
- Harbo, R. M. (2011). Whelks to Whales: Coastal Marine Life of the Pacific Northwest (Revised Second ed.). Harbour Publishing. Turban Snails, Topsnails, 131.
- Harbo, R. M. (2011). Whelks to Whales: Coastal Marine Life of the Pacific Northwest (Revised Second ed.). Harbour Publishing, BC. Whelks, Dogwinkles: 144-145.
- Hiebert, T. C., Butler, B. A., & Shanks, A. L. (1979). *Tegula funebralis* in Oregon Estuarine Invertebrates: Rudys' Illustrated Guide to 140 Common Species (No. 3). University of Oregon Libraries and Oregon Institute of Marine Biology. https://oimb.uoregon.edu/wp-content/uploads/2019/03/T_funebralis_2018.pdf
- Hoffman, D. L. (1980). Defensive Responses of Marine Gastropods (Prosobranchia, Trochidae) to Certain Predatory Seastars and the Dire Whelk, *Searlesia dira* (Reeve). *Pacific Science*, 34(3), 233-243.
- Holyoak, A.R. (1988). Spawning and larval development of the trochid gastropod *Calliostoma ligatum*. *The Veliger*, 30(4), 369-371
- Lloyd, M. J., & Gosselin, L. A. (2007). Role of Maternal Provisioning in Controlling Interpopulation Variation in Hatching Size in the Marine Snail *Nucella ostrina*. *The Biological Bulletin*, 213(3), 316–324. <https://doi.org/10.2307/25066649>
- Meschkat, C., Fretwell, K., & Brian Starzomski, B. (2014). *Northern striped dogwinkle • Nucella ostrina*. *Biodiversity of the Central Coast*. <https://www.centralcoastbiodiversity.org/northern-striped-dogwinkle-bull-nucella-ostrina.html>
- Meschkat, C., Fretwell, K., & Starzomski, B. (2013). *Blue topsnail, Western ridged top shell. Calliostoma ligatum*. *Biodiversity of the Central Coast*. <https://www.centralcoastbiodiversity.org/blue-topsnail-bull-calliostoma-ligatum.html>
- Proudfoot, B., & Fretwell, K. (2015). *Frilled Dogwinkle, Wrinkled Dogwinkle • Nucella Lamellosa*. *Biodiversity of the Central Coast*. <https://www.centralcoastbiodiversity.org/frilled-dogwinkle-bull-nucella-lamellosa.html>.
- Tuskes, P. M. (2019). Calliostomatidae of the northeast Pacific. *Zoosymposia*, 13, 83-96. <http://dx.doi.org/10.11646/zootaxa.13.1.9>