GABA-aminotransferase inactivators

Scheda H02_gabaat_inactivator_myelin_exposure · C3 evidenza contraria da chiarire, danno o nullo da fuori WWOX, F3 ipotesi favorevole dal bersaglio, non provata in WWOX, P3 premessa mancante, nessun elemento favorevole registrato

Data dei dati: 2026-10-28 · **Costruzione del catalogo:** 2026-10-28T23:40:16Z (due date diverse: la prima è l'ultimo aggiornamento dei dati, la seconda la costruzione del catalogo dei composti).

1. Domanda (il bisogno)	Controllo inibitorio; Controllo di crisi e spasmi come bisogno sintomatico; Ipomielinizzazione
2. Candidato / intervento	**GABA-aminotransferase inactivators (vigabatrin; sibling CPP-115): myelin exposure limit in a WWOX-deficient brain with white-matter loss (hypomyelination unresolved in …** (Beneficio parziale utile; Track A (indipendente dalla variante); controversa)
3. Risultato WWOX	new symmetrical MRI signal change in the globus pallidi and thalami during vigabatrin (VABAM): increase – A report of two children selected for having the finding: the number of WWOX children given vigabatrin, how many had follow-up MRI, the doses and … [human]
4. Precedente esterno	myelination on histology after exposure from postnatal day 4 (hypomyelination and gliopathy noted from days 4 to 15): decrease – Vigabatrin only. A lesser degree of hypomyelination persisted at longer exposure windows; vacuolation was attenuated after the recovery period.
5. Controevidenza decisiva	Two children selected for having the finding, with no denominator, cannot be shown to exceed (or fall below) the label's figures: 22 percent MRI changes in vigabatrin-treated non-WWOX infants with infantile spasms against 4 percent on other therapies (retrospective, N=205, arm sizes not stated). (altre voci nel dossier completo: 14) (la prima voce che la scheda o il bisogno elencano; non è un giudizio di peso)
6. Dato mancante	How many WWOX-DEE children have received vigabatrin and how many of them had follow-up MRI (the denominator behind the two reported cases).
7. Prossimo passo	Rodent: a Wwox-deficient juvenile model that survives the exposure window (the neuron-specific deletion is the candidate; survival is to be checked first) against control littermates, vigabatrin against vehicle at an exposure window and level taken from the published rat studies, with confirmed brain exposure, blinded electron microscopy for intramyelinic splits, myelin …
8. Fonti essenziali (al massimo 5)	• PMID 41442931 https://pubmed.ncbi.nlm.nih.gov/41442931/ Vigabatrin-Associated Brain Magnetic Resonance Imaging Abnormalities in Two Children With WW domain-containing oxidoreductase-Related Epileptic Encephalopathy Syndrome. [Pediatr Neurol, 2026] • PMID 34634460 https://pubmed.ncbi.nlm.nih.gov/34634460/ Altered neocortical oscillations and cellular excitability in an in vitro Wwox knockout mouse model of epileptic encephalopathy. [Neurobiol Dis, 2021] • PMID 30290271 https://pubmed.ncbi.nlm.nih.gov/30290271/ Wwox deletion leads to reduced GABA-ergic inhibitory interneuron numbers and activation of microglia and astrocytes in mouse hippocampus. [Neurobiol Dis, 2019] • PMID 34268881 https://pubmed.ncbi.nlm.nih.gov/34268881/ Modeling genetic epileptic encephalopathies using brain organoids. [EMBO Mol Med, 2021] • PMID 36828035 https://pubmed.ncbi.nlm.nih.gov/36828035/ WWOX P47T partial loss-of-function mutation induces epilepsy, progressive neuroinflammation, and cerebellar degeneration in mice phenocopying human SCAR12. [Prog Neurobiol, 2023]

Dossier completo (3-4 pagine, appendice): #export=H02_gabaat_inactivator_myelin_exposure. Materiale di ricerca per la discussione con il team clinico curante: non è consiglio medico e non è una graduatoria di trattamenti. Associazione non è modulazione, e modulazione non è recupero. Generato da un registro del repository (non incluso in questo pacchetto) (schema legend.wwox.repurposing_view/2, shareable). Versione condivisibile: vedi il piè di pagina della pagina per le fonti incluse.