

Class 14

Marina Puffer (PID: A16341339)

Section 1. Differential Expression Analysis

```
library(DESeq2)
```

Loading required package: S4Vectors

Loading required package: stats4

Loading required package: BiocGenerics

Attaching package: 'BiocGenerics'

The following objects are masked from 'package:stats':

IQR, mad, sd, var, xtabs

The following objects are masked from 'package:base':

anyDuplicated, aperm, append, as.data.frame, basename, cbind,
colnames, dirname, do.call, duplicated, eval, evalq, Filter, Find,
get, grep, grepl, intersect, is.unsorted, lapply, Map, mapply,
match, mget, order, paste, pmax, pmax.int, pmin, pmin.int,
Position, rank, rbind, Reduce, rownames, sapply, setdiff, sort,
table, tapply, union, unique, unsplit, which.max, which.min

Attaching package: 'S4Vectors'

The following object is masked from 'package:utils':

findMatches

The following objects are masked from 'package:base':

expand.grid, I, unname

Loading required package: IRanges

Loading required package: GenomicRanges

Loading required package: GenomeInfoDb

Loading required package: SummarizedExperiment

Loading required package: MatrixGenerics

Loading required package: matrixStats

Attaching package: 'MatrixGenerics'

The following objects are masked from 'package:matrixStats':

colAlls, colAnyNAs, colAnys, colAvgsPerRowSet, colCollapse,
colCounts, colCummaxs, colCummins, colCumprods, colCumsums,
colDiffs, colIQRDiffs, colIQRs, colLogSumExps, colMadDiffs,
colMads, colMaxs, colMeans2, colMedians, colMins, colOrderStats,
colProds, colQuantiles, colRanges, colRanks, colSdDiffs, colSds,
colSums2, colTabulates, colVarDiffs, colVars, colWeightedMads,
colWeightedMeans, colWeightedMedians, colWeightedSds,
colWeightedVars, rowAlls, rowAnyNAs, rowAnys, rowAvgsPerColSet,
rowCollapse, rowCounts, rowCummaxs, rowCummins, rowCumprods,
rowCumsums, rowDiffs, rowIQRDiffs, rowIQRs, rowLogSumExps,
rowMadDiffs, rowMads, rowMaxs, rowMeans2, rowMedians, rowMins,
rowOrderStats, rowProds, rowQuantiles, rowRanges, rowRanks,
rowSdDiffs, rowSds, rowSums2, rowTabulates, rowVarDiffs, rowVars,
rowWeightedMads, rowWeightedMeans, rowWeightedMedians,
rowWeightedSds, rowWeightedVars

Loading required package: Biobase

Welcome to Bioconductor

Vignettes contain introductory material; view with
'browseVignettes()'. To cite Bioconductor, see
'citation("Biobase")', and for packages 'citation("pkgname")'.

Attaching package: 'Biobase'

The following object is masked from 'package:MatrixGenerics':

rowMedians

The following objects are masked from 'package:matrixStats':

anyMissing, rowMedians

Data import

```
metaFile <- "GSE37704_metadata.csv"
countFile <- "GSE37704_featurecounts.csv"
```

```
# Import metadata and take a peak
colData = read.csv(metaFile, row.names=1)
head(colData)
```

```
              condition
SRR493366 control_sirna
SRR493367 control_sirna
SRR493368 control_sirna
SRR493369      hoxa1_kd
SRR493370      hoxa1_kd
SRR493371      hoxa1_kd
```

```
# Import countdata
countData = read.csv(countFile, row.names=1)
head(countData)
```

	length	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370
ENSG00000186092	918	0	0	0	0	0
ENSG00000279928	718	0	0	0	0	0
ENSG00000279457	1982	23	28	29	29	28
ENSG00000278566	939	0	0	0	0	0
ENSG00000273547	939	0	0	0	0	0
ENSG00000187634	3214	124	123	205	207	212

	SRR493371
ENSG00000186092	0
ENSG00000279928	0
ENSG00000279457	46
ENSG00000278566	0
ENSG00000273547	0
ENSG00000187634	258

Q. Complete the code below to remove the troublesome first column from countData

```
# Note we need to remove the odd first $length col
countData <- as.matrix(countData[,-1])
head(countData)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000186092	0	0	0	0	0	0
ENSG00000279928	0	0	0	0	0	0
ENSG00000279457	23	28	29	29	28	46
ENSG00000278566	0	0	0	0	0	0
ENSG00000273547	0	0	0	0	0	0
ENSG00000187634	124	123	205	207	212	258

Q. Complete the code below to filter countData to exclude genes (i.e. rows) where we have 0 read count across all samples (i.e. columns).

```
# Filter count data where you have 0 read count across all samples.
rowsums <- rowSums(countData)
zero <- rowsums == 0
countData = countData[!zero, ]
head(countData)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000279457	23	28	29	29	28	46
ENSG00000187634	124	123	205	207	212	258
ENSG00000188976	1637	1831	2383	1226	1326	1504
ENSG00000187961	120	153	180	236	255	357
ENSG00000187583	24	48	65	44	48	64
ENSG00000187642	4	9	16	14	16	16

How many genes left?

```
nrow(countData)
```

```
[1] 15975
```

Running DESeq2

```
dds = DESeqDataSetFromMatrix(countData=countData,
                              colData=colData,
                              design=~condition)
```

Warning in DESeqDataSet(se, design = design, ignoreRank): some variables in design formula are characters, converting to factors

```
dds = DESeq(dds)
```

estimating size factors

estimating dispersions

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing

```
dds
```

```
class: DESeqDataSet
dim: 15975 6
metadata(1): version
assays(4): counts mu H cooks
rownames(15975): ENSG00000279457 ENSG00000187634 ... ENSG00000276345
               ENSG00000271254
rowData names(22): baseMean baseVar ... deviance maxCooks
colnames(6): SRR493366 SRR493367 ... SRR493370 SRR493371
colData names(2): condition sizeFactor
```

```
res = results(dds, contrast=c("condition", "hoxa1_kd", "control_sirna"))
```


Q. Call the `summary()` function on your results to get a sense of how many genes are up or down-regulated at the default 0.1 p-value cutoff.

```
summary(res)
```

```
out of 15975 with nonzero total read count
adjusted p-value < 0.1
LFC > 0 (up)      : 4349, 27%
LFC < 0 (down)    : 4396, 28%
outliers [1]      : 0, 0%
low counts [2]    : 1237, 7.7%
(mean count < 0)
[1] see 'cooksCutoff' argument of ?results
[2] see 'independentFiltering' argument of ?results
```

Volcano plot

```
plot(res$log2FoldChange, -log(res$padj))
```




Q. Improve this plot by completing the below code, which adds color and axis labels

```
# Make a color vector for all genes
mycols <- rep("gray", nrow(res) )

# Color red the genes with absolute fold change above 2
mycols[ abs(res$log2FoldChange) > 2 ] <- "red"

# Color blue those with adjusted p-value less than 0.01
# and absolute fold change more than 2
inds <- (abs(res$padj)<0.1) & (abs(res$log2FoldChange) > 2 )
mycols[ inds ] <- "blue"

plot( res$log2FoldChange, -log(res$padj), col=mycols, xlab="Log2(FoldChange)", ylab="-Log(
```



Adding gene annotation

Q. Use the `mapIDs()` function multiple times to add **SYMBOL, **ENTREZID** and **GENENAME** annotation to our results by completing the code below.**

```
library("AnnotationDbi")
library("org.Hs.eg.db")
```

```
columns(org.Hs.eg.db)
```

[1] "ACCNUM"	"ALIAS"	"ENSEMBL"	"ENSEMBLPROT"	"ENSEMBLTRANS"
[6] "ENTREZID"	"ENZYME"	"EVIDENCE"	"EVIDENCEALL"	"GENENAME"
[11] "GENETYPE"	"GO"	"GOALL"	"IPI"	"MAP"
[16] "OMIM"	"ONTOLOGY"	"ONTOLOGYALL"	"PATH"	"PFAM"
[21] "PMID"	"PROSITE"	"REFSEQ"	"SYMBOL"	"UCSCKG"
[26] "UNIPROT"				


```
res$symbol = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="SYMBOL",
                    multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$entrez = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="ENTREZID",
                    multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$name = mapIds(org.Hs.eg.db,
                  keys=row.names(res),
                  keytype="ENSEMBL",
                  column="GENENAME",
                  multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
head(res, 10)
```

log2 fold change (MLE): condition hoxa1_kd vs control_sirna

Wald test p-value: condition hoxa1 kd vs control sirna

DataFrame with 10 rows and 9 columns

	baseMean	log2FoldChange	lfcSE	stat	pvalue
	<numeric>	<numeric>	<numeric>	<numeric>	<numeric>
ENSG00000279457	29.913579	0.1792571	0.3248216	0.551863	5.81042e-01
ENSG00000187634	183.229650	0.4264571	0.1402658	3.040350	2.36304e-03
ENSG00000188976	1651.188076	-0.6927205	0.0548465	-12.630158	1.43989e-36
ENSG00000187961	209.637938	0.7297556	0.1318599	5.534326	3.12428e-08
ENSG00000187583	47.255123	0.0405765	0.2718928	0.149237	8.81366e-01
ENSG00000187642	11.979750	0.5428105	0.5215599	1.040744	2.97994e-01

ENSG00000188290	108.922128	2.0570638	0.1969053	10.446970	1.51282e-25
ENSG00000187608	350.716868	0.2573837	0.1027266	2.505522	1.22271e-02
ENSG00000188157	9128.439422	0.3899088	0.0467163	8.346304	7.04321e-17
ENSG00000237330	0.158192	0.7859552	4.0804729	0.192614	8.47261e-01
	padj	symbol	entrez		name
	<numeric>	<character>	<character>		<character>
ENSG00000279457	6.86555e-01	NA	NA		NA
ENSG00000187634	5.15718e-03	SAMD11	148398	sterile alpha motif ..	
ENSG00000188976	1.76549e-35	NOC2L	26155	NOC2 like nucleolar ..	
ENSG00000187961	1.13413e-07	KLHL17	339451	kelch like family me..	
ENSG00000187583	9.19031e-01	PLEKHN1	84069	pleckstrin homology ..	
ENSG00000187642	4.03379e-01	PERM1	84808	PPARGC1 and ESRR ind..	
ENSG00000188290	1.30538e-24	HES4	57801	hes family bHLH tran..	
ENSG00000187608	2.37452e-02	ISG15	9636	ISG15 ubiquitin like..	
ENSG00000188157	4.21963e-16	AGRN	375790	agrin	
ENSG00000237330	NA	RNF223	401934	ring finger protein ..	

Q. Finally for this section let's reorder these results by adjusted p-value and save them to a CSV file in your current project directory.

```
res = res[order(res$pvalue),]
write.csv(res, file="deseq_results.csv")
```

Section 2: Pathway Analysis

```
library(pathview)
```

```
#####
Pathview is an open source software package distributed under GNU General
Public License version 3 (GPLv3). Details of GPLv3 is available at
http://www.gnu.org/licenses/gpl-3.0.html. Particullary, users are required to
formally cite the original Pathview paper (not just mention it) in publications
or products. For details, do citation("pathview") within R.
```

```
The pathview downloads and uses KEGG data. Non-academic uses may require a KEGG
license agreement (details at http://www.kegg.jp/kegg/legal.html).
```

```
#####
```

```
library(gage)
```

```
library(gageData)
```

```
data(kegg.sets.hs)
```

```
data(sigmet.idx.hs)
```

```
# Focus on signaling and metabolic pathways only
```

```
kegg.sets.hs = kegg.sets.hs[sigmet.idx.hs]
```

```
# Examine the first 3 pathways
```

```
head(kegg.sets.hs, 3)
```

```
$`hsa00232 Caffeine metabolism`
```

```
[1] "10" "1544" "1548" "1549" "1553" "7498" "9"
```

```
$`hsa00983 Drug metabolism - other enzymes`
```

```
[1] "10" "1066" "10720" "10941" "151531" "1548" "1549" "1551"  
[9] "1553" "1576" "1577" "1806" "1807" "1890" "221223" "2990"  
[17] "3251" "3614" "3615" "3704" "51733" "54490" "54575" "54576"  
[25] "54577" "54578" "54579" "54600" "54657" "54658" "54659" "54963"  
[33] "574537" "64816" "7083" "7084" "7172" "7363" "7364" "7365"  
[41] "7366" "7367" "7371" "7372" "7378" "7498" "79799" "83549"  
[49] "8824" "8833" "9" "978"
```

```
$`hsa00230 Purine metabolism`
```

```
[1] "100" "10201" "10606" "10621" "10622" "10623" "107" "10714"  
[9] "108" "10846" "109" "111" "11128" "11164" "112" "113"  
[17] "114" "115" "122481" "122622" "124583" "132" "158" "159"  
[25] "1633" "171568" "1716" "196883" "203" "204" "205" "221823"  
[33] "2272" "22978" "23649" "246721" "25885" "2618" "26289" "270"  
[41] "271" "27115" "272" "2766" "2977" "2982" "2983" "2984"  
[49] "2986" "2987" "29922" "3000" "30833" "30834" "318" "3251"  
[57] "353" "3614" "3615" "3704" "377841" "471" "4830" "4831"  
[65] "4832" "4833" "4860" "4881" "4882" "4907" "50484" "50940"  
[73] "51082" "51251" "51292" "5136" "5137" "5138" "5139" "5140"  
[81] "5141" "5142" "5143" "5144" "5145" "5146" "5147" "5148"  
[89] "5149" "5150" "5151" "5152" "5153" "5158" "5167" "5169"
```

```

[97] "51728" "5198" "5236" "5313" "5315" "53343" "54107" "5422"
[105] "5424" "5425" "5426" "5427" "5430" "5431" "5432" "5433"
[113] "5434" "5435" "5436" "5437" "5438" "5439" "5440" "5441"
[121] "5471" "548644" "55276" "5557" "5558" "55703" "55811" "55821"
[129] "5631" "5634" "56655" "56953" "56985" "57804" "58497" "6240"
[137] "6241" "64425" "646625" "654364" "661" "7498" "8382" "84172"
[145] "84265" "84284" "84618" "8622" "8654" "87178" "8833" "9060"
[153] "9061" "93034" "953" "9533" "954" "955" "956" "957"
[161] "9583" "9615"

```

The main `gage()` function requires a named vector of fold changes:

```

foldchanges = res$log2FoldChange
names(foldchanges) = res$entrez
head(foldchanges)

```

```

      1266      54855      1465      51232      2034      2317
-2.422719  3.201955 -2.313738 -2.059631 -1.888019 -1.649792

```

Run the `gage` pathway analysis

```

# Get the results
keggres = gage(foldchanges, gsets=kegg.sets.hs)

```

Look at the results:

```

attributes(keggres)

```

```

$names
[1] "greater" "less" "stats"

```

```

# Look at the first few down (less) pathways
head(keggres$less)

```

	p.geomean	stat.mean	p.val
hsa04110 Cell cycle	8.995727e-06	-4.378644	8.995727e-06
hsa03030 DNA replication	9.424076e-05	-3.951803	9.424076e-05
hsa03013 RNA transport	1.375901e-03	-3.028500	1.375901e-03
hsa03440 Homologous recombination	3.066756e-03	-2.852899	3.066756e-03

hsa04114	Oocyte meiosis	3.784520e-03	-2.698128	3.784520e-03
hsa00010	Glycolysis / Gluconeogenesis	8.961413e-03	-2.405398	8.961413e-03
		q.val	set.size	exp1
hsa04110	Cell cycle	0.001448312	121	8.995727e-06
hsa03030	DNA replication	0.007586381	36	9.424076e-05
hsa03013	RNA transport	0.073840037	144	1.375901e-03
hsa03440	Homologous recombination	0.121861535	28	3.066756e-03
hsa04114	Oocyte meiosis	0.121861535	102	3.784520e-03
hsa00010	Glycolysis / Gluconeogenesis	0.212222694	53	8.961413e-03

Now we can use `pathview()` to make a pathway plot with the RNA-seq expression results shown in color.

```
pathview(gene.data=foldchanges, pathway.id="hsa04110")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04110.pathview.png

```
#! [] (hsa04110.pathview.png)
```


Info: Writing image file hsa04110.pathview.pdf

Can also pull out top 5 upregulated pathways

```
## Focus on top 5 upregulated pathways here for demo purposes only
keggrespathways <- rownames(keggres$greater)[1:5]

# Extract the 8 character long IDs part of each string
keggresids = substr(keggrespathways, start=1, stop=8)
keggresids
```

```
[1] "hsa04640" "hsa04630" "hsa00140" "hsa04142" "hsa04330"
```

Pass those IDs in keggresids to the pathview function to draw plots for all the top 5 pathways

```
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04640.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04630.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14

Info: Writing image file hsa00140.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14


Info: Writing image file hsa04142.pathview.png


Info: some node width is different from others, and hence adjusted!

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04330.pathview.png





Q. Can you do the same procedure as above to plot the pathview figures for the top 5 down-regulated pathways?

```
## Focus on top 5 downregulated pathways
keggrespathwaysdown <- rownames(keggres$less)[1:5]

# Extract the 8 character long IDs part of each string
keggresidsdown = substr(keggrespathwaysdown, start=1, stop=8)
keggresidsdown
```

```
[1] "hsa04110" "hsa03030" "hsa03013" "hsa03440" "hsa04114"
```

```
pathview(gene.data=foldchanges, pathway.id=keggresidsdown, species="hsa")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04110.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14

Info: Writing image file hsa03030.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14

Info: Writing image file hsa03013.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14

Info: Writing image file hsa03440.pathview.png



'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/marin/Desktop/BIMM 143/Class 14



Info: Writing image file hsa04114.pathview.png

DNA REPLICATION



Replication complex (Bacteria)




Replication complex (Archaea)




Replication complex (Eukaryotes)



Data on KEGG graph
Rendered by Pathview





	p.geomean	stat.mean	p.val
G0:0007156 homophilic cell adhesion	8.519724e-05	3.824205	8.519724e-05
G0:0002009 morphogenesis of an epithelium	1.396681e-04	3.653886	1.396681e-04
G0:0048729 tissue morphogenesis	1.432451e-04	3.643242	1.432451e-04
G0:0007610 behavior	1.925222e-04	3.565432	1.925222e-04
G0:0060562 epithelial tube morphogenesis	5.932837e-04	3.261376	5.932837e-04
G0:0035295 tube development	5.953254e-04	3.253665	5.953254e-04

	q.val	set.size	exp1
G0:0007156 homophilic cell adhesion	0.1951953	113	8.519724e-05
G0:0002009 morphogenesis of an epithelium	0.1951953	339	1.396681e-04
G0:0048729 tissue morphogenesis	0.1951953	424	1.432451e-04
G0:0007610 behavior	0.1967577	426	1.925222e-04
G0:0060562 epithelial tube morphogenesis	0.3565320	257	5.932837e-04
G0:0035295 tube development	0.3565320	391	5.953254e-04

\$less

	p.geomean	stat.mean	p.val
G0:0048285 organelle fission	1.536227e-15	-8.063910	1.536227e-15
G0:0000280 nuclear division	4.286961e-15	-7.939217	4.286961e-15
G0:0007067 mitosis	4.286961e-15	-7.939217	4.286961e-15
G0:0000087 M phase of mitotic cell cycle	1.169934e-14	-7.797496	1.169934e-14
G0:0007059 chromosome segregation	2.028624e-11	-6.878340	2.028624e-11
G0:0000236 mitotic prometaphase	1.729553e-10	-6.695966	1.729553e-10

	q.val	set.size	exp1
G0:0048285 organelle fission	5.841698e-12	376	1.536227e-15
G0:0000280 nuclear division	5.841698e-12	352	4.286961e-15
G0:0007067 mitosis	5.841698e-12	352	4.286961e-15
G0:0000087 M phase of mitotic cell cycle	1.195672e-11	362	1.169934e-14
G0:0007059 chromosome segregation	1.658603e-08	142	2.028624e-11
G0:0000236 mitotic prometaphase	1.178402e-07	84	1.729553e-10

\$stats

	stat.mean	exp1
G0:0007156 homophilic cell adhesion	3.824205	3.824205
G0:0002009 morphogenesis of an epithelium	3.653886	3.653886
G0:0048729 tissue morphogenesis	3.643242	3.643242
G0:0007610 behavior	3.565432	3.565432
G0:0060562 epithelial tube morphogenesis	3.261376	3.261376
G0:0035295 tube development	3.253665	3.253665

Section 4. Reactome Analysis

Reactome is a database consisting of biological molecules and their relation to pathways and processes. We can conduct over-representation enrichment analysis and pathway-topology analysis Reactome using the previous list of significant genes generated from the differential expression results above. First output the list of significant genes at the 0.05 level as a plain text file:

```
sig_genes <- res[res$padj <= 0.05 & !is.na(res$padj), "symbol"]
print(paste("Total number of significant genes:", length(sig_genes)))
```

```
[1] "Total number of significant genes: 8147"
```

```
write.table(sig_genes, file="significant_genes.txt", row.names=FALSE, col.names=FALSE, quo
```

This file is inputted into the website: “<https://reactome.org/PathwayBrowser/#TOOL=AT>”. Parameters set to “Project to Humans”, then analyzed.

Q: What pathway has the most significant “Entities p-value”? Do the most significant pathways listed match your previous KEGG results? What factors could cause differences between the two methods?

Cell Cycle, Mitotic has the most significant Entities p-value. This somewhat matches the KEGG results. For example, cell cycle is the most downregulated pathway shown in the KEGG analysis. However, many of the other significant up or downregulated pathways, such as DNA replication and Jak-STAT signaling do not appear in the top of reactome’s list of significant p-values. Rather, some other pathways in gene expression and signaling are highlighted as significant, so the discrepancy might be due to the data being pulled from different sources that use different names to describe similar pathways.

```
head(keggres$less)
```

	p.geomean	stat.mean	p.val
hsa04110 Cell cycle	8.995727e-06	-4.378644	8.995727e-06
hsa03030 DNA replication	9.424076e-05	-3.951803	9.424076e-05
hsa03013 RNA transport	1.375901e-03	-3.028500	1.375901e-03
hsa03440 Homologous recombination	3.066756e-03	-2.852899	3.066756e-03
hsa04114 Oocyte meiosis	3.784520e-03	-2.698128	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	8.961413e-03	-2.405398	8.961413e-03

	q.val	set.size	exp1
hsa04110 Cell cycle	0.001448312	121	8.995727e-06
hsa03030 DNA replication	0.007586381	36	9.424076e-05
hsa03013 RNA transport	0.073840037	144	1.375901e-03
hsa03440 Homologous recombination	0.121861535	28	3.066756e-03
hsa04114 Oocyte meiosis	0.121861535	102	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	0.212222694	53	8.961413e-03

```
head(keggres$greater)
```

	p.geomean	stat.mean	p.val
hsa04640 Hematopoietic cell lineage	0.002822776	2.833362	0.002822776
hsa04630 Jak-STAT signaling pathway	0.005202070	2.585673	0.005202070
hsa00140 Steroid hormone biosynthesis	0.007255099	2.526744	0.007255099
hsa04142 Lysosome	0.010107392	2.338364	0.010107392
hsa04330 Notch signaling pathway	0.018747253	2.111725	0.018747253
hsa04916 Melanogenesis	0.019399766	2.081927	0.019399766

	q.val	set.size	exp1
hsa04640 Hematopoietic cell lineage	0.3893570	55	0.002822776
hsa04630 Jak-STAT signaling pathway	0.3893570	109	0.005202070
hsa00140 Steroid hormone biosynthesis	0.3893570	31	0.007255099
hsa04142 Lysosome	0.4068225	118	0.010107392
hsa04330 Notch signaling pathway	0.4391731	46	0.018747253
hsa04916 Melanogenesis	0.4391731	90	0.019399766