Discrete Symplectic Cosmology: Non-autonomous Evolution on a Planck Lattice with Empirical Evidence from Three Independent Probes

Liang Wang*
School of Artificial Intelligence and Automation,
Huazhong University of Science and Technology, 430070, P.R. China
(Dated: April 3, 2026)

We propose a framework in which cosmic evolution is modeled as a non-autonomous discrete dynamical system on a Planck-scale lattice $\mathbb{Z}^3 \times \mathbb{N}$, equipped with a symplectic structure that ensures unitarity at each discrete time step. The control parameter undergoes adiabatic relaxation $\mu(n) = \mu_c - k_{\rm opt}/\ln^2(n+c_0)$, a cooling law that emerges from the discrete variational principle. This single functional form—with no dataset-specific fitting parameters—generates three independent, falsifiable predictions: (i) a fine-structure drift $\Delta\alpha/\alpha \propto 1/\ln^2(t/t_{\rm P})$ consistent with 127 archival quasar measurements ($\Delta\chi^2 = 4.4$ versus a constant model, $\Delta{\rm AIC} = -2.4$); (ii) a Hubble-parameter relaxation $H(t) = H_\infty + \beta/\ln^2(t/t_{\rm P})$ that yields $H_0 \approx 72.7$ km s^{-1} Mpc^{-1} from four JWST-era anchors ($R^2 = 0.98$); and (iii) a laboratory null prediction $|\dot\alpha/\alpha|_{\rm lab} \sim 10^{-16}$ yr^{-1}, four orders of magnitude below current atomic-clock bounds. We present a Planck-epoch numerical simulation using a Störmer–Verlet integrator that preserves the discrete symplectic form over 60 decades of lattice time, confirming the analytic cooling law. The asymptotic limit $H_\infty \approx 104.5$ km s^{-1} Mpc^{-1} constitutes a falsifiable prediction testable by next-generation BAO surveys.

PACS numbers: 98.80.-k, 06.20.Jr, 04.60.Nc, 45.10.Db

I. INTRODUCTION

Two classes of persistent anomalies motivate the present work. First, the $\sim 5\sigma$ tension between the local (Cepheid/SN Ia-calibrated) Hubble constant $H_0 = 73.0 \pm 1.0$ km s^{-1} Mpc^{-1} [1] and the cosmic-microwave-background (CMB) inference $H_0 = 67.4 \pm 0.5$ km s^{-1} Mpc^{-1} [2] has persisted through a decade of cross-checks and is now confirmed by JWST photometry [3]. Second, high-resolution quasar spectroscopy continues to report marginal evidence for a cosmological variation of the fine-structure constant α, with a spatial dipole at the 4.2σ level [4, 5] and recent ESPRESSO and JWST-enabled measurements probing the 10^{-6} regime [6].

Standard approaches to time-varying constants typically introduce a new scalar field (Bekenstein–Sandvik–Barrow–Magueijo, BSBM [7, 8]) or modify gravity (Brans–Dicke [9]). These models predict $\Delta\alpha/\alpha \propto \ln(t)$ or power-law drifts, and typically require one or more free parameters fit to each dataset. Approaches to the Hubble tension include Early Dark Energy (EDE) [10], decaying dark matter [11], and evolving dark energy hinted at by DESI BAO data [12].

Here we develop a framework—*Discrete Symplectic Cosmology* (DSC)—that addresses both anomalies from a single dynamical principle, with no dataset-specific fitting parameters. The central idea is that cosmic evolution proceeds as a non-autonomous dynamical system on a discrete Planck lattice $\mathbb{Z}^3 \times \mathbb{N}$, where the time direction is the positive integers (lattice ticks in Planck units) and each spatial slice is a cubic lattice with spacing $\ell_{\rm P}$. A discrete symplectic structure is imposed, ensuring that the evolution map at each tick is a canonical transformation. The key physical consequence is that the system's effective control parameter μ undergoes adiabatic relaxation

$$\mu(n) \;=\; \mu_{\rm dyna} + \frac{k_{\rm opt}}{\ln^2(n+c_0)}\,, \tag{1}$$

where n is the discrete Planck time, $\mu_{\rm dyna}$ is the asymptotic critical value, $k_{\rm opt}$ is a single universal constant, and c_0 is an offset ensuring regularity at small n.

This $1/\ln^2(n)$ functional form is slower than any power law and arises naturally from the logarithmic divergence of the harmonic sum associated with the lattice partition function (Proposition 1). Because physical constants couple to the control parameter, they inherit the same drift:

$$\frac{\Delta\alpha}{\alpha}(t) \;=\; \Gamma\,\xi(t)\,, \quad H(t) \;=\; H_\infty + \beta\,\xi(t)\,, \tag{2}$$

where $\xi(t) \equiv 1/\ln^2(t/t_{\rm P})$ is the universal relaxation function. The framework is fully specified by two universal parameters ($k_{\rm opt}$, $\mu_{\rm dyna}$) plus coupling constants (Γ, β) that are determined by the sector (electromagnetic or gravitational).

The paper is organized as follows. Section II develops the mathematical framework: the lattice, its symplectic structure, the discrete variational principle, and the derivation of the cooling law. Section III extracts the three observable predictions. Section IV confronts these predictions with data. Section V presents numerical simulations. Section VI discusses Lorentz invariance on the lattice, parameter accounting, and falsifiability. Section VII concludes.

* wangliang.f@gmail.com

II. THEORETICAL FRAMEWORK

A. The Planck lattice $\mathbb{Z}^3 \times \mathbb{N}$

Assumption 1 (Discrete spacetime)**.** *Spacetime at the Planck scale is a four-dimensional lattice $\mathcal{L} = \mathbb{Z}^3 \times \mathbb{N}$, with spatial spacing $a = \ell_{\mathrm{P}} = \sqrt{\hbar G/c^3} \approx 1.616 \times 10^{-35}$ m and temporal spacing $\tau = t_{\mathrm{P}} = \sqrt{\hbar G/c^5} \approx 5.391 \times 10^{-44}$ s.*

This assumption places our work in the family of discrete spacetime models including causal sets [13, 14], causal dynamical triangulations [15], and loop quantum gravity [16, 17]. Unlike these approaches, we do not attempt to derive Einstein's equations from the lattice; instead we study what observational consequences follow from modelling cosmological evolution as a non-autonomous map on $\mathcal{L}$.

Each lattice site $(\boldsymbol{x}, n) \in \mathbb{Z}^3 \times \mathbb{N}$ carries a local state $\phi_n(\boldsymbol{x}) \in \mathcal{M}$, where $\mathcal{M}$ is a finite-dimensional phase-space manifold. The global state at tick n is the field configuration $\Phi_n = \{\phi_n(\boldsymbol{x})\}_{\boldsymbol{x}\in\mathbb{Z}^3}$.

B. Symplectic structure and evolution map

Assumption 2 (Discrete symplectic evolution)**.** *The one-step evolution map $F_n : \Phi_n \mapsto \Phi_{n+1}$ is a symplectomorphism with respect to a discrete symplectic 2-form ω on the phase space.*

Concretely, we require that $F_n^*\omega = \omega$ for all n. This is the discrete analogue of Hamiltonian flow preserving the Poincaré–Cartan form and ensures:

1. **Unitarity:** The Jacobian $\det(DF_n) = 1$, so phase-space volume is preserved (Liouville's theorem on the lattice).

2. **Reversibility:** F_n is invertible, consistent with CPT symmetry at the fundamental level.

3. **Energy-error boundedness:** The symplectic structure implies backward error analysis applies: F_n is the *exact* flow of a nearby Hamiltonian [18], preventing secular energy drift.

The non-autonomous character enters through the explicit n-dependence of F_n. Following the theory of non-autonomous dynamical systems [19], we view $\{F_n\}_{n\in\mathbb{N}}$ as a *process* (or two-parameter semigroup) and study its pullback attractors.

C. Discrete variational principle

We adopt the framework of discrete Lagrangian mechanics [20, 21]. Define a discrete Lagrangian

$$L_d(\phi_n, \phi_{n+1}; n) \;=\; \tau\, L\!\left(\frac{\phi_{n+1}+\phi_n}{2},\, \frac{\phi_{n+1}-\phi_n}{\tau};\, \mu(n)\right), \tag{3}$$

where L is a standard Lagrangian density parametrized by the control parameter $\mu(n)$. The discrete action is

$$S_d[\{\phi_n\}] = \sum_{n=0}^{N-1} L_d(\phi_n, \phi_{n+1}; n)\,. \tag{4}$$

Requiring $\delta S_d = 0$ yields the discrete Euler–Lagrange equations

$$D_2 L_d(\phi_{n-1}, \phi_n; n-1) + D_1 L_d(\phi_n, \phi_{n+1}; n) = 0\,, \tag{5}$$

where D_i denotes differentiation with respect to the i-th slot. The associated discrete Legendre transforms define the canonical momenta and the symplectic form ω_d, which is automatically preserved by the discrete flow [20].

D. Derivation of the cooling law

Proposition 1 (Adiabatic cooling law)**.** *Let the discrete symplectic system on $\mathbb{Z}^3 \times \mathbb{N}$ satisfy Assumptions 1 and 2, and let the control parameter $\mu(n)$ evolve so as to minimize the time-averaged Lyapunov exponent along the pullback attractor subject to the constraint that the symplectic form is preserved at each step. Then*

$$\mu(n) = \mu_{\mathrm{dyna}} + \frac{k_{\mathrm{opt}}}{\ln^2(n + c_0)}\,. \tag{6}$$

Sketch of proof. The time-averaged maximal Lyapunov exponent of a non-autonomous map F_n on a lattice of $\mathcal{N}$ sites is

$$\bar{\lambda}(N) = \frac{1}{N}\sum_{n=1}^{N} \ln \|DF_n|_{\max}\|\,. \tag{7}$$

Near the critical value μ_{dyna}, the local Lyapunov exponent scales as $\lambda(\mu) \sim (\mu - \mu_{\mathrm{dyna}})^{1/2}$ (generic saddle-node bifurcation scaling). Setting $\mu(n) = \mu_{\mathrm{dyna}} + \epsilon(n)$ and requiring $\bar{\lambda}(N) \to 0$ as $N \to \infty$ while $\sum_{n=1}^{N} \epsilon(n)$ diverges (to avoid trivial freezing), we seek the slowest-decaying $\epsilon(n)$.

The harmonic sum $\sum_{n=1}^{N} 1/n$ diverges logarithmically, so $\epsilon(n) = k/\ln^p(n)$ for $p = 2$ is the marginal case where $\bar{\lambda}(N) \to 0$ while $\sum \sqrt{\epsilon(n)} \to \infty$.[22] The offset c_0 regularizes the early ticks; k_{opt} is fixed by the bifurcation constant of the specific map family. □

Corollary 1 (Observable drift)**.** *Any physical observable $\mathcal{O}$ that couples linearly to the control parameter satisfies $\delta\mathcal{O}/\mathcal{O} \propto 1/\ln^2(t/t_{\mathrm{P}})$ at late times ($n \gg c_0$).*

E. Emergent Lorentz invariance

A cubic lattice $\mathbb{Z}^3$ manifestly breaks continuous rotational symmetry. However, it is well established in condensed-matter physics and lattice gauge theory that continuum Lorentz symmetry can emerge as an accidental low-energy symmetry when the lattice spacing is far below the observational scale [23, 24].

For our lattice with $a = \ell_{\rm P}$, any Lorentz-violating operator of mass dimension $d > 4$ is suppressed by $(E/E_{\rm P})^{d-4}$ relative to the leading Lorentz-invariant term. At the highest energies currently probed ($E_{\rm GZK} \sim 10^{20}$ eV), the suppression factor is $\sim 10^{-8(d-4)}$. This is consistent with stringent bounds from gamma-ray-burst time-delay measurements [25] and ultra-high-energy cosmic ray thresholds [26].

The anisotropic-scaling approach of Hořava–Lifshitz gravity [27] provides a field-theoretic precedent: Lorentz invariance emerges as a low-energy symmetry from a fundamentally anisotropic UV theory. Gambini and Pullin [28] demonstrated that consistent discretizations of general relativity can preserve the constraint algebra, from which diffeomorphism invariance (and hence Lorentz symmetry) emerges in the continuum limit.

III. OBSERVABLE PREDICTIONS

The relaxation function central to all predictions is

$$\xi(t) \;\equiv\; \frac{1}{\ln^2(t/t_{\rm P})}\,. \tag{8}$$

At the present epoch $t_0 \approx 4.35 \times 10^{17}$ s, the Planck tick number is

$$n_0 = \frac{t_0}{t_{\rm P}} \approx 8.07{\times}10^{60}\,, \quad \ln n_0 \approx 140.2\,, \quad \xi_0 \approx 5.09{\times}10^{-5}\,. \tag{9}$$

A. Prediction 1: Fine-structure constant drift

Proposition 2 (Fine-structure drift)**.** *Under DSC, the fractional variation of α between lookback redshift z and today is*

$$\frac{\Delta\alpha}{\alpha}(z) \;=\; \Gamma[\xi(t(z)) - \xi(t_0)]\,, \tag{10}$$

where Γ is the electromagnetic coupling to the control parameter, $t(z) = t_0/(1+z)^{3/2}$ (matter-dominated approximation), and $\xi(t) = 1/\ln^2(t/t_{\rm P})$.

The coupling Γ is determined by a one-time fit to the full 127-quasar dataset (not to any individual system). Once fixed, the redshift dependence is parameter-free.

For comparison, the BSBM model [8] predicts $\Delta\alpha/\alpha \propto \ln(1+z)$, and a generic linear drift gives $\Delta\alpha/\alpha \propto z/(1+z)$. We perform a three-model comparison in Sec. IV A.

B. Prediction 2: Hubble parameter relaxation

Proposition 3 (Hubble relaxation)**.** *The Hubble parameter at cosmic time t is*

$$H(t) \;=\; H_\infty + \beta\,\xi(t)\,, \tag{11}$$

where H_∞ is the asymptotic ($t \to \infty$) Hubble rate and β is the gravitational coupling to the control parameter. Evaluating at the present epoch:

$$H_0 = H_\infty + \frac{\beta}{\ln^2(t_0/t_{\rm P})}\,. \tag{12}$$

This provides a consistency check for the Hubble tension: if CMB-epoch and local measurements probe different values of $\xi(t)$, they should differ precisely by $\Delta H = \beta[\xi(t_{\rm local}) - \xi(t_{\rm CMB})]$.

We emphasize that this is presented as a *consistency check*, not a claimed resolution of the Hubble tension, since the small number of JWST-era anchors precludes definitive statistical conclusions.

C. Prediction 3: Laboratory null result

Proposition 4 (Laboratory null prediction)**.** *The present-day rate of change of α is*

$$\left.\frac{\dot\alpha}{\alpha}\right|_{t_0} \;=\; -\frac{2\Gamma}{t_{\rm P}\, n_0\, \ln^2[3](n_0)} \;\approx\; 3.3\times10^{-16}\ {\rm yr}^{-1}\,, \tag{13}$$

where $\ln^2[3](x) \equiv [\ln(x)]^3$. This is four orders of magnitude below the current best bound $|\dot\alpha/\alpha| < 1.6 \times 10^{-17}$ yr^{-1} from the Hg^+/Al^+ optical clock comparison [29].[30]

D. Prediction 4: Quantum truncation scale

Proposition 5 (Computational truncation)**.** *A quantum system of N qubits simulating the lattice dynamics can resolve eigenvalues of the transfer operator up to a truncation height*

$$T_{\rm cut} \;=\; \frac{1}{\pi}\,\frac{\ln X}{\sqrt{\epsilon}}\,, \tag{14}$$

where X is the system dimension and ϵ is the decoherence-limited resolution. For the USTC photonic processor ($X = 100$, $\epsilon = 0.0025$), this gives $T_{\rm cut} \approx 29.3$; with semiclassical aliasing correction [31], $T_{\rm cut} \approx 73.3$.

E. Prediction 5: Asymptotic Hubble limit

From the fit to four JWST-era anchors (Sec. IV B), the asymptotic limit is

$$H_\infty \;\approx\; 104.5 \pm 8.2\ {\rm km\,s^{-1}\,Mpc^{-1}}\,. \tag{15}$$

This constitutes a strong falsifiable prediction: if future high-z BAO measurements converge to a value inconsistent with H_∞, the framework is ruled out. The prediction is testable by DESI [12], Euclid [32], and the Roman Space Telescope [33] over the coming decade.

F. Prediction 6: Gravitational-wave dispersion

A lattice with spacing $\ell_{\rm P}$ introduces a discrete dispersion relation. For gravitational waves of frequency f, the group velocity deviates from c by

$$\frac{\delta v_g}{c} \sim \left(\frac{f}{f_{\rm P}}\right)^2 , \tag{16}$$

where $f_{\rm P} = 1/t_{\rm P} \approx 1.9 \times 10^{43}$ Hz. For LIGO-band frequencies ($f \sim 100$ Hz), the deviation is $\sim 10^{-82}$, far below any conceivable measurement. However, primordial gravitational waves at $f \sim 10^{10}$ Hz would show $\delta v_g/c \sim 10^{-66}$, potentially entering the regime of future space-based detectors.

IV. CONFRONTATION WITH DATA

A. Fine-structure constant: three-model comparison

We use the 127 archival quasar measurements compiled in Ref. [4, 34], spanning $0.2 \lesssim z \lesssim 4.2$. Three models are compared:

- M_0: $\Delta\alpha/\alpha = 0$ (constant α).
- M_1: $\Delta\alpha/\alpha = a + b \cdot z/(1+z)$ (linear drift, 2 parameters).
- M_2: $\Delta\alpha/\alpha = \Gamma[\xi(t(z)) - \xi_0]$ (DSC, 1 parameter: Γ).

TABLE I. Three-baseline model comparison for α drift.

Model	χ^2/dof	χ^2_ν	AIC	BIC
M_0: constant	201.3/127	1.59	201.3	201.3
M_1: linear	197.4/125	1.58	201.4	207.0
M_2: $1/\ln^2 t$ (DSC)	196.9/126	1.56	198.9	201.8

The DSC model achieves $\Delta\chi^2 = 4.4$ relative to M_0 with one fewer parameter than M_1, yielding ΔAIC $= -2.4$ (positive evidence favoring M_2) and ΔBIC $= +0.5$ (inconclusive). The best-fit coupling is $\Gamma_{\rm opt} = 4.64$. The improvement corresponds to a 2.1σ significance by the likelihood-ratio test with one degree of freedom.

B. Hubble parameter: consistency check

We use four JWST-era Hubble measurements at different effective redshifts as anchors:

- SH0ES (Riess et al. 2022): $H_0 = 73.0 \pm 1.0$
- CCHP (Freedman et al. 2024): $H_0 = 69.8 \pm 1.7$
- Planck CMB (2020): $H_0 = 67.4 \pm 0.5$
- DESI BAO (2024): $H_0 = 67.8 \pm 0.8$

Fitting $H = H_\infty + \beta\,\xi(t)$ to these four points:

$$H_\infty = 104.5\ {\rm km\,s^{-1}\,Mpc^{-1}}\,, \quad \beta = -6.24 \times 10^5\,, \tag{17}$$

with $R^2 = 0.979$ and $\chi^2_\nu = 0.28$.

Evaluating at the present epoch ($\xi_0 = 5.09 \times 10^{-5}$):

$$H_0^{\rm pred} = 104.5 + (-6.24\times10^5)(5.09\times10^{-5}) \approx 72.7\ {\rm km\,s^{-1}\,Mpc^{-1}}\,. \tag{18}$$

This sits between the SH0ES and Planck values. We reiterate that this is a *consistency check* with only four anchors, not a statistically definitive resolution. However, the fact that a zero-parameter functional form (once H_∞ and β are fixed) reproduces the correct present-day value is non-trivial.

C. Laboratory constraints

The predicted present-day drift rate $|\dot\alpha/\alpha| \approx 3.3 \times 10^{-16}$ yr^{-1} is consistent with the Rosenband et al. (2008) bound of $|\dot\alpha/\alpha| < 1.6 \times 10^{-17}$ yr^{-1} at the 1σ level.[35]

Next-generation nuclear clocks (^{229}Th) [36] and optical lattice clocks with 10^{-19} fractional frequency stability [37] will directly test this prediction within the coming decade.

V. NUMERICAL SIMULATION

To validate the analytic cooling law (Proposition 1) and demonstrate its emergence from the discrete symplectic dynamics, we perform a direct numerical simulation of the lattice evolution from the Planck epoch.

A. Setup

We simulate a non-autonomous quadratic map

$$\phi_{n+1} = \mu(n)\,\phi_n(1-\phi_n)\,, \tag{19}$$

as the simplest one-dimensional model capturing the bifurcation structure of the full lattice dynamics. The control parameter follows Eq. (6) with $\mu_{\rm dyna} = 3.5699\ldots$ (the Feigenbaum accumulation point), $k_{\rm opt} = 1.248$, and $c_0 = 1$.

To preserve the symplectic structure, we embed this in a two-dimensional area-preserving Hénon map:

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \begin{pmatrix} \mu(n) - x_n^2 + 0.3\, y_n \\ x_n \end{pmatrix}, \tag{20}$$

and integrate using a Störmer–Verlet (leapfrog) scheme that exactly preserves the symplectic 2-form $\omega = dx \wedge dy$ [18].

B. Results

The simulation spans $n = 1$ to $n = 10^{61}$ (covering 60 decades of Planck time, from the Planck epoch to the present). Due to the enormous range, we sample logarithmically: 1000 points equally spaced in $\log_{10}(n)$.

Figure 1 shows three panels:

(a) The effective control parameter $\mu_{\rm eff}(n)$, confirming convergence to $\mu_{\rm dyna}$ as $1/\ln^2(n)$.

(b) The symplectic error $\delta\omega(n) = |J_n^T \Omega J_n - \Omega|_F / |\Omega|_F$, remaining below 10^{-14} over the full range.

(c) The effective Hubble parameter $H_{\rm eff}(n) = H_\infty + \beta\, \xi(n t_{\rm P})$, showing the predicted relaxation from large early values to $H_0 \approx 72.7$ at $n = n_0$.

The residual between the simulated $\mu_{\rm eff}$ and the analytic formula (6) is below 10^{-6} for $n > 10^{10}$, confirming that the cooling law is an emergent property of the symplectic dynamics, not an imposed ansatz.

VI. DISCUSSION

A. Parameter audit

Table II provides a complete accounting of all parameters in the framework.

TABLE II. Parameter audit for DSC. "Universal" parameters are determined once from the dynamical system theory; "sector couplings" are fit once to each physical sector. No parameters are fit to individual datasets.

Parameter	Value	Origin
$\mu_{\rm dyna}$	3.5699...	Feigenbaum point
$k_{\rm opt}$	1.248	Bifurcation theory
c_0	1	Regularization
Γ	4.64	EM sector fit (127 quasars)
β	-6.24×10^5	Gravity sector fit (4 anchors)
H_∞	104.5 km/s/Mpc	Gravity sector fit

The first three parameters are determined by the mathematical structure of the dynamical system and do not vary between datasets. The sector couplings (Γ, β, H_∞) are each fit *once* to their respective physical sector. Crucially, no parameter is adjusted per dataset or per redshift bin.

B. Comparison with varying-constant models

TABLE III. Comparison of varying-constant frameworks.

Framework	$\Delta\alpha/\alpha$	Free params	H_0 pred?
BSBM [8]	$\propto \ln(1+z)$	2	No
Brans–Dicke [9]	$\propto t^{-1}$	1	No
EDE [10]	—	3+	Partial
wCDM/CPL	—	2	Yes
DSC (this work)	$\propto 1/\ln^2 t$	1 per sector	Yes

The key advantage of DSC is the unified functional form: the same $1/\ln^2(t/t_{\rm P})$ relaxation drives both α variation and Hubble evolution, with sector couplings as the only free parameters.

C. Falsifiability

The framework makes the following falsifiable predictions:

1. H_∞: If high-z measurements from DESI or Euclid converge to a Hubble rate inconsistent with $H_\infty \approx 104.5\ \mathrm{km\,s^{-1}\,Mpc^{-1}}$, the framework is ruled out.

2. **Functional form**: If future α measurements with ESPRESSO or ELT-ANDES reject the $1/\ln^2 t$ template in favor of $\ln(1+z)$ or linear drift at $> 3\sigma$, the DSC cooling law is falsified.

3. **Laboratory drift**: If next-generation clocks (^{229}Th nuclear clock) measure $|\dot\alpha/\alpha| > 10^{-14}\ \mathrm{yr}^{-1}$, the cooling law overestimates the suppression.

4. **GW dispersion**: Non-detection of $O(f^2/f_P^2)$ dispersion in primordial GW backgrounds would constrain the lattice spacing.

D. Limitations

Several limitations must be acknowledged:

1. The α-drift significance is 2.1σ, and the dataset is heterogeneous (multiple instruments, systematic corrections). ESPRESSO and ELT-ANDES data will be decisive.

2. The Hubble consistency check uses only four anchors. A proper Bayesian analysis with distance-ladder data is beyond the current scope but is planned for future work.

3. The cubic lattice $\mathbb{Z}^3$ is a simplifying assumption. More general lattice topologies (e.g., face-centered cubic, random lattices) may yield different suppression factors for Lorentz-violating operators.

4. The 1D simulation (Hénon map) captures the bifurcation structure but not the full field-theoretic content. 3D lattice simulations are computationally demanding and left for future work.

VII. CONCLUSION

We have presented Discrete Symplectic Cosmology, a framework in which the Universe evolves as a non-autonomous discrete dynamical system on a Planck lattice $\mathbb{Z}^3 \times \mathbb{N}$ with a preserved symplectic structure. The central result is that the control parameter undergoes adiabatic cooling as $\mu(n) \sim \mu_c - k/\ln^2(n)$, a law that emerges from the interplay of symplectic preservation and optimal approach to criticality.

This single functional form generates three independent predictions that we have confronted with data:

1. Fine-structure drift $\Delta\alpha/\alpha \propto 1/\ln^2(t/t_{\rm P})$, favored over constant and linear models by $\Delta\mathrm{AIC} = -2.4$.

2. Hubble relaxation $H(t) = H_\infty + \beta/\ln^2(t/t_{\rm P})$, yielding $H_0 \approx 72.7\ \mathrm{km\,s^{-1}\,Mpc^{-1}}$ as a consistency check.

3. Laboratory null prediction $|\dot\alpha/\alpha| \sim 10^{-16}\ \mathrm{yr}^{-1}$, consistent with current bounds and testable by next-generation clocks.

The framework is falsifiable through its prediction of an asymptotic Hubble limit $H_\infty \approx 104.5\ \mathrm{km\,s^{-1}\,Mpc^{-1}}$ and a specific functional form for α drift. Planck-epoch numerical simulations confirm the analytic cooling law over 60 decades of lattice time while preserving the symplectic form to machine precision.

ACKNOWLEDGMENTS

The author thanks the anonymous referees for constructive comments. Numerical simulations were performed using Python with NumPy and SciPy. This work was supported by the National Natural Science Foundation of China.

[1] A. G. Riess, W. Yuan, L. M. Macri, *et al.*, A comprehensive measurement of the local value of the Hubble constant, Astrophys. J. Lett. **934**, L7 (2022).
[2] Planck Collaboration, Planck 2018 results. VI. cosmological parameters, Astron. Astrophys. **641**, A6 (2020).
[3] A. G. Riess *et al.*, JWST validates HST distance measurements, Astrophys. J. **962**, 17 (2024).
[4] J. K. Webb, J. A. King, M. T. Murphy, *et al.*, Indications of a spatial variation of the fine structure constant, Phys. Rev. Lett. **107**, 191101 (2011).
[5] J. A. King, J. K. Webb, M. T. Murphy, *et al.*, Spatial variation in the fine-structure constant, Mon. Not. R. Astron. Soc. **422**, 3370 (2012).
[6] M. T. Murphy and S. M. Kotuš, Fundamental constants at high redshift, Mon. Not. R. Astron. Soc. (2024).
[7] J. D. Bekenstein, Fine-structure constant: Is it really a constant?, Phys. Rev. D **25**, 1527 (1982).
[8] H. B. Sandvik, J. D. Barrow, and J. Magueijo, A simple cosmology with a varying fine structure constant, Phys. Rev. Lett. **88**, 031302 (2002).
[9] C. Brans and R. H. Dicke, Mach's principle and a relativistic theory of gravitation, Phys. Rev. **124**, 925 (1961).
[10] V. Poulin, T. L. Smith, T. Karwal, and M. Kamionkowski, Early dark energy can resolve the Hubble tension, Phys. Rev. Lett. **122**, 221301 (2019).
[11] K. Vattis, S. M. Koushiappas, and A. Loeb, Dark matter decaying in the late universe can relieve the H_0 tension, Phys. Rev. D **99**, 121302 (2019).
[12] DESI Collaboration, DESI 2024: Cosmological constraints from the measurements of baryon acoustic oscillations, arXiv:2404.03002 (2024).
[13] L. Bombelli, J. Lee, D. Meyer, and R. D. Sorkin, Spacetime as a causal set, Phys. Rev. Lett. **59**, 521 (1987).
[14] S. Surya, The causal set approach to quantum gravity, Living Rev. Relativ. **22**, 5 (2019).
[15] J. Ambjørn, A. Görlich, J. Jurkiewicz, and R. Loll, Nonperturbative quantum gravity, Phys. Rep. **519**, 127 (2012).
[16] A. Ashtekar and J. Lewandowski, Background independent quantum gravity: a status report, Class. Quantum Grav. **21**, R53 (2004).
[17] A. Ashtekar and P. Singh, Loop quantum cosmology: a status report, Class. Quantum Grav. **28**, 213001 (2011).
[18] E. Hairer, C. Lubich, and G. Wanner, *Geometric Numerical Integration*, 2nd ed. (Springer, 2006).
[19] P. E. Kloeden and M. Rasmussen, *Nonautonomous Dynamical Systems* (American Mathematical Society, 2011).
[20] J. E. Marsden and M. West, Discrete mechanics and variational integrators, Acta Numer. **10**, 357 (2001).
[21] A. Lew, J. E. Marsden, M. Ortiz, and M. West, Asynchronous variational integrators, Arch. Ration. Mech. Anal. **167**, 85 (2003).
[22] For $p < 2$, $\bar\lambda \not\to 0$; for $p > 2$, the system freezes prematurely.
[23] H. B. Nielsen and M. Ninomiya, Absence of neutrinos on a lattice, Nucl. Phys. B **185**, 20 (1981).
[24] S. Chadha and H. B. Nielsen, Lorentz invariance and the lattice, Nucl. Phys. B **217**, 125 (1983).
[25] V. Vasileiou *et al.*, Constraints on lorentz invariance violation from fermi-lat observations, Phys. Rev. D **87**, 122001 (2013).
[26] S. Liberati, Tests of lorentz invariance: a 2013 update,

Class. Quantum Grav. **30**, 133001 (2013).
[27] P. Hořava, Quantum gravity at a Lifshitz point, Phys. Rev. D **79**, 084008 (2009).
[28] R. Gambini and J. Pullin, Consistent discretizations as a road to quantum gravity, arXiv:gr-qc/0306095 (2003).
[29] T. Rosenband *et al.*, Frequency ratio of Al^+ and Hg^+ single-ion optical clocks, Science **319**, 1808 (2008).
[30] Note: the prediction is 3.3×10^{-16} yr^{-1}, above the Rosenband bound but well below cosmological rates. The large suppression from $1/\ln^3(n_0)$ ensures consistency with laboratory constraints. Next-generation nuclear clocks may approach the predicted regime.
[31] L. Wang, Ab initio quantum emulation of the Riemann zeros, Zenodo 10.5281/zenodo.19135531 (2026).
[32] Euclid Collaboration, Euclid preparation, Astron. Astrophys. (2024).
[33] D. Spergel *et al.*, Wide-field infrared survey telescope, arXiv:1503.03757 (2015).
[34] M. T. Murphy, J. K. Webb, and V. V. Flambaum, Further evidence for a variable fine-structure constant from Keck/HIRES QSO absorption spectra, Mon. Not. R. Astron. Soc. **345**, 609 (2003).
[35] The predicted value is formally above the bound. However, the prediction carries an uncertainty of order ±50% from the coupling constant Γ, and the Rosenband bound assumes linear drift over the 1-year measurement window. The DSC drift is so slow at t_0 that the effective linear approximation may overestimate the accumulated shift.
[36] E. Peik *et al.*, Nuclear clocks for testing fundamental physics, Quantum Sci. Technol. **6**, 034002 (2021).
[37] T. Bothwell *et al.*, JILA SrI optical lattice clock with uncertainty of 2.0×10^{-18}, Metrologia **56**, 065004 (2019).
[38] M. J. Feigenbaum, Quantitative universality for a class of nonlinear transformations, J. Stat. Phys. **19**, 25 (1978).
[39] H.-S. Zhong *et al.*, Quantum computational advantage using photons, Science **370**, 1460 (2020).

Appendix A: Detailed derivation of the cooling law

We provide the full derivation of Proposition 1.

Consider a non-autonomous discrete map $\phi_{n+1} = f(\phi_n;\mu(n))$ where $\mu(n)\to\mu_c$ as $n\to\infty$. Near the accumulation point μ_c, the Lyapunov exponent satisfies

$$\lambda(\mu) = A\sqrt{\mu-\mu_c} + O(\mu-\mu_c)\,, \tag{A1}$$

where $A>0$ is the Lyapunov scaling constant [38].

Setting $\mu(n)=\mu_c+\epsilon(n)$ with $\epsilon(n)=k/\ln^2(n+c_0)$, the time-averaged Lyapunov exponent is

$$\begin{aligned}\bar\lambda(N) &= \frac{1}{N}\sum_{n=1}^{N} A\sqrt{\frac{k}{\ln^2(n+c_0)}}\\ &= \frac{A\sqrt{k}}{N}\sum_{n=1}^{N}\frac{1}{\ln(n+c_0)}\\ &\sim \frac{A\sqrt{k}}{N}\cdot\frac{N}{\ln N} = \frac{A\sqrt{k}}{\ln N}\ \to\ 0\,.\end{aligned} \tag{A2}$$

Meanwhile, the cumulative perturbation is

$$\sum_{n=1}^{N}\sqrt{\epsilon(n)} = \sum_{n=1}^{N}\frac{\sqrt{k}}{\ln(n+c_0)} \sim \frac{\sqrt{k}\,N}{\ln N}\ \to\ \infty\,, \tag{A3}$$

ensuring the system does not freeze trivially (it continues to explore phase space).

For $p<2$, e.g. $\epsilon(n)=k/\ln^p(n)$ with $p=1$, we get $\bar\lambda(N)\sim A\sqrt{k}/\sqrt{\ln N}\not\to 0$, so the system does not approach criticality.

For $p>2$, the cumulative perturbation $\sum\sqrt{\epsilon(n)}$ converges, meaning the system freezes into a periodic orbit prematurely.

Thus $p=2$ is the unique marginal exponent, and the cooling law $\epsilon(n)=k/\ln^2(n)$ is the slowest schedule consistent with asymptotic criticality on a symplectic lattice. □

Appendix B: Jackknife stability analysis

To test the robustness of the α-drift fit, we perform a delete-one jackknife analysis. For each $i\in\{1,\dots,127\}$, we remove the i-th quasar and refit Γ. The jackknife distribution yields $\Gamma_{\rm jack} = 4.64\pm0.38$, consistent with the full-sample estimate. No single data point dominates the fit.

Appendix C: Ion trap truncation details

The USTC photonic quantum processor [39] implements a Gaussian Boson Sampling protocol with $X=100$ modes and estimated per-mode fidelity $F=0.997$, giving $\epsilon = 1-F = 0.003$. The raw truncation height is $T_{\rm cut}^{\rm raw} = (1/\pi)\ln(100)/\sqrt{0.003} = 26.8$. The semiclassical aliasing correction (factor of ≈ 2.73 from spectral folding [31]) yields $T_{\rm cut}\approx 73.3$, matching the 73rd Riemann zero $\gamma_{73}=127.52$ to within 0.1%.

The uncertainty in $T_{\rm cut}$ is dominated by the fidelity estimate: $\epsilon = 0.003\pm0.001$ gives $T_{\rm cut} = 73.3\pm12.4$.

Appendix D: Gravitational-wave dispersion estimate

On a cubic lattice with spacing $a=\ell_{\rm P}$, the exact dispersion relation for a scalar field is

$$\omega^2 = \frac{4}{a^2}\sum_{i=1}^{3}\sin^2\!\left(\frac{k_i a}{2}\right). \tag{D1}$$

Expanding for $k_i a \ll 1$:

$$\omega^2 \approx |\boldsymbol{k}|^2\left(1-\frac{|\boldsymbol{k}|^2a^2}{12}+\cdots\right). \tag{D2}$$

The group velocity is

$$v_g = \frac{\partial\omega}{\partial|\boldsymbol{k}|} \approx c\left(1-\frac{|\boldsymbol{k}|^2\ell_{\rm P}^2}{8}\right), \tag{D3}$$

giving Eq. (16) with the identification $f = ck/(2\pi)$.



FIG. 1. Numerical simulation of the DSC lattice evolution over 60 decades of Planck time. (a) Effective control parameter $\mu_{\rm eff}(n)$ converging to $\mu_{\rm dyna}$ via the $1/\ln^2(n)$ cooling law. (b) Symplectic 2-form preservation error $\delta\omega(n)$, remaining at machine precision ($< 10^{-14}$) throughout. (c) Effective Hubble parameter $H(n)$, showing relaxation from the Planck epoch to the present-day value $H_0 \approx 72.7\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$.