Integer Quantum Resonance and Antiresonance of the Kicked Rotor: An Exact Parity Sheet

Route-A source-local certificate HCS-C337

3 September 2026

Abstract

We solve the positive-integer resonance sheet of the quantum kicked rotor in a fixed free-after-kick convention. Parity of the integer resonance index gives a complete dichotomy: an exact Bessel propagator with ballistic momentum variance, or a state-independent two-kick antiresonance.

1 Frozen Floquet owner

Let

$$\mathcal{H} = L^2(\mathbb{T}, d\theta/(2\pi)), \qquad |n\rangle(\theta) = e^{\mathrm{i}n\theta}, \qquad \widehat{n} = -\mathrm{i}\partial_\theta.$$

The periodic H^1 realization of $\widehat{n}$ is self-adjoint, and $\widehat{n}^2$ is self-adjoint on periodic H^2. For $\kappa \in \mathbb{R}$ put

$$K_\kappa = e^{-\mathrm{i}\kappa\cos\theta}, \qquad U_\tau = e^{-\mathrm{i}\tau\widehat{n}^2/2} K_\kappa.$$

Here K_κ denotes unit-modulus multiplication. Spectral calculus and bounded multiplication make both factors, hence U_τ, unitary. The ordering is part of the definition: one kick followed by free rotation.

We restrict to

$$\tau = 2\pi\ell, \qquad \ell \in \mathbb{Z}_{>0}. \tag{1}$$

No assertion below concerns another rational sheet or a detuned value.

Lemma 1 (Free-phase parity)**.** *At (1), the free factor is I for even ℓ and the half-turn*

$$(Rf)(\theta) = f(\theta + \pi)$$

for odd ℓ.

Proof. On $|n\rangle$ the free eigenvalue is $e^{-\mathrm{i}\pi\ell n^2}$. Since $n^2 - n = n(n-1)$ is even, this equals $e^{-\mathrm{i}\pi\ell n}$. It is one for even ℓ and $(-1)^n$ for odd ℓ. The latter is precisely the Fourier action of R. □

Theorem 2 (Integer resonance and antiresonance)**.** *Fix $m \in \mathbb{Z}$ and $t \in \mathbb{Z}_{\geq 0}$.*

1. *If ℓ is even, then*
$$\langle n|U_{2\pi\ell}^t|m\rangle = (-\mathrm{i})^{n-m} J_{n-m}(\kappa t). \tag{2}$$
Consequently the momentum law is $J_{n-m}(\kappa t)^2$, its mean is m, and
$$\mathrm{Var}(n) = \frac{\kappa^2 t^2}{2}, \qquad \mathbb{E}_m[\widehat{n}^2/2]_t = \frac{m^2}{2} + \frac{\kappa^2 t^2}{4}. \tag{3}$$
2. *If ℓ is odd, then $U_{2\pi\ell} = RK_\kappa$ and*
$$U_{2\pi\ell}^2 = I. \tag{4}$$
At even t the amplitude is δ_{nm}, while at odd t it is
$$(-1)^n(-\mathrm{i})^{n-m} J_{n-m}(\kappa). \tag{5}$$

Proof. For even ℓ, Lemma 1 gives $U = K_\kappa$ and $U^t = K_{t\kappa}$. The Jacobi–Anger expansion

$$e^{-\mathrm{i}x\cos\theta} = \sum_{q\in\mathbb{Z}} (-\mathrm{i})^q J_q(x) e^{\mathrm{i}q\theta}$$

with $q = n - m$ proves (2). The moment statements follow from Fourier orthogonality:

$$\sum_q J_q(x)^2 e^{\mathrm{i}qu} = J_0(2x\sin(u/2)).$$

Two derivatives at $u = 0$ give mean displacement zero and variance $x^2/2$; translation by m gives (3).

For odd ℓ, the free factor is R. Since the half-turn reverses the cosine,

$$RK_\kappa R = K_{-\kappa} = K_\kappa^{-1}.$$

Thus $(RK_\kappa)^2 = I$. Even powers are I, and odd powers are RK_κ. Applying R after the Jacobi–Anger expansion multiplies the nth Fourier coefficient by $(-1)^n$, which proves (5). □

The phrase *operator parity owner* records the first paper round: the all-time dichotomy is an operator theorem, not a finite-lattice simulation.

References

[1] I. Dana, E. Eisenberg, and N. Shnerb, "Antiresonance and localization in quantum dynamics," *Physical Review E* 54 (1996), 5948–5963. DOI: 10.1103/PhysRevE.54.5948.

[2] I. Dana and D. L. Dorofeev, "General Quantum Resonances of the Kicked Particle," *Physical Review E* 73 (2006), 026206. DOI: 10.1103/PhysRevE.73.026206; arXiv:nlin/0509035.

[3] C. Ryu, M. F. Andersen, A. Vaziri, M. B. d'Arcy, J. M. Grossman, K. Helmerson, and W. D. Phillips, "High-Order Quantum Resonances Observed in a Periodically Kicked Bose–Einstein Condensate," *Physical Review Letters* 96 (2006), 160403. DOI: 10.1103/PhysRevLett.96.160403.