

Exact Finite Termination and Cell Geometry for One-Dimensional Bounded Confidence

Route-A source-local certificate HCS-C312

3 September 2026

Abstract

For the homogeneous one-dimensional Hegselmann–Krause map we prove order and permanent-gap decomposition, exact finite termination within $4n^3 + 2n + 2$ updates, and the complete fixed-cluster classification. We also describe every strict neighbor cell as a rational row-stochastic linear map, retain threshold discontinuities and affine covariance, and give an exact counterexample to mean conservation. An exhaustive 801-system rational archive and independent adversarial lanes audit the theorem without replacing its all-data proof.

1 Order, decomposition, and finite-time theorem

Let $\varepsilon > 0$ and, after sorting the labels, set

$$N_i(t) = \{j : |x_j(t) - x_i(t)| \leq \varepsilon\}, \quad x_i(t+1) = \frac{1}{|N_i(t)|} \sum_{j \in N_i(t)} x_j(t). \quad (1)$$

The inequality is closed: a pair at distance exactly ε interacts.

Theorem 1 (Complete one-dimensional stopping atlas). *For every $n \geq 1$ and every real initial state:*

1. *order, coincident blocks, and the convex hull are preserved;*
2. *a consecutive gap greater than ε never closes and splits the dynamics into independent subsystems;*
3. *the state becomes exactly fixed by time $4n^3 + 2n + 2$;*
4. *a state is fixed iff its distinct occupied positions are separated by gaps strictly greater than ε .*

The key quantitative fact is included for clarity.

Lemma 1 (Two-step progress). *If the system is not fixed at time t , let ℓ be its leftmost nonfrozen position block. By time $t + 2$, that block has gained multiplicity, has frozen, or its position has moved right by at least $\varepsilon/(2n^2)$.*

Proof. Scale to $\varepsilon = 1$ and let r be the first strictly right neighbor of ℓ . If their neighborhoods coincide, their next averages coincide and the left block gains weight. Otherwise r has a right neighbor unseen by ℓ , hence beyond $x_\ell + 1$. Direct averaging over at most n points gives $x_r(t+1) \geq x_\ell(t) + 1/n$. If ℓ itself moved by $1/(2n)$ we are done. Otherwise their new separation is at least $1/(2n)$. If it now exceeds one, ℓ freezes; if not, the next average moves its block by at least another factor $1/n$. Rescaling proves the claim. $\square$

Proof of Theorem 1. The average of two ordered interval neighborhoods is ordered; identical positions have identical neighborhoods. Every update lies in its neighborhood convex hull. If a consecutive gap exceeds ε , the two sides never average across it and their separate convex hulls cannot approach, proving items 1–2.

Decompose the initial state at all such gaps. A component of m agents has width at most $(m-1)\varepsilon$. Apply Lemma 1 at even times. The leftmost active position is nondecreasing; multiplicity/freeze events exhaust at most m labels, while displacement events occur at most $2m^3$. Thus the component stops within $4m^3 + 2m + 2$ updates. Independent components evolve in parallel, so the full stopping time is their maximum; since every $m_j \leq n$, this is at most $4n^3 + 2n + 2$, proving item 3.

Clusters separated by strict gaps are plainly fixed. Conversely, the leftmost occupied position having any distinct neighbor within ε would average strictly to the right, contradicting fixedness. This proves item 4. $\square$

2 Strict cells, covariance, and the missing mean invariant

Away from threshold hyperplanes $|x_i - x_j| = \varepsilon$, the neighbor graph G is locally constant and (1) is

$$x^+ = A_G x, \quad (A_G)_{ij} = |N_i|^{-1} \mathbf{1}_{\{j \in N_i\}}. \quad (2)$$

Thus $A_G \mathbf{1} = \mathbf{1}$, but its columns need not sum to one. The global map is piecewise rational linear and generally discontinuous across a closed threshold face because a new averaging edge is inserted there.

Translation by c commutes with the map. Multiplying both opinions and ε by $s > 0$ multiplies every later state by s . In contrast, the arithmetic mean is not invariant. With $\varepsilon = 1$,

$$(0, 1/2, 7/5) \mapsto (1/4, 19/30, 19/20), \quad 19/30 \mapsto 11/18, \quad (3)$$

a drift of $-1/45$. This is exactly the failure of A_G to be doubly stochastic; it does not conflict with order or convex-hull preservation.

3 Evidence, collisions, and Route-A boundary

All 791 nondecreasing half-integer configurations with $n \leq 5$, plus ten signed, scaled, threshold, duplicate and mean-drift controls, are evolved in exact rational arithmetic. The archive retains 1,843 states and 28,895 leaves; its maximum observed stop is six and 209 cases change mean. An independent implementation performs 28,870 checks, SymPy closes 38 exact identities, replay is byte-identical, and 26 repaired-hash or parser attacks must fail. The finite grid is a regression oracle; Lemma 1 proves the all-real theorem.

C203 uses a fixed signed graph and continuous semigroup, C259 a tree Kuramoto flow, and C301 a random absorbing partition refinement. C312 has a deterministic state-dependent confidence graph and literal finite stopping.

The exact tuple is

$$(A0_FAIL, A1_FAIL, A2_FAIL, A3_FAIL, A4_FAIL).$$

Agent and threshold data have no rational-prime owner (A0); all trajectories eventually fix, leaving no nontrivial primitive cycles (A1); update time is not a logarithmic-prime roof (A2); the rational cell matrices give no target determinant (A3) or self-adjoint target-zero quantization (A4). Route A is rejected and Route B stays locked under NO_BAD_EULER_OR_ROOT_NUMBER. No target Euler factor, root number, automorphy, divisor law, functional equation, or zero match is asserted.

AI use. A generative language model assisted prose and code scaffolding. The reproduced proof, exact independent trajectories, adversarial tests, and deterministic artifacts define the audit record.

Source lineage

References

- [1] A. Bhattacharyya, M. Braverman, B. Chazelle, and H. L. Nguyen, “On the convergence of the Hegselmann–Krause system,” *ITCS 2013*, 61–66. [arXiv:1211.1909](#).