

An Exact Stability, Crossing, and Jordan Atlas for Erlang-2 Distributed-Delay Feedback

Route-A source-local certificate HCS-C343

3 September 2026

Abstract

We solve a scalar negative-feedback equation with a normalized Erlang shape-two memory kernel. Compatible histories admit an exact three-dimensional linear-chain realization, whose cubic Routh determinant gives a necessary-and-sufficient stability threshold and a transverse imaginary crossing.

1 Frozen memory owner

Fix

$$a, b \geq 0, \quad r > 0, \quad K_r(s) = r^2 s e^{-rs} \quad (s \geq 0),$$

and consider

$$\dot{x}(t) = -ax(t) - b \int_0^\infty K_r(s)x(t-s) ds. \quad (1)$$

The kernel is normalized; it has mean $2/r$, variance $2/r^2$, and Laplace transform $r^2/(\lambda + r)^2$. A compatible fading-memory history is one for which the following integrals exist and permit the standard differentiation under the integral:

$$z_1(t) = \int_0^\infty r e^{-rs} x(t-s) ds, \quad z_2(t) = \int_0^\infty K_r(s)x(t-s) ds.$$

Lemma 1 (Exact linear chain). *For compatible initialization, equation (1) is equivalent to*

$$\dot{x} = -ax - bz_2, \quad \dot{z}_1 = r(x - z_1), \quad \dot{z}_2 = r(z_1 - z_2). \quad (2)$$

It is not asserted that arbitrary values of (z_1, z_2) encode a prescribed prehistory.

Proof. Integration by parts gives the two filter equations in (2); the first equation is (1). Conversely, subtract the displayed convolutions from a solution of the two stable filter equations. The differences solve homogeneous first-order equations and vanish initially, so they vanish for all later times. Thus compatible chain solutions recover the delay equation exactly. $\square$

Put $X = (x, z_1, z_2)^\top$. Then $\dot{X} = MX$, where

$$M = \begin{pmatrix} -a & 0 & -b \\ r & -r & 0 \\ 0 & r & -r \end{pmatrix},$$

and direct expansion gives

$$\begin{aligned} p(\lambda) &= \det(\lambda I - M) = (\lambda + a)(\lambda + r)^2 + br^2 \\ &= \lambda^3 + c_2\lambda^2 + c_1\lambda + c_0, \\ c_2 &= a + 2r, \quad c_1 = r(r + 2a), \quad c_0 = r^2(a + b). \end{aligned} \quad (3)$$

Theorem 2 (Sharp stability and imaginary crossing). *For $b > 0$, the zero solution is exponentially stable if and only if*

$$0 < b < b_H, \quad b_H = \frac{2(a + r)^2}{r}. \quad (4)$$

At $b = b_H$,

$$p(\lambda) = (\lambda + a + 2r)(\lambda^2 + r(r + 2a)), \quad (5)$$

so the roots are $-a - 2r$ and $\lambda_\pm = \pm i\omega_H$, with $\omega_H = \sqrt{r(r + 2a)}$. The conjugate pair is simple and crosses from left to right:

$$\operatorname{Re} \lambda'_+(b_H) = \frac{r^2}{2\{r(r + 2a) + (a + 2r)^2\}} > 0. \quad (6)$$

For $b > b_H$, exactly two roots lie in the open right half-plane.

Proof. For $b > 0$, all c_j are positive. The remaining cubic Routh–Hurwitz minor is

$$c_2c_1 - c_0 = r\{2(a + r)^2 - br\}.$$

This proves the equivalence in (4). At equality, multiplication verifies (5). Implicit differentiation of $p(\lambda, b) = 0$ at $\lambda = i\omega_H$ yields (6). Above the threshold the first Routh column has signs $+, +, -, +$, hence two sign changes and exactly two open-right-half-plane roots. $\square$

The phrase *chain equivalence owner* records the first manuscript round: this is an all-parameter analytic theorem, not stability inferred from sampled roots.