

An Exact Finite-Size Bulk–Edge and Quench Atlas for the SSH Chain

Route-A source-local certificate HCS-C318

3 September 2026

Abstract

For the balanced open Su–Schrieffer–Heeger chain of M cells, we derive the full characteristic polynomial and prove that a strictly hyperbolic edge pair exists if and only if $w/v > (M+1)/M$. We give its exact splitting, two-ended eigenvectors, decay bound, critical linear taper, and fixed-ratio asymptotic. We then separate this finite threshold from the periodic bulk transition, resolve finite-ring parity and every hopping face, and derive an entire matrix-sinc propagator valid at singular couplings. An exact mode-quench corollary retains the finite momentum-grid caveat; independent symbolic, numerical, replay, and hostile tests define the claim boundary.

1 Finite hyperbolic edge threshold

Let $M \geq 2$ and $v, w \geq 0$. In the ordered basis $(A_1, \dots, A_M, B_1, \dots, B_M)$, the open Hamiltonian is

$$H_O = \begin{pmatrix} 0 & T \\ T^* & 0 \end{pmatrix}, \quad T = vI + wS_-, \quad (S_-)_{j,j-1} = 1. \quad (1)$$

Thus $\Gamma = \text{diag}(I, -I)$ obeys $\Gamma H_O \Gamma = -H_O$.

Theorem 1 (Characteristic and sharp finite edge theorem). *Put $q_M(y) = \det(yI - TT^*)$. Then*

$$q_0 = 1, \quad q_1 = y - v^2, \quad q_m = (y - v^2 - w^2)q_{m-1} - v^2w^2q_{m-2}. \quad (2)$$

For $v, w > 0$, set $x = (E^2 - v^2 - w^2)/(2vw)$. With U_m the Chebyshev polynomial of the second kind,

$$\det(EI - H_O) = (vw)^M \left[U_M(x) + \frac{w}{v} U_{M-1}(x) \right]. \quad (3)$$

There is exactly one secular root $x = -\cosh \kappa < -1$, and hence exactly one eigenvalue pair $\pm E_e$ outside the trigonometric sector, if and only if

$$\frac{w}{v} > \frac{M+1}{M}. \quad (4)$$

For the unique $\kappa > 0$ determined by

$$\frac{w}{v} = \frac{\sinh((M+1)\kappa)}{\sinh(M\kappa)}, \quad (5)$$

the positive energy and one unnormalised positive-energy vector are

$$E_e = v \frac{\sinh \kappa}{\sinh(M\kappa)} = w \frac{\sinh \kappa}{\sinh((M+1)\kappa)}, \quad (6)$$

$$a_j = (-1)^{j-1} \sinh((M+1-j)\kappa), \quad b_j = (-1)^{j-1} \sinh(j\kappa). \quad (7)$$

Thus (a, b) has energy E_e and $(a, -b)$ has energy $-E_e$. The profiles decay strictly inward:

$$\frac{|a_{j+1}|}{|a_j|} < e^{-\kappa}, \quad \frac{|b_j|}{|b_{j+1}|} < e^{-\kappa} \quad (1 \leq j < M). \quad (8)$$

At equality in (4), $x = -1$ and $E = v/M$. After common division of a, b by κ (equivalently, after overall normalisation), the $\kappa \downarrow 0$ profiles are $a_j = (-1)^{j-1}(M+1-j)$ and $b_j = (-1)^{j-1}j$.

Proof. The leading diagonal entry of TT^* is v^2 ; every subsequent one is $v^2 + w^2$, and the off-diagonal entries are vw . Its continuant is (2). Normalising this recurrence gives (3); in particular, the factor $(vw)^M$ cannot be dropped. The M roots are real and simple because TT^* is an irreducible Jacobi matrix. Also $\det T = v^M$, so $E = 0$ is impossible when $v > 0$.

For $x = -\cosh \kappa$, the identity $U_m(-\cosh \kappa) = (-1)^m \sinh((m+1)\kappa)/\sinh \kappa$ reduces the secular equation to (5). Its logarithmic derivative is

$$(M+1) \coth((M+1)\kappa) - M \coth(M\kappa) > 0,$$

because $s \mapsto s \coth(sk)$ is strictly increasing. The ratio therefore rises from $(M+1)/M$ to infinity. No root lies above $x = 1$, where both Chebyshev terms are positive, so this proves the iff statement and uniqueness. Direct hyperbolic addition verifies $Tb = E_e a$ and $T^*a = E_e b$, including both expressions in (6). Finally,

$$\frac{\sinh(n\kappa)}{\sinh((n+1)\kappa)} = e^{-\kappa} \frac{1 - e^{-2n\kappa}}{1 - e^{-2(n+1)\kappa}} < e^{-\kappa},$$

which proves (8). Taking $\kappa \downarrow 0$ gives the threshold energy; applying the common $1/\kappa$ rescaling before that limit gives the nonzero tapers. $\square$

Corollary 2 (Finite splitting). *For fixed $r = w/v > 1$ and all sufficiently large M , the hyperbolic root exists, $\kappa_M \rightarrow \log r$, and*

$$E_e \sim w(1 - r^{-2})r^{-M}. \quad (9)$$

Consequently the finite pair is never exactly zero for $v > 0$, although its splitting is exponentially small.

Proof. Writing $z = e^{-\kappa}$ turns (5) into $r = z^{-1}(1 - z^{2M+2})/(1 - z^{2M})$, hence $z \rightarrow r^{-1}$. Equation (6) becomes $E_e = vz^{M-1}(1 - z^2)/(1 - z^{2M})$, which gives (9). $\square$

2 Bulk winding, boundary faces, and unitary propagation

Close the chain and use momenta $k_n = 2\pi n/M$. Our orientation convention is

$$q(k) = v + we^{ik}, \quad h(k) = \begin{pmatrix} 0 & \overline{q(k)} \\ q(k) & 0 \end{pmatrix}. \quad (10)$$

Theorem 3 (Periodic bulk and finite parity). *The Bloch energies are*

$$E_{\pm}(k) = \pm \sqrt{v^2 + w^2 + 2vw \cos k}. \quad (11)$$

For the continuum symbol and the sampled M -cell ring, respectively, let Δ_{∞} and Δ_M denote the minimum positive-band distance to zero. Then

$$\Delta_{\infty} = |v - w|, \quad \Delta_M = \begin{cases} |v - w|, & M \text{ even}, \\ \sqrt{v^2 + w^2 - 2vw \cos(\pi/M)}, & M \text{ odd}. \end{cases} \quad (12)$$

The corresponding central band gaps are $2\Delta_{\infty}$ and $2\Delta_M$. Whenever the loop avoids zero, its counterclockwise winding is 1 for $w > v$ and 0 for $v > w$; at $v = w > 0$ it is undefined. At this critical line a finite ring has a two-dimensional zero sector if and only if M is even. For odd M it instead has $\Delta_M = 2v \sin(\pi/(2M)) > 0$.

Proof. Squaring (10) gives $|q(k)|^2 I$, proving the dispersion. Over the continuum, $\cos k$ is minimized at $k = \pi$. The finite grid samples that point exactly for even M ; for odd M its nearest points are $\pi \pm \pi/M$, which proves (12). The curve $q(k)$ is the counterclockwise circle of radius w centred at v . It encloses zero exactly when $w > v$. On the critical line it reaches zero only at $k = \pi$, which belongs to the finite grid exactly for even M ; the corresponding 2×2 fiber is zero. $\square$

Theorems 1 and 3 exhibit a finite separation: the bulk is topological for $w/v > 1$, but the open pair is strictly hyperbolic only for $w/v > (M+1)/M$. Equality in the latter condition is a band-edge taper, not an exponentially localized state.

Theorem 4 (All hopping faces). *For the open chain the following statements are exact.*

1. *If $w = 0 < v$, there are M dimers and $\pm v$ each has multiplicity M .*

2. If $v = 0 < w$, there are $M - 1$ dimers, $\pm w$ each has multiplicity $M - 1$, and A_1, B_M span a two-dimensional zero sector.
3. If $v = w = 0$, the kernel has dimension $2M$.
4. If $v = w > 0$, then $\text{spec}(H_O) = \{2v \cos(\ell\pi/(2M + 1)) : 1 \leq \ell \leq 2M\}$, so zero is absent.

For $M = 1$, declared separately, the open intercell bond is absent and the energies are $\pm v$; the periodic wrap merges with the intracell bond and the energies are $\pm(v + w)$.

Proof. The first three statements follow by listing the isolated sites and bonds. The fourth is the sine diagonalisation of the uniform path on $2M$ sites; its denominator $2M + 1$ prevents a zero cosine. The last statement follows from the two explicit 2×2 matrices. $\square$

Theorem 5 (Entire boundary-safe propagator). *Let $\text{sinc } z = \sin(z)/z$ with $\text{sinc } 0 = 1$. Entire matrix functional calculus gives, on every face in Theorem 4,*

$$e^{-itH_O} = \begin{pmatrix} \cos(t\sqrt{TT^*}) & -it \text{sinc}(t\sqrt{TT^*})T \\ -it \text{sinc}(t\sqrt{T^*T})T^* & \cos(t\sqrt{T^*T}) \end{pmatrix}. \quad (13)$$

It is unitary and satisfies $\Gamma e^{-itH_O} \Gamma = e^{itH_O} = (e^{-itH_O})^*$.

Proof. The even and odd powers have block forms

$$H_O^{2n} = \text{diag}((TT^*)^n, (T^*T)^n), \quad H_O^{2n+1} = \begin{pmatrix} 0 & (TT^*)^n T \\ (T^*T)^n T^* & 0 \end{pmatrix}.$$

Separating the exponential series proves (13). Because cosine and sinc are entire in the squared operators, no inverse or positive singular-value assumption enters. Unitarity follows from self-adjointness, and the last identity from chirality. $\square$

3 Quench corollary, evidence, and Route-A boundary

Consider strictly positive, gapped initial and final hoppings. Let $q_i(k), q_f(k)$ use (10), and start in the lower initial Bloch state.

Corollary 6 (Exact mode zero with finite-grid caveat). *Its final-evolution amplitude is*

$$g_k(t) = \cos(E_f(k)t) + ic(k) \sin(E_f(k)t), \quad c(k) = \frac{\text{Re}(\overline{q_i(k)} q_f(k))}{|q_i(k)| |q_f(k)|}. \quad (14)$$

A continuum mode vanishes at a real time if and only if

$$(v_i - w_i)(v_f - w_f) < 0. \quad (15)$$

In that case

$$\cos k_* = -\frac{v_i v_f + w_i w_f}{v_i w_f + w_i v_f}, \quad t_n = \frac{\pi/2 + n\pi}{E_f(k_*)}. \quad (16)$$

A finite ring has such an exact mode zero only if a sampled $k_m = 2\pi m/M$ has this cosine.

Proof. Pauli-matrix exponentiation gives (14). A complex zero requires both $\cos(E_f t) = 0$ and $c(k) = 0$. The numerator of c is

$$A + B \cos k, \quad A = v_i v_f + w_i w_f, \quad B = v_i w_f + w_i v_f > 0.$$

It has a root in $[-1, 1]$ exactly when $A < B$, since $A, B > 0$; but $A - B = (v_i - w_i)(v_f - w_f)$. Gapped endpoints make the inequality strict, giving (15)–(16). A finite Fourier decomposition contains only its sampled momenta, proving the last sentence. $\square$

For example, the quench $(v_i, w_i) = (3, 1)$ to $(v_f, w_f) = (1, 5)$ has $\cos k_* = -1/2$. In the audited range, and in general, the finite ring hits that value exactly when 3 divides M . Opposite bulk phases alone do not guarantee an exact zero for an arbitrary finite ring. This is a single-particle mode statement, not a many-body dynamical-phase-transition claim.

Evidence. The content-addressed evidence has 55 open-polynomial rows, 33 rational hyperbolic witnesses, 11 threshold rows, 70 periodic rows containing 595 momentum cells, 33 boundary rows, 30 propagator rows, and six quench rows: 7,161 audited scalar leaves. A producer-independent checker performs 10,948 checks using exact Sturm sequences and a separate matrix exponential. SymPy closes 9,181 identities, two isolated runs replay byte for byte, and 53 repaired-hash/parser mutations must fail. These finite grids are regressions; Theorems 1–5 and Corollary 6 have the stated analytic domains.

Collision boundary. C308 studies a one-site non-Hermitian nonreciprocal Hatano–Nelson skin chain and excludes topology. The present object is balanced, Hermitian, bipartite, and chiral; its distinguishing theorem is the strict finite bulk–edge threshold separation. C267 is an infinite uniform-field Wannier–Stark ladder, C297 a two-site PT dimer, and C138 a metric-graph flux winding problem. No resolvent, pseudospectral, or skin-effect claim is reused here.

Route-A result and nonclaims. Under NO_BAD_EULER_OR_ROOT_NUMBER, the strict tuple is

(A0_FAIL, A1_FAIL, A2_FAIL, A3_FAIL, A4_NATURAL_QUANTIZATION).

The cells and Bloch momentum carry no rational-prime payload (A0); the momentum loop is not a primitive-orbit ledger (A1); the finite determinant is not an Euler product (A2); and no target functional equation, counting law, or divisor appears (A3). The finite Hermitian matrix is already a natural source quantization (A4), but its lattice levels are not target zeros and it is not a Hilbert–Pólya operator. Route A is rejected and Route B remains locked. No target Euler factor, root number, automorphy, zero correspondence, disorder, interaction, or self-consistent phonon claim is made.

AI use. A generative language model assisted drafting and code scaffolding. The displayed derivations, independent recomputation, adversarial tests, and deterministic artifacts define the audit record.

Source lineage

References

- [1] W. P. Su, J. R. Schrieffer, and A. J. Heeger, “Solitons in Polyacetylene,” *Physical Review Letters* 42 (1979), 1698–1701. DOI: [10.1103/PhysRevLett.42.1698](https://doi.org/10.1103/PhysRevLett.42.1698).
- [2] W. P. Su, J. R. Schrieffer, and A. J. Heeger, “Soliton excitations in polyacetylene,” *Physical Review B* 22 (1980), 2099–2111. DOI: [10.1103/PhysRevB.22.2099](https://doi.org/10.1103/PhysRevB.22.2099).
- [3] J. K. Asbóth, L. Oroszlány, and A. Pályi, “The Su–Schrieffer–Heeger Model,” in *A Short Course on Topological Insulators* (Springer, 2016). DOI: [10.1007/978-3-319-25607-8_1](https://doi.org/10.1007/978-3-319-25607-8_1).