

SYNCHRONOUS MEX DYNAMICS ON COMPLETE MULTIPARTITE GRAPHS: EXACT FIBRES, TWO-CYCLES, AND BASINS

ANONYMOUS

ABSTRACT. Every vertex of a finite graph is updated simultaneously to the least nonnegative colour absent from its open neighbourhood. On $K_{a_1, \dots, a_k}$, the first image is constant on each part and induces $T(y)_i = \text{mex}\{y_j : j \neq i\}$. We retain the labelled information discarded by this quotient through the exact first-image fibre $N_{\mathbf{a}, q}(y)$. One formula uses inclusion–exclusion over missing lower colours; a materially different formula uses the part support of every named colour and multivariate labelled generating functions.

The quotient has exactly $k!$ fixed states and

$$b_k = \sum_{m=0}^{k-2} \frac{k!}{(k-m)!}$$

two-cycles. More strongly, a repeated coordinate in a graph image can only be its maximum; this forces the second iterate of every labelled colouring to be recurrent. Closed products give the fibre of every recurrent target. Together with explicit quotient preimages they yield every orbit basin and its depth-zero, one, and two layers. The 2003 serial-daemon mex correction, Grundy fixed points, generic labelled-EGF machinery, and zeta bookkeeping receive zero contribution credit. The residual scope is only this synchronous multipartite conjunction. Ownership, priority, and external circulation remain on hold.

1. INTRODUCTION

The local instruction *take the least colour absent from the open neighbourhood* is classical in Grundy colouring. In particular, Hedetniemi et al. [5] use the same value, shifted to positive colours, in self-stabilizing algorithms controlled by a daemon. Later parallel, asynchronous, local-search, and dynamic-colouring work includes [2, 3]. The local correction, Grundy fixed points, and scheduler-dependent algorithmic convergence therefore receive zero contribution credit here.

We instead fix an unconditional Jacobi clock: every vertex updates on every round from the unchanged old colouring. This scheduler does not generally converge; even the uniform state can alternate. Complete multipartite graphs give a tractable family, because vertices in one part have identical open neighbourhoods. Yet replacing each part by one coordinate erases the ordered size vector $\mathbf{a} = (a_1, \dots, a_k)$.

The key device is to retain the weight of every first-round quotient state. For a part vector y , let $N_{\mathbf{a}, q}(y)$ be the number of labelled vertex colourings whose first image is y . We derive this weight twice. Inclusion–exclusion counts violated presence requirements. A second proof fixes, for every named colour, the set of parts in which it occurs, and counts onto assignments. Their equality is a useful exact bridge rather than an algorithmic complexity claim.

The temporal classification then has three steps. An elementary identity for the quotient keeps unique colours below a global mex and replaces all other coordinates by that mex. A separate image-shape lemma says that only the maximum first-image value can repeat. Together they put the second graph iterate directly in an explicit fixed/two-cycle list. Finally, closed recurrent fibres and quotient preimages turn the classification into all basin and layer counts.

2020 *Mathematics Subject Classification*. 05C15, 37P35, 05A18.

Key words and phrases. finite dynamics, mex, Grundy colouring, complete multipartite graph, functional graph, fibre, basin.

We do not treat arbitrary graphs or asynchronous schedules. Conflict guards and the closed-neighbourhood variant are also outside our scope. A bounded owner search did not locate this exact synchronous multipartite functional graph, but a search non-hit is not novelty or priority evidence. External dissemination remains **HOLD**.

2. THE GRAPH MAP AND ITS QUOTIENT

Let $k \geq 1$, let $a_i \geq 1$, and let

$$G = K_{a_1, \dots, a_k}, \quad V(G) = V_1 \sqcup \dots \sqcup V_k, \quad |V_i| = a_i.$$

All vertices are labelled. Put $n = \sum_i a_i$ and

$$\Delta = n - \min_i a_i.$$

For an integer $q \geq \Delta + 1$, let $C_q = \{0, \dots, q-1\}$. The canonical phase uses $q = \Delta + 1$; the larger-palette extension will make surplus-colour dependence explicit.

Definition 2.1 (synchronous open-neighbourhood mex). For $c : V(G) \rightarrow C_q$, define

$$\Phi(c)(v) = \text{mex}\{c(u) : uv \in E(G)\}.$$

Every value on the right is read from c , and all vertices update simultaneously.

Every neighbourhood has at most $\Delta < q$ vertices, so its mex belongs to C_q . Vertices in one part have the same open neighbourhood. Hence $\Phi(c)$ is part-monochromatic, represented by

$$y(c) = (y_1(c), \dots, y_k(c)), \quad y_i(c) = \text{mex } c(V(G) \setminus V_i).$$

Subsequent rounds are governed by

$$(1) \quad T(y)_i = \text{mex}\{y_j : j \neq i\}.$$

Thus $\Phi^t(c)$ is represented by $T^{t-1}(y(c))$ for $t \geq 1$. Define the exact first-image fibre

$$N_{\mathbf{a},q}(y) = |\{c \in C_q^{V(G)} : y(c) = y\}|.$$

3. TWO EXACT FIBRE FORMULAS

3.1. Bad-event inclusion–exclusion. For $y \in C_q^k$, put

$$\mathcal{E}(y) = \{(i, r) : 1 \leq i \leq k, 0 \leq r < y_i\}.$$

For $J \subseteq \mathcal{E}(y)$ and $1 \leq h \leq k$, define the set

$$B_h(y, J) = \{y_i : i \neq h\} \cup \{r : (i, r) \in J, i \neq h\}.$$

Repeated values in this union are counted once.

Theorem 3.1 (inclusion–exclusion fibre). *For every $y \in C_q^k$,*

$$(2) \quad N_{\mathbf{a},q}(y) = \sum_{J \subseteq \mathcal{E}(y)} (-1)^{|J|} \prod_{h=1}^k (q - |B_h(y, J)|)^{a_h}.$$

Impossible targets automatically have value zero.

Proof. The equality $y_i = \text{mex } c(V \setminus V_i)$ requires colour y_i to be absent outside V_i , and every $r < y_i$ to occur there. Impose all the absence requirements first. For $(i, r) \in \mathcal{E}(y)$, let the bad event be that r is also absent outside V_i . Under a selected set J of bad events, a vertex in V_h cannot use y_i for $i \neq h$ and cannot use r when $(i, r) \in J$ with $i \neq h$. Its forbidden set is exactly $B_h(y, J)$. Vertex choices now factor by parts, and inclusion–exclusion over J gives (2). $\square$

3.2. Colour supports and labelled EGFs. For a named colour r , let $P_r \subseteq [k]$ be the set of parts in which it occurs. Define $\mathcal{A}_r(y)$ to consist of the supports $P \subseteq [k]$ satisfying

$$(3) \quad y_i = r \implies P \subseteq \{i\},$$

$$(4) \quad y_i > r \implies P \not\subseteq \{i\}.$$

Let $s(a, d) = d!S(a, d)$ be the number of onto maps from an a -element labelled set to d named colours, with $s(a, 0) = 0$ for $a \geq 1$. Put

$$G_r^y(x_1, \dots, x_k) = \sum_{P \in \mathcal{A}_r(y)} \prod_{i \in P} (e^{x_i} - 1).$$

Theorem 3.2 (support and EGF fibre). *For every $y \in C_q^k$,*

$$(5) \quad N_{\mathbf{a}, q}(y) = \left(\prod_{i=1}^k a_i! \right) [x_1^{a_1} \cdots x_k^{a_k}] \prod_{r=0}^{q-1} G_r^y(x_1, \dots, x_k),$$

$$(6) \quad = \sum_{\substack{P_r \in \mathcal{A}_r(y) \\ 0 \leq r < q}} \prod_{i=1}^k s(a_i, d_i), \quad d_i = |\{r : i \in P_r\}|.$$

If $M = \max_i y_i$, the factors with $r > M$ combine to

$$\exp((q-1-M)(x_1 + \cdots + x_k)).$$

Proof. Conditions (3)–(4) are exactly the absence and lower-colour presence conditions for each target coordinate. After all supports are fixed, part V_i uses d_i named colours, each at least once, giving $s(a_i, d_i)$. Multiplication over parts and summation over supports proves (6).

In labelled-EGF form, one named colour with support P contributes $\prod_{i \in P} (e^{x_i} - 1)$. Multiplying over named colours and extracting all labelled vertices proves (5); see [4] for the generic labelled construction. For $r > M$, every support is allowed, and

$$\sum_{P \subseteq [k]} \prod_{i \in P} (e^{x_i} - 1) = e^{x_1 + \cdots + x_k},$$

which gives the last assertion. $\square$

4. THE RECURRENT QUOTIENT AND TWO-ROUND COLLAPSE

For $y \in C_q^k$, let $g = \text{mex}\{y_1, \dots, y_k\}$.

Lemma 4.1 (quotient support reduction).

$$T(y)_i = \begin{cases} y_i, & y_i < g \text{ and } y_i \text{ occurs exactly once in } y, \\ g, & \text{otherwise.} \end{cases}$$

Proof. Every value below g occurs in y , while g does not. Removing coordinate i creates a missing value below g exactly when $y_i < g$ and that value was unique. Otherwise all lower values remain and g remains missing. $\square$

Write $[m]_0 = \{0, \dots, m-1\}$. For $0 \leq m \leq k-2$ and an injection $\iota : [m]_0 \hookrightarrow [k]$, define

$$x_{\iota, m}^\pm(i) = \begin{cases} r, & i = \iota(r), \\ m, & i \notin \text{im } \iota \text{ for } x^-, \\ m+1, & i \notin \text{im } \iota \text{ for } x^+. \end{cases}$$

The last two lines mean that the uninjected coordinates are filled by m in x^- and by $m+1$ in x^+ .

Theorem 4.2 (complete quotient recurrence). *The recurrent states of T are exactly:*

- (1) the $k!$ permutations of $0, \dots, k-1$, all fixed; and
- (2) the pairs $x_{\iota, m}^- \leftrightarrow x_{\iota, m}^+$ above.

Every quotient state enters this list in at most two rounds. The number of two-cycles is

$$b_k = \sum_{m=0}^{k-2} \frac{k!}{(k-m)!},$$

with an empty sum for $k = 1$.

Proof. Lemma 4.1 shows that one quotient update retains some unique values below g and fills every other coordinate by g . Let m be the least nonnegative integer missing from the retained unique values. A second use of the lemma retains precisely the initial segment $0, \dots, m-1$ and fills all other coordinates by m , unless only one coordinate remains, in which case the result is a permutation. Direct substitution shows that every permutation is fixed and that each displayed pair is swapped. This proves exhaustion. For fixed m , the number of injections is $k!/(k-m)!$. $\square$

Graph images satisfy a stronger shape restriction.

Lemma 4.3 (only the maximum image value repeats). *Assume $k \geq 2$. If $y_i(c) = y_j(c) = r$ for $i \neq j$, then $r = \max_{\ell} y_{\ell}(c)$.*

Proof. The target condition at i makes colour r absent from $V \setminus V_i$; the condition at j makes it absent from $V \setminus V_j$. Those two sets cover all vertices, so r is globally absent. A target larger than r would require r to occur in its open neighbourhood, a contradiction. $\square$

Theorem 4.4 (two-round global recurrence). *For every labelled colouring c ,*

$$T(y(c)) \in \mathcal{R}_k,$$

where $\mathcal{R}_k$ is the recurrent list in theorem 4.2. Equivalently, $\Phi^2(c)$ is recurrent, so every original-state preperiod is at most two.

Proof. Put $M = \max_i y_i(c)$ and $g = \text{mex}\{y_1(c), \dots, y_k(c)\}$. If $g \leq M$, lemma 4.3 makes every value below g unique. Lemma 4.1 therefore retains one copy of $0, \dots, g-1$ and fills the remaining coordinates by g , giving a permutation or $x_{\iota, g}^-$.

If $g = M+1$, every value $0, \dots, M$ occurs. Lemma 4.3 makes $0, \dots, M-1$ unique. If M is unique, $y(c)$ is a fixed permutation. If it repeats, the quotient update retains the lower initial segment and replaces every M by $M+1$, giving $x_{\iota, M}^+$. $\square$

Every periodic graph colouring belongs to the image of Φ , hence is part-monochromatic. The converse is immediate from the quotient. Thus the graph map has

$$R_k = k! + 2b_k$$

recurrent states. Standard finite-cycle bookkeeping [1] gives:

Corollary 4.5 (zeta function).

$$\zeta_{\Phi}(z) = (1-z)^{-k!}(1-z^2)^{-b_k}.$$

5. CLOSED RECURRENT FIBRES

The general fibre formulas simplify on every recurrent target. Empty products equal one, and $0^0 = 1$ has its combinatorial meaning.

Proposition 5.1 (recurrent-target products). (i) *Let x be a fixed permutation and let i_r be the part with $x_{i_r} = r$. With $h = q - k$,*

$$(5.1) \quad N_{\mathbf{a}, q}(x) = \left[\prod_{r=0}^{k-2} ((h+1)^{a_{i_r}} - h^{a_{i_r}}) \right] (h+1)^{a_{i_{k-1}}}.$$

(ii) *For $x_{\iota, m}^-$, put $R = [k] \setminus \text{im } \iota$, $A_R = \sum_{j \in R} a_j$, $i_r = \iota(r)$, and $v = q - m - 1$. Then*

$$(5.2) \quad N_{\mathbf{a}, q}(x_{\iota, m}^-) = \left[\prod_{r=0}^{m-1} ((v+1)^{a_{i_r}} - v^{a_{i_r}}) \right] v^{A_R}.$$

(iii) For $x_{\iota, m}^+$, put $u = q - m - 2$ and

$$\begin{aligned} L_1 &= \prod_{r=0}^{m-1} ((u+2)^{a_{ir}} - (u+1)^{a_{ir}}), \\ L_0 &= \prod_{r=0}^{m-1} ((u+1)^{a_{ir}} - u^{a_{ir}}), \\ Q &= (u+1)^{A_R}, \\ Q_{\geq 2} &= Q - u^{A_R} - \sum_{j \in R} ((u+1)^{a_j} - u^{a_j}) u^{A_R - a_j}. \end{aligned}$$

Then

$$(5.3) \quad N_{\mathbf{a}, q}(x_{\iota, m}^+) = (L_1 - L_0)Q + L_0 Q_{\geq 2}.$$

Proof. For a fixed permutation, low colour r is confined to V_{i_r} . It is required there for $r < k-1$, while colour $k-1$ is optional; the h larger colours are free. This gives (5.1).

For x^- , every low colour is confined to and required in its injected part. The repeated target m makes colour m globally absent, and the v larger colours are free. This gives (5.2).

For x^+ , colour $m+1$ is globally absent. If colour m occurs in an injected low part, it automatically occurs outside every part in R ; this contributes $(L_1 - L_0)Q$. If it occurs in no low part, it must occur in at least two distinct R -parts. The expression $Q_{\geq 2}$ subtracts from all R -assignments the cases where colour m is absent or occurs in exactly one part. This gives (5.3). $\square$

6. EVERY BASIN AND DEPTH LAYER

The quotient preimages needed below are also explicit.

Proposition 6.1 (recurrent quotient preimages). *For a fixed permutation x with $x_{i_r} = r$,*

$$(6.1) \quad T^{-1}(x) = \{y : y_{i_r} = r \ (r < k-1), \ y_{i_{k-1}} \in \{k-1, \dots, q-1\}\}.$$

For a two-cycle indexed by (ι, m) ,

$$(6.2) \quad T^{-1}(x_{\iota, m}^-) = \{y : y_{i_r} = r \ (r < m), \ y_j \in \{m+1, \dots, q-1\} \ (j \in R)\},$$

$$(6.3) \quad \begin{aligned} T^{-1}(x_{\iota, m}^+) &= \{y : y_{i_r} = r \ (r < m), \ y_j \in \{m\} \cup \{m+2, \dots, q-1\} \ (j \in R), \\ &\quad |\{j \in R : y_j = m\}| \geq 2\}. \end{aligned}$$

Proof. Apply lemma 4.1. To output a fixed permutation, every value below $k-1$ must be unique at its prescribed coordinate, while the last coordinate can carry any value at least $k-1$. To output x^- , the prescribed low values are unique and all other coordinates lie strictly above m . To output x^+ , the global mex is $m+1$; the low values are prescribed, $m+1$ is absent, and m must repeat. $\square$

Put

$$S_{\mathbf{a}, q} = \sum_{x \in \mathcal{R}_k} N_{\mathbf{a}, q}(x), \quad H_x = \sum_{\substack{y \in C_q^k \\ T(y) = x}} N_{\mathbf{a}, q}(y).$$

The sums defining H_x use the finite sets in proposition 6.1; impossible first images simply have fibre zero.

Theorem 6.2 (global and orbitwise layers). *The exact numbers of labelled graph colourings at depths zero, one, and two are*

$$(6.4) \quad \boxed{D_0 = R_k, \quad D_1 = S_{\mathbf{a}, q} - R_k, \quad D_2 = q^n - S_{\mathbf{a}, q}.}$$

TABLE 1. Canonical $K_{1,2}$ signal with $q = 3$. The unequal fixed fibres retain the ordered part sizes after quotient collapse.

recurrent orbit	recurrent target fibres	basin size	layer profile
fixed $(0, 1)$	$N(0, 1) = 4$	4	$(1, 3, 0)$
fixed $(1, 0)$	$N(1, 0) = 6$	10	$(1, 5, 4)$
$(0, 0) \leftrightarrow (1, 1)$	8, 3	13	$(2, 9, 2)$
all orbits		27	$(4, 17, 6)$

If $\mathcal{O} \subseteq \mathcal{R}_k$ is one fixed orbit or one two-cycle, then its basin size and its three layers are

$$(6.5) \quad B_{\mathcal{O}} = \sum_{x \in \mathcal{O}} H_x,$$

$$(6.6) \quad B_{\mathcal{O},0} = |\mathcal{O}|,$$

$$(6.7) \quad B_{\mathcal{O},1} = \sum_{x \in \mathcal{O}} N_{\mathbf{a},q}(x) - |\mathcal{O}|,$$

$$(6.8) \quad B_{\mathcal{O},2} = B_{\mathcal{O}} - \sum_{x \in \mathcal{O}} N_{\mathbf{a},q}(x).$$

Proof. A state has depth at most one exactly when its first image belongs to $\mathcal{R}_k$; summing its fibres gives $S_{\mathbf{a},q}$. Removing the R_k recurrent states gives D_1 , and [theorem 4.4](#) puts every remaining state at depth two.

Moreover, $\Phi^2(c) = x$ exactly when $T(y(c)) = x$. Hence H_x counts the states whose second image is x , and summing over one recurrent orbit gives its whole basin. Its recurrent states form layer zero. Its states of depth at most one are precisely the fibres over its recurrent targets. Subtracting successively proves the displayed formulas. $\square$

7. CONTROLS, CREDIT BOUNDARY, AND CONCLUSION

The deterministic verifier enumerates every labelled colouring in fifteen parameter lanes. It independently compares literal fibres, [theorem 3.1](#), support/onto counts from [theorem 3.2](#), the products in [proposition 5.1](#), all quotient preimages, pointwise depths, and every basin layer. The grid includes $K_{1,2,3}$ with 46,656 states, $K_{2,2,2}$ with 15,625, enlarged palettes, and $k = 1$. A fresh run passes 201,922 exact assertions. Computation is control evidence, not a proof of the parameterized statements.

The closest located rule-level owner is Hedetniemi et al. [5]: the local mex correction and Grundy fixed-point interpretation receive full prior credit. Generic labelled generating functions [4] and finite-map zeta products [1] also receive zero credit. The residual is only the synchronous complete-multipartite conjunction: two exact part-sensitive fibre formulas, the repeated-maximum image shape, two-round recurrence, closed recurrent fibres, and all basin layers.

No polynomial-time evaluation claim is made; inclusion–exclusion can be exponential. No arbitrary-graph or asynchronous conclusion is implied. The owner search remains bounded, and external ownership, novelty, priority, and dissemination remain on hold.

REFERENCES

- [1] Michael Artin and Barry Mazur. On periodic points. *Annals of Mathematics*, 81(1):82–99, 1965. doi: 10.2307/1970384.
- [2] Jakob Bossek, Frank Neumann, Pan Peng, and Dirk Sudholt. Time complexity analysis of randomized search heuristics for the dynamic graph coloring problem. *Algorithmica*, 83(10):3148–3179, 2021. doi: 10.1007/s00453-021-00838-3.

- [3] Jesun Sahariar Firoz, Marcin Zalewski, and Andrew Lumsdaine. A synchronization-avoiding distance-1 grundy coloring algorithm for power-law graphs. In *2019 28th International Conference on Parallel Architectures and Compilation Techniques*, pages 421–432, 2019. doi: 10.1109/PACT.2019.00040.
- [4] Philippe Flajolet and Robert Sedgewick. *Analytic Combinatorics*. Cambridge University Press, 2009. doi: 10.1017/CBO9780511801655.
- [5] Stephen T. Hedetniemi, David P. Jacobs, and Pradip K. Srimani. Linear time self-stabilizing colorings. *Information Processing Letters*, 87(5):251–255, 2003. doi: 10.1016/S0020-0190(03)00299-0.