

Cut and First-Conjugate Geometry of the Standard Heisenberg Group: A Complete Hamiltonian and Distance Atlas

HCS Research Program

1 September 2026

Abstract

For the standard real Heisenberg group we integrate every unit-speed normal sub-Riemannian geodesic and close the global minimizing geometry, rather than stopping at an explicit orbit formula. We exclude nonconstant abnormal extremals, compute the exponential Jacobian, prove equality of the first cut, conjugate, and rotational Maxwell times, and determine the cut locus from the identity. A strictly monotone implicit angle gives the Carnot–Carathéodory distance, including both degenerate endpoint faces. All signs and factors are frozen to one frame convention; no extension to arbitrary Carnot groups is claimed.

1 Frozen model

On $H^1 = \mathbb{R}^3$ put

$$X = \partial_x - \frac{y}{2}\partial_z, \quad Y = \partial_y + \frac{x}{2}\partial_z, \quad [X, Y] = \partial_z,$$

and make X, Y orthonormal. The normal Hamiltonian is

$$H = \frac{1}{2} \left(p_x - \frac{y}{2}p_z \right)^2 + \frac{1}{2} \left(p_y + \frac{x}{2}p_z \right)^2.$$

Write $\lambda = p_z$ and impose $H = 1/2$.

Theorem 1 (Complete identity-based atlas). *Every unit-speed normal geodesic from the identity has an angle ϕ and is, when $\lambda = 0$,*

$$(x, y, z) = (t \cos \phi, t \sin \phi, 0),$$

while for $\lambda \neq 0$ it is

$$\begin{aligned} x &= \frac{\sin(\phi + \lambda t) - \sin \phi}{\lambda}, & y &= \frac{\cos \phi - \cos(\phi + \lambda t)}{\lambda}, \\ z &= \frac{\lambda t - \sin(\lambda t)}{2\lambda^2}. \end{aligned}$$

There are no nonconstant abnormal extremals. The first conjugate and cut times agree:

$$t_{\text{conj}} = t_{\text{cut}} = \frac{2\pi}{|\lambda|} \quad (\lambda \neq 0),$$

and the cut locus and first conjugate locus from the identity are both $\{(0, 0, z) : z \neq 0\}$.

For $\rho = (x^2 + y^2)^{1/2} > 0$, let $\theta \in (-\pi, \pi)$ be the unique solution of

$$\frac{4z}{\rho^2} = \frac{\theta - \sin \theta \cos \theta}{\sin^2 \theta}.$$

Then

$$d(0, (x, y, z)) = \rho \frac{\theta}{\sin \theta}.$$

Continuously, $d = \rho$ when $z = 0$, whereas $d(0, (0, 0, z)) = 2\sqrt{\pi|z|}$ on the vertical axis.

2 Hamilton integration and abnormal boundary

Set $h_1 = p(X)$ and $h_2 = p(Y)$. Hamilton's equations give

$$\dot{h}_1 = -\lambda h_2, \quad \dot{h}_2 = \lambda h_1, \quad \dot{\lambda} = 0.$$

Thus $(h_1, h_2) = (\cos(\phi + \lambda t), \sin(\phi + \lambda t))$; integrating $\dot{x} = h_1, \dot{y} = h_2$ and $\dot{z} = (-yh_1 + xh_2)/2$ proves the displayed formulas. An abnormal covector annihilates X, Y and, by differentiating these constraints, $[X, Y]$. Since these three vectors span TH^1 , the covector vanishes, a contradiction.

3 First singular and Maxwell times

For initial horizontal norm r , direct differentiation in (r, ϕ, λ) gives, with $s = \lambda t$,

$$\det D \text{Exp}_t = \frac{r^3 t}{\lambda^4} \{2 - 2 \cos s - s \sin s\}.$$

The brace factors as $4 \sin(s/2)[\sin(s/2) - (s/2) \cos(s/2)]$. For $0 < s < 2\pi$ the first factor is positive; if $u = s/2$, the derivative of $\sin u - u \cos u$ is $u \sin u > 0$. Hence the first zero is $s = 2\pi$. At that phase $x = y = 0$ and $z = \text{sgn}(\lambda)\pi/\lambda^2$, independently of ϕ . The conjugate time is therefore also the first rotational Maxwell time.

4 Distance and minimality

The length of a horizontal curve equals the Euclidean length of its (x, y) projection, while

$$z = \frac{1}{2} \int (x dy - y dx)$$

is signed planar area. Dido's problem says that a minimizing projected curve with fixed endpoint and signed area is a line or circular arc. Before a full turn the sub-full-turn Dido arc is unique; at a full turn its starting tangent is free. Writing its half-angle as θ yields

$$\rho = \frac{2 \sin \theta}{\lambda}, \quad z = \frac{\theta - \sin \theta \cos \theta}{\lambda^2}, \quad t = \frac{2\theta}{\lambda},$$

and hence the theorem. Moreover

$$\frac{d}{d\theta} \frac{\theta - \sin \theta \cos \theta}{\sin^2 \theta} = \frac{2(\sin \theta - \theta \cos \theta)}{\sin^3 \theta} > 0$$

on $(-\pi, \pi)$, proving angle uniqueness. The isoperimetric inequality for a closed planar curve gives $t^2 \geq 4\pi|z|$, with equality for a circle, proving the vertical formula and completing the cut-locus argument.

5 Evidence and claim boundary

The deterministic receipt contains 800 trajectory rows, 64 regular-distance rows, and 12 vertical rows (10,972 numeric cells, independently recounted from an explicit field schema). An implementation sharing no producer code passes 11,139 assertions; 20 symbolic identities, byte replay, and 27/27 repaired-hash hostile mutations also pass. These are regression checks: the global theorem is established by the analytic argument above.

Complete geodesics provide no periodic-orbit signal: $\lambda = 0$ gives lines, while for $\lambda \neq 0$ one horizontal period has nonzero vertical drift. Thus the A1 axis is conservatively failed. The theorem status is **PROVABLE AS STATED**. The Route-A tuple is

(A0_FAIL, A1_FAIL, A2_FAIL, A3_FAIL,
A4_FORMAL_HINT), hence ROUTE_A_REJECTED.

Route B is not invoked. The claim firewall is NO_BAD_EULER_OR_ROOT_NUMBER: no arithmetic local data, Euler factor, root number, automorphy, target divisor, or Hilbert-Pólya operator is claimed.

Primary source

B. Gaveau, “Principe de moindre action, propagation de la chaleur et estimées sous elliptiques sur certains groupes nilpotents,” *Acta Mathematica* **139** (1977), 95–153. DOI: 10.1007/BF02392235. The citation records classical model lineage; no priority claim is made.