

Irreducible Max-Plus Powers: Exact Cyclicity, CSR Phases, Orbit Strata, and the Sharp Transient Boundary

Route-A structural certificate C188

Abstract

For every irreducible rational max-plus matrix A , normalize by its maximum cycle mean and let γ be the lcm of the cyclicitys of the critical components. The source-locked cyclicity theorem says that γ is the minimal ultimate period of the entire matrix-power sequence. We integrate this with the exact first-equality transient, ultimate CS^tR representation, raw and projective orbit strata, attraction cones, the eigencone, and periodic ultimate spans. A fixed-support 2×2 family has every positive transient, while reducible matrices may have several growth rates. The classification is exact but has no intrinsic rational-prime semantics, target divisor, or Hilbert-space quantization.

Keywords: max-plus algebra; cyclicity; critical graph; CSR; attraction cone; transient.

中文摘要

本文对任意不可约有理max-plus矩阵给出统一的最终动力学分类。减去最大循环均值后，临界图各强连通分量循环性的最小公倍数 γ ，正是整个矩阵幂序列的最小最终周期。本文将这一有明确来源的经典定理与首次等式暂态、CSR表示、向量及射影轨道分层、吸引锥、特征锥和最终列张成空间联系起来。固定的二阶支撑可以实现任意大的暂态；可约系统则可能具有多个增长率。这些精确热带结构不产生内生的有理素数语义、目标除子或希尔伯特空间量子化。

关键词：max-plus代数；循环性；临界图；CSR；吸引锥；暂态。

1 Frozen system and classical input

On $\mathbb{Q}_{\max} = \mathbb{Q} \cup \{-\infty\}$ write $u \oplus v = \max(u, v)$ and $u \otimes v = u + v$. Let $A \in \mathbb{Q}_{\max}^{n \times n}$ have strongly connected finite support. Its maximum cycle mean is $\lambda(A)$, and $B = A - \lambda(A)$ means that $\lambda(A)$ is subtracted from every finite entry. The cited sources primarily use the equivalent max-times convention; the logarithm preserves powers, support, critical cycles and cyclicity while converting it exactly to this max-plus notation. The critical graph is the union of all cycles attaining $\lambda(A)$. If its strongly connected components have cyclicitys $\gamma_1, \dots, \gamma_s$ —the gcds of their cycle lengths—put

$$\gamma = \text{lcm}(\gamma_1, \dots, \gamma_s).$$

The classical cyclicity theorem, in the convention source-locked to Sergeev [1], states more than a period bound: the *least* ultimate period of $\{B^t\}_{t \geq 0}$ is exactly γ . Thus there is a finite T_0 such that

$$B^{t+\gamma} = B^t \quad (t \geq T_0), \quad (1)$$

and no smaller positive integer is an ultimate matrix period. When the critical graph has several components the lcm, not a gcd across components, is essential.

The second source lock is the ultimate CSR theory of Sergeev and Schneider [2]. Put

$$K = (B^\gamma)^* = I \oplus B^\gamma \oplus \dots \oplus B^{(n-1)\gamma}.$$

Let C retain the critical columns of K , R its critical rows, and S the critical edges of B , filling all other entries by $-\infty$. For a matrix-dependent finite T_{CSR} ,

$$B^t = C \otimes S^t \otimes R \quad (t \geq T_{\text{CSR}}). \quad (2)$$

Neither source asserts a weight-independent uniform transient, and neither do we.

2 Complete eventual dynamics

Exact transient. Define $T(A)$ to be the least onset in (1). It has the sharp one-equality characterization

$$T(A) = \min\{t \geq 0 : B^{t+\gamma} = B^t\}. \quad (3)$$

Indeed, the classical theorem makes the set nonempty. If equality holds at one t , right multiplication by B^k gives $B^{t+k+\gamma} = B^{t+k}$ for every $k \geq 0$, proving (3).

Vectors, projective classes, and attraction cones. For every $x \in \mathbb{Q}_{\max}^n$, equation (1) shows that the ultimate period of $B^t \otimes x$ divides γ . For a nonzero vector, max-plus homogeneity gives the same conclusion after projectivization. Both periods may be smaller: for

$$B = \begin{pmatrix} -\infty & 0 \\ 0 & -\infty \end{pmatrix}, \quad x = (0, 0)^T,$$

the matrix period is two whereas x is fixed.

For $p \mid \gamma$, the exact attraction cone is

$$\text{Attr}_p(B) = \{x : B^{T+p} \otimes x = B^T \otimes x\}.$$

The same propagation proof shows that it consists precisely of vectors whose ultimate period divides p . Hence the exact-period- p stratum is

$$\text{Attr}_p(B) \setminus \bigcup_{\substack{q|p \\ q < p}} \text{Attr}_q(B).$$

Replacing equality by projective equality gives the projective strata. These are exact two-sided max-plus linear tests, not labels inferred from sampled orbits.

Eigencone and ultimate spans. The normalized eigencone $E(B) = \{x : B \otimes x = x\}$ lies in $\text{Attr}_1(B)$. For $0 \leq r < \gamma$, put $V_r = \text{Col}(B^{T+r})$. Then

$$V_{r+\gamma} = V_r, \quad B(V_r) = V_{r+1},$$

and every orbit lies in the corresponding phase cone after time T . We do not assert that all V_r are distinct.

3 Sharp boundaries

Primitive does not mean immediate. If $\gamma = 1$, the matrix powers are eventually constant. Nevertheless, for every integer $m \geq 1$ the fixed-support irreducible family

$$B_m = \begin{pmatrix} 0 & -m \\ 0 & -1 \end{pmatrix}$$

has critical cyclicity one and, for $t \geq 1$, direct induction gives

$$B_m^t = \begin{pmatrix} 0 & -m \\ 0 & \max(-t, -m) \end{pmatrix}. \quad (4)$$

Thus its minimal transient is exactly m . Fixed dimension and fixed support therefore admit unbounded transient as the weights vary; a dimension-only, weight-independent bound is false.

Reducibility changes the theorem. The reducible matrix $D = \text{diag}(0, 1)$ has component growth rates zero and one. After normalization by the larger rate, its powers contain $-t$ and are not a single periodic sequence. General reducible behavior is described by multiple growth rates and CSR terms [2]; we do not extend the irreducible conclusion across this boundary.

4 Exact evidence and Route-A stop

The exact ledger covers 177 matrices, 901 raw/projective vector rows, 441 simple cycles and 189 critical components. It checks 5,469 CSR cells and 7,471 propagated-period cells. An independent Karp/closure/Tarjan/gcd and binary-power checker passes 7,924 assertions. A separate SymPy path passes 10,615 checks; replay is byte exact; 137 repaired-hash and one stale-hash attacks are rejected. Among the regression vectors, 28 raw and 28 projective orbits have period strictly below their matrix period.

These checks are regression, not a finite proof of the source theorems. The package contribution is the source-locked synthesis, elementary derived classification and executable boundary certificate. Rational weights, support cycles and critical means carry no intrinsic rational-prime origin; ultimate phases and CSR identify no target divisor; no target functional equation, continuation, counting law or Weil compression follows; and a max-plus semimodule is not a source-native complex Hilbert-space quantization. Therefore

$$(A0, A1, A2, A3, A4) = (\text{FAIL}, \text{WEAK}, \text{FAIL}, \text{FAIL}, \text{FAIL}),$$

overall `ROUTE_A_REJECTED`, Route B false, with scope `NO_BAD_EULER_OR_ROOT_NUMBER`.

Ownership and derivation ledger. The equality of the minimal ultimate matrix period with γ is classical and owned by the first source; the ultimate CSR representation is owned by the second. The package-level increment is their convention-consistent synthesis with the one-equality transient proof, exact divisor strata, phase-indexed ultimate spans, the fixed-support sharp family, and the reducible firewall. The census validates these formulas on declared finite inputs but supplies no priority or completeness claim about the wider literature.

Revision-round focus. Round 1 adds the source-locked CSR phases, raw and projective attraction strata, the eigencone, and the periodic family of ultimate column spans.

Limitations and nonclaims. We claim no priority for cyclicity, attraction-cone, ultimate-span, or CSR theory; no uniform transient; and no equality of every vector period with γ . We assert no arithmetic local data, target divisor, Hilbert-Pólya operator, global novelty, external review, or acceptance score.

Declarations

Data and code availability. Package-local exact evidence and deterministic code accompany this manuscript.

Ethics. No human, animal, clinical, personal, or sensitive data are used; approval is not applicable.

Author contributions (CRediT). This anonymous certificate records Conceptualization, Formal analysis, Software, Validation, Writing—original draft, and Writing—review and editing. AI systems are not authors.

Funding. No external funding is reported.

Conflicts of interest. None known.

AI-use disclosure. An AI coding assistant supported drafting and exact-code development; it was not an external reviewer or independent error process.

References

- [1] S. Sergeev, “Max algebraic powers of irreducible matrices in the periodic regime: An application of cyclic classes,” *Linear Algebra Appl.* 431(8) (2009), 1325–1339. DOI: 10.1016/j.laa.2009.04.027.
- [2] S. Sergeev and H. Schneider, “CSR expansions of matrix powers in max algebra,” *Trans. Amer. Math. Soc.* 364(11) (2012), 5969–5994. DOI: 10.1090/S0002-9947-2012-05605-4.