

Rectangular Promotion: Cyclic Sieving, the Complete Cycle Ledger, and the Finite Koopman Spectrum

Route-A structural certificate C187

Abstract

For every rectangular standard Young-tableau set $X_{a,b} = \text{SYT}(b^a)$, we turn Rhoades's attributed cyclic-sieving theorem into a complete finite dynamical classification. Root-of-unity evaluations of the unshifted q -hook polynomial give every fixed iterate; Möbius inversion gives all least periods and cycles, hence the Artin–Mazur zeta, finite Koopman determinant, traces, and every spectral multiplicity. Evacuation reverses promotion. The theorem says $j^{ab} = 1$, not that the order always equals ab . The resulting finite unitary is natural but carries no intrinsic rational-prime semantics or target divisor.

Keywords: promotion; standard Young tableaux; cyclic sieving; q -hook formula; Koopman spectrum; evacuation.

中文摘要

本文对任意矩形标准杨表上的升运算给出完整有限动力学分类。以 Rhoades 的未平移 q 钩长循环筛分定理为归属明确的输入，单位根取值给出每个迭代的固定点数；Möbius 反演继而给出所有最小周期、循环、有限动力学 ζ 函数、Koopman 行列式、迹与谱重数。疏散算子将升运算共轭到其逆。统一结论是 $j^{ab} = 1$ ，而不是所有矩形的实际阶都等于 ab 。该有限酉算子虽属源系统自身，却没有内生的有理素数语义或目标除子匹配。

关键词：升运算；标准杨表；循环筛分； q 钩长公式；Koopman 谱；疏散。

1 Frozen convention and source theorem

Fix positive integers a, b , put $N = ab$, and let $X_{a,b} = \text{SYT}(b^a)$. We use the promotion convention j in Rhoades [1]: remove N , slide the hole northwest, increment the retained entries, and insert 1. Removing 1 and sliding southeast is the inverse clock and is not silently substituted.

For $[r]_q = (q^r - 1)/(q - 1)$ define the *unshifted* polynomial

$$F_{a,b}(q) = \frac{[N]_q!}{\prod_{c \in b^a} [h(c)]_q}.$$

Rhoades's rectangular standard-tableau CSP states, with $\omega_N = e^{2\pi i/N}$,

$$\# \text{Fix}(j^d) = F_{a,b}(\omega_N^d) \quad (d \in \mathbb{Z}). \quad (1)$$

The same source records the classical consequence $j^N = 1$, with order background attributed to Haiman [2]. It explicitly warns that the actual order need not be N . Thus the cyclic group of order N may act nonfaithfully.

The polynomial convention matters. The neighboring column-strict result in the source uses a shifted Schur specialization; importing that shift into (1) would change fixed counts. We use exactly the standard-tableau formula displayed above.

2 From fixed points to every cycle

Let P_ℓ be the number of tableaux of least promotion period ℓ , and let $C_\ell = P_\ell/\ell$. Since $j^N = 1$, only $\ell \mid N$ occur. A tableau is fixed by j^d precisely when its least period divides d , so divisor Möbius inversion gives

$$P_\ell = \sum_{d \mid \ell} \mu(\ell/d) F_{a,b}(\omega_N^d), \quad C_\ell = \frac{P_\ell}{\ell}. \quad (2)$$

This is an all-rectangle consequence of the source theorem, not an inference from a finite table. It also yields the actual promotion order as $\text{lcm}\{\ell : C_\ell > 0\}$, a divisor of N .

Shape	$ X_{a,b} $	nonzero cycle counts	actual order
$1 \times b$ or $a \times 1$	1	$C_1 = 1$	1
2×2	2	$C_2 = 1$	$2 < 4$
2×3	5	$C_2 = C_3 = 1$	6
3×3	42	$C_3 = 2, C_9 = 4$	9
3×4	462	$C_3 = 2, C_4 = 3, C_6 = 4, C_{12} = 35$	12

For the sharp boundary, $F_{2,2}(q) = 1 + q^2$. Its evaluations give no fixed tableau at one promotion step and two at the second, hence one two-cycle. A uniform claim of exact order N is therefore false even away from the one-dimensional singleton families.

3 Zeta, Koopman determinant, and reversal

The finite Artin–Mazur series and the cycle blocks give

$$\zeta_j(z) = \exp\left(\sum_{d \geq 1} \frac{\#\text{Fix}(j^d)}{d} z^d\right) = \prod_{\ell|N} (1 - z^\ell)^{-C_\ell}, \quad (3)$$

$$\det(I - zU_{a,b}) = \prod_{\ell|N} (1 - z^\ell)^{C_\ell} = \zeta_j(z)^{-1}, \quad (4)$$

where $(U_{a,b}f)(T) = f(jT)$ on counting $\ell^2(X_{a,b})$. Each cycle is a cyclic permutation block. Consequently

$$\text{Tr}(U_{a,b}^d) = \#\text{Fix}(j^d), \quad \text{mult}(\omega_N^k) = \sum_{\substack{\ell|N \\ N|k\ell}} C_\ell.$$

Summing these multiplicities over $k \bmod N$ gives $\sum_\ell \ell C_\ell = |X_{a,b}|$, so no spectral sector is missing.

Schützenberger evacuation e is an involution and satisfies $eje = j^{-1}$ on the rectangle [1]. Its permutation matrix E is a unitary reversor, $EUE = U^{-1}$. Since all matrices are real, E followed by coefficientwise conjugation is an antiunitary involution with the same reversal law. This is a natural source operator statement; it is not a post-hoc diagonal encoding.

4 Exact evidence and Route-A stop

The formula ledger covers all 36 rectangles with $1 \leq a, b \leq 6$: 441 iterate rows, 162 period rows, and 441 root-multiplicity rows. Its largest population is the 6×6 value 1,671,643,033,734,960, handled symbolically. An independent checker constructs 37,401 tableaux across 26 declared small rectangles and passes 230,034 assertions, including promotion/demotion, evacuation and all direct cycles. A separate SymPy path passes 3,065 checks; replay is byte exact; 107 repaired-hash and one stale-hash attacks are rejected.

These finite checks are regression evidence, not a new proof of the Kazhdan–Lusztig CSP. Source ownership remains with Rhoades and Haiman. The package contribution is the source-locked Route-A synthesis and executable boundary certificate.

Promotion has natural primitive cycles and a source-native finite unitary, but rectangle dimensions, hook lengths and cyclotomic orders do not intrinsically encode rational primes or prime powers. The finite root-of-unity determinant has no established target divisor, functional equation, continuation, counting law, or Weil compression. Therefore

$$(A0, A1, A2, A3, A4) = (\text{FAIL}, \text{WEAK}, \text{FAIL}, \text{FAIL}, \text{NATURAL QUANTIZATION}),$$

overall `ROUTE_A_REJECTED`, with Route B false and scope `NO_BAD_EULER_OR_ROOT_NUMBER`.

Revision-round focus. Round 2 adds evacuation reversal, separates the attributed infinite theorem from finite regression, and locks the semantic mutation audit and strict Route-A stopping boundary.

Limitations and nonclaims. We do not claim priority for cyclic sieving, the q-hook formula, the order bound, or evacuation. We do not claim exact order N uniformly or use finite enumeration as an infinite proof. No rational-prime semantics, target divisor, Hilbert–Polya operator, external review, or acceptance score is asserted.

Declarations

Data and code availability. Package-local exact evidence and deterministic code accompany this manuscript.

Ethics. No human, animal, clinical, personal, or sensitive data are used; approval is not applicable.

Author contributions (CRediT). This anonymous certificate records Conceptualization, Formal analysis, Software, Validation, Writing—original draft, and Writing—review and editing. AI systems are not authors.

Funding. No external funding is reported.

Conflicts of interest. None known.

AI-use disclosure. An AI coding assistant supported drafting and exact-code development; it was not an external reviewer or independent error process.

References

- [1] B. Rhoades, “Cyclic sieving, promotion, and representation theory,” *J. Combin. Theory Ser. A* 117(1) (2010), 38–76. DOI: 10.1016/j.jcta.2009.03.017.
- [2] M. D. Haiman, “Dual equivalence with applications, including a conjecture of Proctor,” *Discrete Math.* 99 (1992), 79–113. DOI: 10.1016/0012-365X(92)90368-P.