

A Complete Cycle Atlas for Projective Möbius Dynamics over Finite Fields

Route-A source-local certificate HCS-C260

31 August 2026

Abstract

For every prime power $q = p^r$, we classify the permutation induced by each $g \in \text{PGL}_2(\mathbb{F}_q)$ on $\mathbb{P}^1(\mathbb{F}_q)$. Four cases give the complete cycle multiset, all iterated fixed counts, primitive cycles, the finite Artin–Mazur zeta, and the finite Koopman spectrum. Exhaustive direct permutation checks cover 18 prime and extension fields. All statements are source-local and use no target spectral data.

1 The four cycle types

Represent g by $A \in \text{GL}_2(\mathbb{F}_q)$, modulo nonzero scalar. In the semisimple cases let d be the order of the eigenvalue ratio.

Theorem 1 (complete projective cycle atlas). *Exactly one of the following rows applies.*

<i>type</i>	<i>condition</i>	<i>cycles on $\mathbb{P}^1(\mathbb{F}_q)$</i>	<i>$F_n = \# \text{Fix}(g^n)$</i>
<i>identity</i>	$g = 1$	1^{q+1}	$q + 1$
<i>unipotent</i>	<i>nontrivial, order p</i>	$1^1 p^{q/p}$	$1 + q \mathbf{1}_{p n}$
<i>split</i>	$d > 1, d \mid q - 1$	$1^2 d^{(q-1)/d}$	$2 + (q - 1) \mathbf{1}_{d n}$
<i>nonsplit</i>	$d > 1, d \mid q + 1$	$d^{(q+1)/d}$	$(q + 1) \mathbf{1}_{d n}$

For every $m \geq 1$, the numbers of points and cycles of least period m are

$$P_m = \sum_{e|m} \mu(m/e) F_e, \quad C_m = P_m/m. \quad (1)$$

Proof. The scalar case is immediate. A nonscalar repeated-root matrix has one rational eigenline and is projectively conjugate to $x \mapsto x + 1$. It fixes infinity and partitions the affine line into q/p cycles of length p . If the characteristic polynomial splits with distinct roots, conjugate the two rational eigenlines to 0 and infinity. The map is $x \mapsto \rho x$; ρ has order $d \mid q - 1$ and acts freely on $\mathbb{F}_q^*$.

In the irreducible case the two eigenlines lie over $\mathbb{F}_{q^2}$ and Frobenius interchanges them. Thus $\rho^q = \rho^{-1}$ and $d \mid q + 1$. A proper nonidentity power cannot acquire a rational eigenline: it retains the same two nonrational eigenlines. Hence the cyclic action is semiregular on all $q + 1$ rational projective points. The fixed formulas sum cycle lengths dividing n , and Möbius inversion gives (1). $\square$

2 Zeta and finite Koopman closure

For a finite permutation with c_L cycles of length L ,

$$\zeta_g(t) = \exp \left(\sum_{n \geq 1} \frac{F_n}{n} t^n \right) = \prod_L (1 - t^L)^{-c_L}. \quad (2)$$

Consequently the four rows give, respectively,

$$(1 - t)^{-(q+1)}, \quad (1 - t)^{-1} (1 - t^p)^{-q/p}, \quad (1 - t)^{-2} (1 - t^d)^{-(q-1)/d}, \quad (1 - t^d)^{-(q+1)/d}. \quad (3)$$

On counting-measure $\ell^2(\mathbb{P}^1(\mathbb{F}_q))$, the Koopman operator is a permutation unitary. Every L -cycle contributes each L th root of unity once, and

$$\det(I - tU_g) = \prod_L (1 - t^L)^{c_L} = \zeta_g(t)^{-1}. \quad (4)$$

3 Exact regression and boundary

Two implementations cover $q = 2, 3, 4, 5, 7, 8, 9, 11, 13, 16, 17, 19, 23, 25, 27, 29, 31, 32$. The independent checker rebuilds every projective permutation and derives its type only from fixed-point geometry and cycle lcm. In total it checks 155,346 transformations, 4,367,094 state images, and closes 6,159,318 assertions. Symbolic algebra closes 193 identities; byte replay is exact; semantic mutation rejects 40/40 repaired-hash changes. Finite enumeration is a regression oracle, not the proof for arbitrary q .

Finite-field arithmetic provides only `AO_WEAK_ARITHMETIC_RELATION`; Theorem 1 earns `A1_PASS_ANALYTIC`, while (4) gives `A4_NATURAL_QUANTIZATION`. There is no rational-prime orbit dictionary, target divisor, or target global analytic structure. The strict tuple is

`(AO_WEAK_ARITHMETIC_RELATION, A1_PASS_ANALYTIC, A2_FAIL, A3_FAIL, A4_NATURAL_QUANTIZATION)`.

The verdict is `ROUTE_A_EXPLORATORY`; Route B is disabled. Under `NO_BAD_EULER_OR_ROOT_NUMBER`, (2) is only a finite source zeta, and (4) is not a Hilbert–Polya identification.

References

- [1] E. Sakzad, M.-R. Sadeghi, and D. Panario, *Adv. Math. Commun.* **6** (2012), 347–361, [doi:10.3934/amc.2012.6.347](https://doi.org/10.3934/amc.2012.6.347).
- [2] C. Forsyth, J. Gurev, and S. Shrima, *Proc. Amer. Math. Soc.* **144** (2016), 4583–4590, [doi:10.1090/proc/13126](https://doi.org/10.1090/proc/13126).
- [3] G. E. Wall, *Bull. Aust. Math. Soc.* **22** (1980), 339–364, [doi:10.1017/S0004972700006675](https://doi.org/10.1017/S0004972700006675).