

Exact Spectra for Hyperplane Chamber Walks: A Route-A Stress Test

HCS-C192 theorem and evidence package

27 August 2026

Abstract

For every finite real hyperplane arrangement and every probability measure on its faces, the Brown–Diaconis theorem diagonalizes the induced chamber walk. We record the resulting characteristic polynomial, finite determinant, power traces, unique-stationarity criterion, exact sampler, and mixing bound. Exact rational coordinate and braid fixtures test the formulas but do not prove the all-family theorem. The complete operator structure still fails Route A: without target arithmetic semantics, its determinant is only a formal hint.

1 Face dynamics and the flat spectrum

Let $\mathcal{A}$ be a finite real hyperplane arrangement in V , with faces $\mathcal{F}$ and chambers $\mathcal{C}$. In sign coordinates the face product is $(FG)_H = F_H$ when $F_H \neq 0$ and $(FG)_H = G_H$ otherwise. A face law w defines

$$K(C, C') = \sum_{F: FC=C'} w(F).$$

Write $L(\mathcal{A})$ for the intersection poset and μ for its Möbius function. Brown and Diaconis [1, Theorem 1] prove that K is diagonalizable, with flat-indexed data

$$\lambda_W = \sum_{F \subseteq W} w(F), \quad m_W = |\mu(W, V)|.$$

Consequently, even when several flats yield the same numerical eigenvalue,

$$\begin{aligned} \chi_K(x) &= \prod_{W \in L(\mathcal{A})} (x - \lambda_W)^{m_W}, \\ \det(I - zK) &= \prod_{W \in L(\mathcal{A})} (1 - z\lambda_W)^{m_W}, \\ \mathrm{tr}(K^\ell) &= \sum_W m_W \lambda_W^\ell. \end{aligned}$$

These are finite-dimensional deductions from the cited theorem, not a new diagonalization claim. In particular, diagonalizable does not mean reversible or self-adjoint.

2 Stationarity and convergence

A face law is *separating* if each $H \in \mathcal{A}$ is avoided by some positive-weight face. Brown–Diaconis Theorem 2 proves that separation is equivalent to unique stationarity. Under separation, sample all positive-weight faces without replacement, choosing each next face proportionally to its remaining weight. Their product is a chamber with the stationary law π .

For every start C and $\ell \geq 0$, their coupling argument gives

$$\|K_C^\ell - \pi\|_{\mathrm{TV}} \leq - \sum_{W \neq V} \mu(W, V) \lambda_W^\ell \leq \sum_{H \in \mathcal{A}} \lambda_H^\ell.$$

The middle term is exactly the probability that the product of ℓ i.i.d. sampled faces is not yet a chamber [1, Section 4B].

Stopping terminology. Sampling with replacement and stopping when the accumulated face product first becomes a chamber gives the same stationary marginal: earlier repeated factors may be deleted to recover weighted sampling without replacement. This is an exact stationary stopping sampler and the stopping event drives the displayed coupling bound. We do not call it a strict strong stationary time. That term would additionally require independence of the stopped chamber and the stopping time, a statement not supplied by the cited theorem and false for general unequal-weight coupon orders.

Nonseparating boundary. Let $\mathcal{A}_0$ contain precisely the hyperplanes that contain every positive-weight face. Brown and Diaconis show that the $\mathcal{A}_0$ -chambers index closed components of the walk. Each component has one stationary law, and the set of all stationary laws is exactly the simplex with one vertex per $\mathcal{A}_0$ -chamber. Thus nonseparation has a complete component classification, not merely a failure of uniqueness. Their Section 6 carries Theorems 1 and 2 to oriented-matroid covector face semigroups; we import only that stated extension and no broader affine or realization claim.

3 Finite exact regression

Two independent rational implementations and a separate SymPy oracle check eight coordinate/braid fixtures. The census contains 316 faces, 94 chambers, 75 flats, 1,604 transition cells, 62 stationary probabilities, 24 mixing rows, and 48 trace rows. The independent checker passes 20,609 assertions; SymPy passes 3,398 exact checks. These finite examples are regression oracles only.

The producer generates coordinate sign cubes and ordered-set-partition covectors, then enumerates weighted face orders. The checker imports no producer code: it reconstructs face closure and the support lattice from the serialized signs, recomputes Möbius values recursively, and obtains the stationary law by subset dynamic programming. SymPy separately recomputes the matrix polynomials, traces, and geometric multiplicities. Byte replay is exact, and 74 repaired-payload semantic attacks plus one stale-hash attack are rejected.

Statement	Owner/evidence	Ceiling
flat spectrum and multiplicity	Brown–Diaconis Theorem 1	no self-adjointness
sampler, uniqueness, mixing	Theorem 2 and Section 4	no strict-SST independence
nonseparating simplex	Section 4B remarks	no irreducibility across components
oriented-matroid transfer	Section 6	only the stated covector framework
matrix identities	elementary deduction here	no target determinant
finite census	exact regression code	no proof by enumeration

Table 1: Claim ownership and release ceilings.

4 Route-A verdict

The exact result has no intrinsic rational-prime or target-zero index, recovers no target arithmetic data, supplies no target functional equation, and yields no target counting law. Its finite determinant is a natural operator object but no target divisor is identified. Hence

$$(A_0, A_1, A_2, A_3, A_4) = (\text{FAIL}, \text{FAIL}, \text{FAIL}, \text{FAIL}, \text{FORMAL_HINT}),$$

with overall rejection and no Route-B invocation. No local or Euler factors, root numbers, automorphy, Hilbert–Pólya operator, global novelty, or external review are claimed.

References

- [1] K. S. Brown and P. Diaconis, Random walk and hyperplane arrangements, *Ann. Probab.* 26 (1998), 1813–1854, doi:10.1214/aop/1022855884.