

JOINT ABSORPTION AND SPINE-FLIP LAWS FOR LOCAL TRIAD DYNAMICS ON TRIANGULAR BOOKS

ANONYMOUS

ABSTRACT. For the established $p = 1/3$ local triad dynamics, we specialize to the triangular book $B(3, r) = K_{1,1,r}$ and count active imbalanced-triad update epochs. The page-imbalance count is a reflected one-dimensional quotient. We solve its transform jointly marked by absorption time and common-spine flips in an explicit Chebyshev-rational form. The mean absorption time is the quadratic $k(r + 2 - k)/2$, with sharp extrema over the starting count. The mean together with the probability of odd spine parity exactly identifies the book size and initial imbalance count, subject to an explicit integer feasibility criterion, while either statistic alone has concrete collisions. A private-edge block supplies a uniform absorption certificate. All formulas include the $r = 1$, $r = 2$, $z = 0$, coincident-arrow, and clock-convention boundaries. An exact-arithmetic audit performs 191,278 rational and integer checks.

1. OWNED PROCESS, SPECIAL CARRIER, AND THEOREM

Let

$$B_r = B(3, r) = K_{1,1,r}, \quad r \geq 1,$$

be the graph obtained by joining r triangles along one common *edge*, called the spine. Thus each page has two private edges. Give every physical edge a sign in $\{\pm 1\}$, and call a page imbalanced when the product of its three edge signs is negative. At an update epoch, choose a currently imbalanced page uniformly, choose one of its three physical edges uniformly, and flip that sign. Stop when all pages are balanced.

Write $x_i \in \{0, 1\}$ for the imbalance bit of page i and $K = \sum_i x_i$. Let T be the number of these *active update epochs* until absorption, and let J be the number of spine flips before absorption. Expectations conditional on any sign state with $K = k$ are denoted by $\mathbb{E}_k$; strong lumpability below makes this notation unambiguous.

This stochastic rule is the $p = 1/3$ specialization of local triad dynamics of Antal–Krapivsky–Redner [1, 2]. The 2005 all-triad clock samples balanced pages too and hence inserts no-op holds; the process above is its chain embedded at active epochs. The 2006 formulation directly selects an imbalanced triad. Istrate owns the same probabilistic kernel on general graphs and its triadic-dual/XOR representation [3]; later hypergraph particle-system and drift methods are also background [4]. Signed-book work owns the carrier and its static switching-class count indexed by negative pages [5]. Table 1 makes the resulting claim boundary explicit.

TABLE 1. Primary-source subtraction. The middle column receives zero contribution credit; the right column is the special-carrier residual.

Source	Owned input	Residual used here
Antal–Krapivsky–Redner [1, 2]	local triad dynamics and the $p = 1/3$ equiprobable-edge update	exact joint law after specialization to $B(3, r)$ and the active clock
Istrate and Istrate–Bonchis–Marin [3, 4]	same general-graph kernel, triadic dual/XOR encoding, hypergraph particle systems, drift, and convergence-time program	reflected count recurrence, closed transform, and coarse inverse on the one-spine carrier only
Sehrawat–Bhattacharjya [5]	signed-book carrier and static switching classes indexed by page count	no carrier, switching, coloring, or static-class claim
Generic finite-chain tools	Bellman/resolvent rationality, Chebyshev facts, tail sums, and quadratic concavity	only the boundary-complete formula conjunction stated below

For $0 \leq k \leq r$, define the joint transform

$$(1) \quad F_k(z, u) = \mathbb{E}_k[z^T u^J], \quad F_0(z, u) = 1.$$

Let U_j denote the Chebyshev polynomial of the second kind, with $U_{-1} = 0$, $U_0 = 1$, and $U_{j+1}(x) = 2xU_j(x) - U_{j-1}(x)$.

Theorem 1 (joint law, clock, inverse, and absorption). *For the process on B_r started with $K = k$, $1 \leq k \leq r$, the following hold.*

(i) *The full sign chain is strongly lumpable by K , with count transitions*

$$(2) \quad k \longrightarrow k-1 \quad \text{with probability } \frac{2}{3}, \quad k \longrightarrow r-k \quad \text{with probability } \frac{1}{3}.$$

If the targets coincide, the two masses add.

(ii) *For $r \geq 2$ and $z \neq 0$, put*

$$(3) \quad \xi = \frac{9 + z^2(4 - u^2)}{12z}.$$

Then, as identities of reduced rational functions,

$$(4) \quad F_k(z, u) = U_{k-1}(\xi)F_1(z, u) - U_{k-2}(\xi),$$

$$(5) \quad F_1(z, u) = \frac{3U_{r-2}(\xi) - 2zU_{r-3}(\xi) + zu}{3U_{r-1}(\xi) - 2zU_{r-2}(\xi)}.$$

For probability-transform evaluation, these identities hold on $|z| < 1$, $|u| \leq 1$, with removable points taken from the Bellman system. The boundary values are

$$(6) \quad r = 1: \quad F_1 = \frac{z(2+u)}{3}; \quad r = 2: \quad F_1 = \frac{2z}{3-zu}; \quad z = 0: \quad F_k = \mathbf{1}_{\{k=0\}}.$$

(iii) *The mean is*

$$(7) \quad m_k := \mathbb{E}_k T = \frac{k(r+2-k)}{2}.$$

For $r > 1$, its unique minimum over nonabsorbing counts is at $k = 1$, with value $(r+1)/2$. Its maxima are

$$(8) \quad \begin{cases} k = (r+2)/2, & m_k = (r+2)^2/8, & r \text{ even,} \\ k = (r+1)/2, (r+3)/2, & m_k = ((r+2)^2 - 1)/8, & r \text{ odd.} \end{cases}$$

For $r = 1$, the sole nonabsorbing state is both extremizers and has mean one.

(iv) *If $q_k = \mathbb{P}_k(J \text{ is odd})$, then*

$$(9) \quad \mathbb{E}_k[(-1)^J] = \frac{r+2-2k}{r+2}, \quad q_k = \frac{k}{r+2}.$$

Given exact candidate observations (m, q) for a nonabsorbing start, first require $0 < q < 1$ and set

$$(10) \quad R = \sqrt{\frac{2m}{q(1-q)}}.$$

The pair is feasible if and only if R is an integer with $R \geq 3$ and $k = qR$ is an integer satisfying $1 \leq k \leq R-2$. In that case

$$(11) \quad r = R-2, \quad k = qR$$

are uniquely recovered. Both observations are necessary in general.

(v) *For every integer $n \geq 0$,*

$$(12) \quad \mathbb{P}_k(T > nr) \leq \left[1 - \left(\frac{2}{3}\right)^r\right]^n.$$

In particular, absorption is almost sure and T has an exponential tail.

The theorem recovers only the carrier size and initial imbalance *count*. It neither reconstructs a full edge-sign configuration nor supplies stability under noisy observations.

2. STRONG LUMPING AND AN ABSORPTION CERTIFICATE

Proof of Theorem 1(i) and (v). Flipping either private edge of the selected active page clears that page and changes no other imbalance bit. Both private choices therefore send every state of count k to the count $k - 1$. Flipping the common spine toggles all r imbalance bits and sends count k to $r - k$. The probabilities are the same for every full sign state within the count fibre, which proves strong lumpability and (2). If $k - 1 = r - k$, the private and spine choices are different physical updates with the same quotient target, so their probabilities combine rather than producing two arrows.

Pre-generate the private/spine type of the next r edge choices. At each active epoch the type is private with probability $2/3$, independently of the state. If all r types are private, each performed update lowers K by one, so absorption occurs within the block, possibly before all pre-generated choices are used. This event has probability $(2/3)^r$. Conditional on survival to any block boundary, the same bound applies. Iteration with the Markov property proves (12), and hence almost-sure absorption. $\square$

On the friendship or Dutch-windmill graph, the triangles share only a vertex, not an edge. Every physical edge is then private, so $K \mapsto K - 1$ and $T = K$ deterministically. That carrier does not satisfy (2).

3. THE SPINE-MARKED CHEBYSHEV TRANSFORM

First-step conditioning in the quotient gives, for $1 \leq k \leq r$,

$$(13) \quad F_k = z \left(\frac{2}{3} F_{k-1} + \frac{u}{3} F_{r-k} \right).$$

The factor u marks precisely a common-spine flip; it does not mark all sign flips or the terminal number of negative edges.

Proof of Theorem 1(ii). Work first in the rational-function field with $z \neq 0$. For $1 \leq k < r$, equations (13) at k and $r - k$ give

$$(14) \quad u F_{r-k} = \frac{3}{z} F_k - 2 F_{k-1}, \quad F_{r-k-1} = \frac{3}{2z} F_{r-k} - \frac{u}{2} F_k.$$

Insert the second identity into (13) at $k + 1$, then use the first. No division by u is needed, and simplification gives

$$(15) \quad F_{k+1} = \frac{9 + z^2(4 - u^2)}{6z} F_k - F_{k-1} = 2\xi F_k - F_{k-1}.$$

Since $F_0 = 1$, the Chebyshev solution of (15) is (4). At the terminal count, (13) becomes

$$(16) \quad F_r = \frac{2z}{3} F_{r-1} + \frac{zu}{3}.$$

Substituting (4) into (16) and collecting the F_1 terms gives (5).

For $r = 1$, every edge choice absorbs in one epoch; two choices are private and one is the marked spine, which gives the first formula in (6). When $r = 2$, a spine flip from $k = 1$ is a self-loop, so (13) directly gives $F_1 = 2z/(3 - zu)$. Equivalently, the unreduced Chebyshev ratio is

$$\frac{3 + zu}{6\xi - 2z}, \quad 6\xi - 2z = \frac{(3 - zu)(3 + zu)}{2z};$$

the removable factor $3 + zu$ must be cancelled before pointwise evaluation. Finally, ξ is undefined at $z = 0$, whereas (13) gives $F_0 = 1$ and $F_k = 0$ for $k > 0$. Thus (4)–(5) are read as reduced rational identities with these Bellman continuations. $\square$

Uniform choices matter. Nonuniform page weights generally destroy count lumpability, while nonuniform physical-edge weights change the coefficients in both (2) and (13).

4. QUADRATIC CLOCK AND EXACT COARSE INVERSE

Proof of Theorem 1(iii). The exponential tail justifies first-step conditioning of the mean:

$$(17) \quad m_0 = 0, \quad m_k = 1 + \frac{2}{3} m_{k-1} + \frac{1}{3} m_{r-k}.$$

Use (17) at k , $r - k$, and $k + 1$. Eliminating the two reflected terms yields

$$m_{k+1} - 2m_k + m_{k-1} = -1 \quad (1 \leq k < r).$$

The terminal equation is $m_r = 1 + (2/3)m_{r-1}$. Together with $m_0 = 0$, these conditions have the unique quadratic solution (7).

For $r > 1$, strict concavity makes the minimum occur at an endpoint; since $m_1 = (r+1)/2 < m_r = r$, it is uniquely at $k = 1$. The vertex is $(r+2)/2$, so the nearest admissible integer or integers give exactly (8). The one-state case $r = 1$ is immediate. $\square$

Proof of Theorem 1(iv). Put $h_k = \mathbb{E}_k[(-1)^J]$. A private flip preserves the parity sign and a spine flip reverses it, whence

$$(18) \quad h_0 = 1, \quad h_k = \frac{2}{3}h_{k-1} - \frac{1}{3}h_{r-k}.$$

The affine values $(r+2-2k)/(r+2)$ satisfy every equation. Iterating the bounded first-step identity to $T \wedge n$ and using almost-sure absorption and bounded convergence identifies them with $\mathbb{E}_k[(-1)^J]$. The second identity in (9) follows from $\mathbb{E}[(-1)^J] = 1 - 2\mathbb{P}(J \text{ odd})$.

Let $R_0 = r+2$. Equations (7) and (9) give

$$m = \frac{k(R_0 - k)}{2}, \quad q(1 - q) = \frac{k(R_0 - k)}{R_0^2}.$$

Therefore (10) equals the positive integer R_0 , and $qR_0 = k$ is an integer in $[1, R_0 - 2]$. Conversely, if the stated feasibility conditions hold, the pair $r = R - 2$, $k = qR$ is admissible and the two displayed identities reproduce (m, q) . This proves both feasibility and uniqueness.

The central value $q = 1/2$ is regular: it occurs when r is even and $k = (r+2)/2$. Neither scalar alone suffices. Indeed,

$$(r, k) = (1, 1), (4, 2) \quad \text{both give } q = \frac{1}{3}, \quad (r, k) = (2, 2), (3, 1) \quad \text{both give } m = 2.$$

$\square$

5. EXACT PRESSURE, CONTROLS, AND LIMITATIONS

A standard-library verifier enumerates every nonzero imbalance vector through $r = 9$, solves independent rational Bellman systems, and compares them with the closed formulas. Table 2 records the frozen lanes.

TABLE 2. Deterministic exact-arithmetic falsification. Enumeration supports debugging; the preceding symbolic arguments prove the all-parameter theorem.

Lane	Exact assertions
Chebyshev elimination and transform vectors	4,416
Inverse states and absorption certificate	180,600
Literal strong lumpability	2,026
Mean, parity, and extrema	3,958
$r = 1$, $r = 2$, and $z = 0$ boundaries	278
Total	191,278

The exact control uses only integers and rational numbers. It is neither a proof of the theorem nor an ownership, novelty, priority, or release certificate. The primary-source screen is bounded, and any direct owner of the special-book law would require claim subtraction anew.

Internally adjacent objects are separated by the literal update. A triangular-book edge-deletion process removes edges; here every edge remains and the spine flip reflects k to $r - k$. Vertex-push orientation chains are stationary group walks, whereas this process selects an active imbalanced page and absorbs. Deterministic prefix-XOR feedback and unequal-spider first passage use different state spaces and proof engines; the bivariate spine-marked reflection law, rather than XOR notation or a mean alone, is the organizing object here.

LIMITATIONS

The results require uniform selection of active pages and physical edges, use the active update-epoch clock, and assume the known carrier $B(3, r) = K_{1,1,r}$. They do not cover the all-triad physical clock, nonuniform weights, arbitrary graphs, friendship/windmill carriers, noisy inverse data, or recovery of a full sign state. The inverse uses exact (m, q) and identifies only (r, k) .

DATA AVAILABILITY

No external data were used. The paper-local verifier and its frozen transcript contain the complete deterministic exact control.

ETHICS STATEMENT

This mathematical study involved no human participants, animals, personal data, or field intervention.

AUTHOR CONTRIBUTIONS

The anonymous author performed the mathematical derivations, exact checks, source-boundary audit, and manuscript preparation.

CONFLICT OF INTEREST

The author declares no conflict of interest.

FUNDING

No external funding is declared.

EXTERNAL STATUS

This artifact remains `HOLD_EXTERNAL`; it is not cleared for posting, submission, circulation, or author contact.

REFERENCES

- [1] T. Antal, P. L. Krapivsky, and S. Redner. Dynamics of social balance on networks. *Physical Review E*, 72(3):036121, 2005. doi: 10.1103/PhysRevE.72.036121. URL <https://arxiv.org/abs/cond-mat/0506476>.
- [2] T. Antal, P. L. Krapivsky, and S. Redner. Social balance on networks: The dynamics of friendship and enmity. *Physica D: Nonlinear Phenomena*, 224(1–2):130–136, 2006. doi: 10.1016/j.physd.2006.09.028. URL <https://arxiv.org/abs/physics/0605183>.
- [3] Gabriel Istrate. On the dynamics of social balance on general networks (with an application to XOR-SAT). *Fundamenta Informaticae*, 91(2):341–356, 2009. doi: 10.3233/FI-2009-0047. URL <https://arxiv.org/abs/0811.0381>.
- [4] Gabriel Istrate, Cosmin Bonchis, and Mircea Marin. Interactive particle systems on hypergraphs, drift analysis and the WalkSAT algorithm, 2019. URL <https://arxiv.org/abs/1909.12353>.
- [5] Deepak Sehrawat and Bikash Bhattacharjya. Chromatic polynomials of signed book graphs. *Theory and Applications of Graphs*, 9(1):Article 4, 2022. doi: 10.20429/tag.2022.090104. URL <https://arxiv.org/abs/2206.08580>.