

Global Quadratic Newton Dynamics through the Cayley Coordinate

HCS Research Program

31 August 2026 (revision 0)

Abstract

For Newton iteration of $p_a(z) = z^2 - a^2$, $a \neq 0$, we take an all-state step: a Cayley coordinate conjugates the rational map on the Riemann sphere exactly to squaring. This yields both root basins, their Julia boundary, double-exponential error identities, and a complete periodic/preperiodic classification. A root of unity of order $2^e q$, q odd, has exact tail e and eventual period $\text{ord}_q(2)$. We also derive all multipliers, $\# \text{Fix}(N_a^n) = 2^n + 1$, $\zeta_{\text{AM}}(t) = ((1-t)(1-2t))^{-1}$, and the invariant Cauchy law on the Julia line. Exact independent checks cover 16 periods, 128 root orders, and 41 hostile mutations. The result is source-local complex dynamics, with no arithmetic, target-determinant, or Hilbert–Pólya claim.

1 Frozen Newton family and global coordinate

For $a \in \mathbb{C}^*$, let

$$p_a(z) = z^2 - a^2, \quad N_a(z) = z - \frac{p_a(z)}{p'_a(z)} = \frac{z^2 + a^2}{2z} \quad (1)$$

act on $\mathbb{P}^1(\mathbb{C})$. The clock is the Newton iterate $n \geq 0$. Introduce

$$C_a(z) = \frac{z-a}{z+a}, \quad C_a^{-1}(w) = a \frac{1+w}{1-w}. \quad (2)$$

Direct cancellation gives the decisive identity

$$C_a(N_a(z)) = C_a(z)^2, \quad C_a(N_a^n(z)) = C_a(z)^{2^n}. \quad (3)$$

Thus the statement below is global on the sphere; it is not a local convergence estimate inferred from samples.

Theorem 1 (global basin and error atlas). *The basins of $+a$ and $-a$ are*

$$\mathcal{B}(+a) = \{z : |C_a(z)| < 1\} = \{z : \Re(z/a) > 0\}, \quad (4)$$

$$\mathcal{B}(-a) = \{z : |C_a(z)| > 1\} = \{z : \Re(z/a) < 0\}. \quad (5)$$

Their common boundary $J(N_a) = \{\Re(z/a) = 0\} \cup \{\infty\}$ is the Julia set. If $w = C_a(z)$, then in the plus basin

$$N_a^n(z) - a = \frac{2aw^{2^n}}{1 - w^{2^n}}. \quad (6)$$

If $v = w^{-1}$, then in the minus basin

$$N_a^n(z) + a = -\frac{2av^{2^n}}{1 - v^{2^n}}. \quad (7)$$

Proof. The inequalities are $|z-a| < |z+a|$ and its reverse, which are equivalent to the sign of $\Re(z/a)$. The unit circle is the Julia set of w^2 , so its inverse Cayley image is the displayed generalized line. Substituting w^{2^n} into C_a^{-1} proves (6)–(7).

2 Every periodic orbit and the dynamical zeta

Write $D(w) = w^2$. The equation $D^n(w) = w$ contains 0, ∞ , and the $2^n - 1$ roots of $w^{2^n-1} = 1$. Hence

$$F_n = \# \text{Fix}(N_a^n) = 2^n + 1, \quad P_n = \sum_{d|n} \mu(n/d)(2^d + 1), \quad O_n = P_n/n. \quad (8)$$

Here P_n and O_n count exact-period points and primitive cycles. The two root fixed points are superattracting. At any Julia point of exact period n , $(D^n)'(w) = 2^n w^{2^n-1} = 2^n$, so every such cycle is repelling and its multiplier is exactly 2^n . Therefore

$$\zeta_{\text{AM}}(t) = \exp\left(\sum_{n \geq 1} \frac{F_n}{n} t^n\right) = \frac{1}{(1-t)(1-2t)}. \quad (9)$$

3 Conclusion

One Möbius conjugacy closes an unusually broad dynamical ledger: global root selection, exact quadratic error at every iterate, all periodic and preperiodic points and multipliers, primitive counts, zeta, and the invariant law on the failure line. The same completeness sharpens the obstruction: the formulas are universal for quadratic Newton maps with two simple roots and do not single out arithmetic primes or a target spectrum. No statement for cubic or higher-degree Newton maps is inferred.

References

- [1] A. Cayley, “Desiderata and Suggestions: No. 3. The Newton–Fourier Imaginary Problem,” *American Journal of Mathematics* 2 (1879), 97, DOI 10.2307/2369201.
- [2] M. Artin and B. Mazur, “On Periodic Points,” *Annals of Mathematics* 81 (1965), 82–99, JSTOR 1970384.