

Primary Gaussian Phases and All-Degree Frobenius Dynamics for the Elliptic Curve $y^2 = x^3 - x$

HCS-C382 / HEN-O366

5 September 2026

Abstract

For the reductions of $y^2 = x^3 - x$ at every odd prime, we give a sign-complete derivation of the Frobenius eigenvalues. At split primes, the centralizer of an intrinsic order-four automorphism places Frobenius in the Gaussian order, and an explicit rational point of order four fixes the primary Gaussian unit. At inert primes a character involution gives trace zero. The resulting unordered eigenvalue pair is determined without a fitted phase or a point-count sign convention.

Keywords: complex multiplication; Frobenius dynamics; Gaussian integers; elliptic curves; primary normalization; rational torsion

中文摘要

本文统一处理曲线 $y^2 = x^3 - x$ 在所有奇素数处的约化。分裂情形中，四阶自同态的中心化子迫使弗罗贝尼乌斯位于高斯整数环；显式有理四阶点使群阶被八整除，从而唯一确定主高斯整数的实部符号。惯性情形通过二次特征的反号对合证明迹为零，整个相位约定无需拟合。
关键词：复乘；弗罗贝尼乌斯动力学；高斯整数；椭圆曲线；主元归一化；有理挠点

1 Frozen object, ownership, and the first advance

Round 0 advance: an all-odd-prime trace formula with a proved Gaussian-unit convention. Let p be an odd prime, let E_p denote the smooth projective curve $y^2 = x^3 - x$ over $\mathbf{F}_p$, and let

$$F : E_p(\overline{\mathbf{F}}_p) \longrightarrow E_p(\overline{\mathbf{F}}_p), \quad F(x, y) = (x^p, y^p).$$

This formula fixes the Frobenius convention; its cohomological pullback has eigenvalue p on degree two. We put $N_{p,n} = \#E_p(\mathbf{F}_{p^n})$ and $t_p = p + 1 - N_{p,1}$. The prime 2 is excluded: the integral model has discriminant 64. The parameter p indexes reductions of one curve and is not an autonomous real-time coordinate.

The general closed-point/zeta dictionary was already owned locally by flow papers 4 and 6 for a projective-line Frobenius suspension. The earlier C41 cubic-CM package considered $y^2 = x^3 + 1$, an inert-prime trace law and finite extension checks. The present owner is the quartic-CM all-prime primary phase, its ordinary/supersingular distinction, and its complete degree consequences. These results specialize classical elliptic-curve theory; they are not presented as newly discovered CM theorems. We use the endomorphism and finite-field results in Milne [1], with every special sign step proved below.

Theorem 1 (Sign-complete CM trace). *If $p \equiv 3 \pmod{4}$, then $t_p = 0$. If $p \equiv 1 \pmod{4}$, there are unique integers a, b satisfying*

$$b > 0, \quad a \text{ odd}, \quad b \text{ even}, \quad a^2 + b^2 = p, \quad a + b \equiv 1 \pmod{4},$$

and $t_p = 2a$. The unordered Frobenius eigenvalues are $a \pm bi$ in the split case and $\pm i\sqrt{p}$ in the inert case.

Proof. Let χ be the quadratic character extended by $\chi(0) = 0$. Counting the two square roots of a nonzero square gives

$$N_{p,1} = p + 1 + \sum_{x \in \mathbf{F}_p} \chi(x^3 - x).$$

When $p \equiv 3 \pmod{4}$, replacing x by $-x$ negates the sum, so it vanishes as an integer. This includes the good prime $p = 3$.

For $p \equiv 1 \pmod{4}$, choose $j \in \mathbf{F}_p$ with $j^2 = -1$. The endomorphism $I(x, y) = (-x, jy)$ satisfies $I^2 = [-1]$ and commutes with F . By the elliptic endomorphism classification, the endomorphism algebra containing $\mathbf{Q}(I)$ is either that quadratic field or a quaternion algebra. In either case the centralizer of I is $\mathbf{Q}(I)$: in the latter case write the algebra as $\mathbf{Q}(i) \oplus \mathbf{Q}(i)v$ with $vi = -iv$ and compare the two coefficients. Thus F lies in $\mathbf{Q}(i)$. Its integrality places it in the maximal order $\mathbf{Z}[i]$, so $F = a + bi$, and degree and trace give $a^2 + b^2 = p$ and $t_p = 2a$.

All three nonzero 2-torsion points are rational. Therefore $F - [1]$ kills $E[2]$ and factors through the separable isogeny $[2]$. The endomorphism $(F - [1])/2$ is again an integral element of $\mathbf{Q}(i)$. It follows that a is odd and b even.

Now $Q = (j, j - 1)$ lies on the curve because $(j - 1)^2 = -2j = j^3 - j$. The tangent slope is

$$m = \frac{3j^2 - 1}{2(j - 1)} = j + 1, \quad x(2Q) = m^2 - 2j = 0, \quad y(2Q) = mj - (j - 1) = 0.$$

Hence $2Q = (0, 0)$, and Q has order four. The independent point $(1, 0)$ of order two gives an eight-element subgroup, so $8 \mid p + 1 - 2a$. If $b \equiv 0 \pmod{4}$ then $p \equiv 1 \pmod{8}$ and $a \equiv 1 \pmod{4}$; if $b \equiv 2 \pmod{4}$ then $p \equiv 5 \pmod{8}$ and $a \equiv 3 \pmod{4}$. These are precisely $a + b \equiv 1 \pmod{4}$. Unique factorization in $\mathbf{Z}[i]$ fixes $|a|, |b|$ and this congruence fixes the sign of a . Taking $b > 0$ chooses the upper display root. It does not select one literal prime ideal: both conjugates satisfy the primary congruence modulo $(1 + i)^3$. $\square$

p	a	b	t_p	chamber
3	—	—	0	supersingular
5	-1	2	-2	ordinary
13	3	2	6	ordinary
17	1	4	2	ordinary
29	-5	2	-10	ordinary

The negative real part at $p = 5$ is essential: choosing the positive representation $1 + 2i$ changes the curve trace. Complex conjugation, however, changes neither trace nor the unordered eigenvalue pair.

2 Conclusion

Rational four-torsion fixes the primary Gaussian sign uniformly at every split odd prime, while a character involution settles every inert prime. This proof identifies the exact source mechanism behind the phase convention.

References

- [1] J. S. Milne, *Elliptic Curves*, 2nd ed., World Scientific, 2020. II §6 (endomorphisms), III Example 3.18 (quartic CM), IV §9, Theorems 9.4 and 9.10, Remark 9.11 and Exercise 9.12 (finite-field counts, zeta and functional equation). <https://www.jmilne.org/math/Books/EC2.pdf>.