

The Chaplygin Sleigh for Every Signed Mass Offset: Exact Scattering, Physical Reconstruction, and the Recurrence Boundary

Route-A structural certificate C199

Abstract

We give a convention-complete theorem for the classical Chaplygin sleigh with mass $m > 0$, central inertia $J > 0$, and arbitrary signed longitudinal offset a . For $a \neq 0$, every non-equilibrium positive-energy reduced orbit is an explicit heteroclinic: its blade rotates through an energy-independent angle and its contact point has two rigorously defined asymptotic lines. We also classify the stable equilibrium half-axis, half-plane Poisson structure, necessarily singular reduced density and its configuration-Haar lift, reversor, and all zero-offset and zero-angular-velocity boundaries. A finite exact/high-precision ledger checks conventions but is not used to prove the continuous theorem. The Poisson form gives only a formal Route-A hint; the absence of an intrinsic prime carrier and target determinant forces rejection.

Keywords: Chaplygin sleigh; nonholonomic mechanics; heteroclinic scattering; reconstruction; invariant measure; reversibility.

中文摘要

本文对质量、转动惯量均为正且质心纵向偏置可取任意符号的经典 Chaplygin sleigh 给出完整定理。偏置非零时，全部非平衡正能轨道均为显式异宿轨道；刀刃转角与能量无关，接触点具有两条严格定义的渐近直线。本文还统一分类稳定平衡半轴、半平面泊松结构、必然奇异的不变测度、时间反演以及零偏置等全部边界。有限精确计算只校验约定，不代替连续参数证明；路线 Δ 最终仍被拒绝。

关键词：非完整力学；异宿散射；重构；不变测度；时间反演。

1 Convention, equations, and ownership

Let $r = (x, y)$ be the knife contact point, $\theta \in \mathbb{S}^1$ the *blade* angle, u the signed contact velocity along the blade, and $\omega = \dot{\theta}$. Let $a \in \mathbb{R}$ be the signed longitudinal offset from contact point to centre of mass and put $I_c = J + ma^2$. In physical time, the reduced equations and reconstruction are

$$\dot{r} = ue(\theta), \quad \dot{\theta} = \omega, \quad \dot{u} = a\omega^2, \quad \dot{\omega} = -\frac{ma}{I_c}u\omega, \quad e(\theta) = (\cos \theta, \sin \theta). \quad (1)$$

They imply

$$H = \frac{m}{2}u^2 + \frac{I_c}{2}\omega^2 = \text{constant}. \quad (2)$$

The model, its nonholonomic reduction and Hamiltonization context are classical [1, 2, 3]. We claim a signed-parameter synthesis and certificate, not priority for those results.

2 All nonzero-offset scattering orbits

Theorem 1 (signed-offset normal form). *Assume $a \neq 0$, $H > 0$, and $\omega(0) \neq 0$. Define*

$$R = \sqrt{2H/m}, \quad A = \frac{m|a|R}{I_c}, \quad \eta = \frac{\sqrt{I_c/m}}{|a|}, \quad \sigma = \text{sgn } \omega(0).$$

There are unique $t_0 \in \mathbb{R}$ and θ_0 modulo 2π such that, with $s = A(t - t_0)$,

$$u = \text{sgn}(a)R \tanh s, \quad \omega = \sigma R \sqrt{m/I_c} \text{sech } s, \quad \theta = \theta_0 + \sigma \eta \arcsin(\tanh s). \quad (3)$$

It joins $(-\text{sgn}(a)R, 0)$ to $(\text{sgn}(a)R, 0)$ and its blade-angle deflection is

$$\Delta\theta = \theta_+ - \theta_- = \sigma\pi\eta, \quad (4)$$

which is independent of H .

Proof. The sign of a nonzero ω cannot change by uniqueness. On the energy ellipse write $u = \text{sgn}(a)R \tanh s$ and $\omega = \sigma R \sqrt{m/I_c} \text{sech } s$. The first reduced equation gives $\dot{s} = A$; substitution gives the second. Since $d\{\arcsin(\tanh s)\}/ds = \text{sech } s$, integration of $\dot{\theta} = \omega$ proves (3). Taking $s \rightarrow \pm\infty$ proves both endpoints and (4). $\square$

The endpoint table makes the signed convention explicit:

	$t \rightarrow -\infty$	$t \rightarrow +\infty$
u	$-\text{sgn}(a)R$	$\text{sgn}(a)R$
ω	0^σ	0^σ
θ	$\theta_0 - \sigma\eta\pi/2$	$\theta_0 + \sigma\eta\pi/2$

3 Contact-point reconstruction and two lines

Put $q = \tanh s \in (-1, 1)$. Since $dt = dq/[A(1 - q^2)]$, the full $SE(2)$ reconstruction reduces exactly to

$$r(q) - r(0) = \frac{\text{sgn}(a)R}{A} \int_0^q \frac{y \mathbf{e}(\theta_0 + \sigma \eta \arcsin y)}{1 - y^2} dy. \quad (5)$$

This is a one-dimensional quadrature, not a claim of an elementary primitive.

Proposition 1 (affine asymptotes). *Let $v_{\pm} = u_{\pm} \mathbf{e}(\theta_{\pm})$. There are unique vectors $b_{\pm}$ such that*

$$r(t) = b_{\pm} + v_{\pm} t + o(1) \quad (t \rightarrow \pm\infty).$$

Thus the contact trajectory scatters between two oriented affine lines.

Proof. As $s \rightarrow +\infty$, $u - u_+ = O(e^{-2s})$ while $\theta - \theta_+ = O(e^{-s})$; hence $u \mathbf{e}(\theta) - v_+ = O(e^{-s})$, integrable in physical time because $s = At + O(1)$. Therefore $r(t) - v_+ t$ has a finite limit. The negative end is identical. Uniqueness follows by subtracting two candidate intercepts. $\square$

The blade angle is not always the velocity heading. The physical heading is θ when $u > 0$ and $\theta + \pi$ when $u < 0$ (modulo 2π). Therefore (4) must not be silently relabelled as a velocity-heading deflection.

4 Stability, Poisson form, measure, and reversibility

The entire line $\omega = 0$ is reduced equilibrium. At $(u_*, 0)$ its linearization has eigenvalues 0 and

$$\lambda_{\perp} = -\frac{ma}{I_c} u_*. \quad (6)$$

Hence precisely $au_* > 0$ is transversely stable. On either open half-plane,

$$\{u, \omega\} = \frac{a}{I_c} \omega \quad (7)$$

is a nondegenerate Poisson bracket and $\dot{f} = \{f, H\}$ reproduces the reduced flow. The density $\rho = 1/|\omega|$ satisfies $\nabla \cdot (\rho \dot{u}, \rho \dot{\omega}) = 0$ there.

No positive C^1 reduced invariant density can cross a nonzero reduced equilibrium: at $(u_*, 0)$, invariance would give $0 = \rho \nabla \cdot F = -\rho m a u_*/I_c$, a contradiction. Thus the reduced singularity is structural, not a removable normalization. Multiplying by configuration Haar volume gives an invariant full-flow measure off the singular line, and no smooth density of this Haar-factor form crosses it. However, $(u_*, 0)$ with $u_* \neq 0$ still reconstructs as translation in the full five-dimensional flow. The pointwise reduced argument therefore does not exclude every possible configuration-dependent full-flow C^1 density.

Finally,

$$\mathcal{R}(r, \theta, u, \omega) = (r, \theta, -u, -\omega)$$

satisfies $F(\mathcal{R}z) = -D\mathcal{R}F(z)$, so it is a physical-time reversor.

5 All degenerate boundaries and recurrence

If $a \neq 0$ but $\omega_0 = 0$, the reduced state is fixed and r is a straight line; $H = 0$ is the fully fixed case. Otherwise $\dot{u} = a\omega^2$ is strictly one-signed, excluding nonconstant recurrence.

If $a = 0$, both u and ω are constant. When $\omega \neq 0$,

$$r(t) - r(0) = \frac{u}{\omega} (\sin(\theta_0 + \omega t) - \sin \theta_0, -\cos(\theta_0 + \omega t) + \cos \theta_0),$$

and the complete $SE(2)$ state is periodic with period $2\pi/|\omega|$. When $\omega = 0$, motion is straight (fixed if also $u = 0$). Thus $a = 0$ is a sharp recurrence boundary, not a regular member of the scattering family.

6 Independent certificate and strict Route-A stop

The deterministic ledger uses six rational (m, J, a, H) families, both signs of a and ω , 12 heteroclinic cases, 36 states, and four $a = 0$ boundaries. An independent checker imports no producer and closes 737 assertions under recursively exact schema keys. A separate SymPy derivation checks 25 structural and 36 rational hyperbolic identities. Replay is byte exact; twelve repaired-hash attacks (including unknown-key and four mathematical attacks) plus one stale-hash attack are rejected. These tests audit signs and conventions; Theorem 1 and the proofs above establish the continuous result.

The half-plane bracket and the quantum comparison in [4] motivate only a formal operator hint. There is no intrinsic rational-prime primitive-orbit ledger, prime-power repetition law, target Fredholm determinant, target divisor/functional equation, or constructed same-clock operator with the required spectrum. Consequently

$$(A0, A1, A2, A3, A4) = (\text{FAIL}, \text{FAIL}, \text{FAIL}, \text{FAIL}, \text{FORMAL HINT}),$$

overall `ROUTE_A_REJECTED`, Route B false, under `NO_BAD_EULER_OR_ROOT_NUMBER`.

Claim firewall. We claim no arithmetic local data, Euler factor, root number, automorphy, target analytic structure, Hilbert–Pólya operator, a smooth global reduced or configuration-Haar-factor density, exclusion of all configuration-dependent full-flow densities, finite-regression proof, exhaustive priority, external review, or acceptance score.

Revision focus. Round 2 adds independent executable closure, source ownership, Route-A evaluation, counterclaim firewall and disclosures.

Declarations

Data and code availability. Package-local synthetic exact evidence and deterministic code accompany the manuscript; no observational data are used.

Ethics. No human, animal, personal or sensitive data are used; approval is not applicable.

Author contributions (CRediT). This anonymous certificate records Conceptualization, Formal analysis, Software, Validation, Writing—original draft, and Writing—review and editing. AI systems are not authors.

Funding. No external funding is reported.

Conflicts of interest. None known.

AI-use disclosure. An AI coding assistant supported derivation, drafting and exact-code development; it was not an external reviewer or an independent peer-review process.

References

- [1] A. M. Bloch, “Asymptotic Hamiltonian dynamics: the Toda lattice, the three-wave interaction and the non-holonomic Chaplygin sleigh,” *Physica D* 141 (2000), 297–315. DOI: 10.1016/S0167-2789(00)00046-4.
- [2] A. V. Borisov and I. S. Mamaev, “The dynamics of a Chaplygin sleigh,” *J. Appl. Math. Mech.* 73 (2009), 156–161. DOI: 10.1016/j.jappmathmech.2009.04.005.
- [3] N. K. Moshchuk, “On the motion of Chaplygin’s sledge,” *J. Appl. Math. Mech.* 51 (1987), 426–430. DOI: 10.1016/0021-8928(87)90079-7.
- [4] A. M. Bloch and A. G. Rojo, “Quantization of a Nonholonomic System,” *Phys. Rev. Lett.* 101 (2008), 030402. DOI: 10.1103/PhysRevLett.101.030402.