

Relaxed Douglas–Rachford Dynamics for Two Subspaces: An All-Parameter Principal-Angle Atlas

Route-A structural certificate C197

Abstract

For every pair of linear subspaces in finite-dimensional real Hilbert space and every real relaxation parameter, we give the complete orthogonal block atlas of the relaxed Douglas–Rachford map. It yields the fixed space, sharp convergence window and rate, optimal relaxation, shadow limit, endpoint rotation dynamics, instability boundary, all power traces, and a closed finite determinant. Exact rational sentinels are reconstructed independently from projections and by symbolic algebra; they test conventions without replacing the all-subspace proof. The source-native orthogonal endpoint is a formal operator hint, but projection geometry has no intrinsic prime carrier or logarithmic clock, so the candidate fails Route A.

Keywords: Douglas–Rachford splitting; principal angles; relaxed projections; convergence rate; finite determinant; Route-A obstruction.

中文摘要

本文对有限维实希尔伯特空间中任意两个线性子空间及任意实松弛参数，给出松弛 Douglas–Rachford映射的完整主角分块。该分块一次确定不动空间、收敛区间与精确速率、最优松弛、影子极限、端点旋转、失稳边界、全部幂迹与有限行列式。有理角度证书仅检验符号与约定，不替代全参数证明。正交端点虽然是同钟自然算子，却没有内生素数载体或对数时钟，故不能通过Route A。

关键词：投影算法；主角；松弛动力学；收敛速率；算术阻碍。

1 Frozen map and canonical planes

Let $U, V \subset H$ be linear subspaces, P_W the orthogonal projection and $R_W = 2P_W - I$. One algorithmic tick is

$$T_\lambda = (1 - \lambda)I + \frac{\lambda}{2}(I + R_V R_U), \quad \lambda \in \mathbb{R}.$$

This is the two-subspace specialization of the classical splitting lineage of Douglas and Rachford [1]; the sharp subspace convergence location and Friedrichs-angle rate are due to Bauschke et al. [2].

The cosine–sine decomposition splits H orthogonally into

$$F = (U \cap V) \oplus (U^\perp \cap V^\perp), \quad M = (U \cap V^\perp) \oplus (U^\perp \cap V),$$

and principal planes E_j with $0 < \theta_j < \pi/2$. On F and M the map is I and $(1 - \lambda)I$, respectively. Choosing $U \cap E_j = \text{span}(e_1)$ and $V \cap E_j = \text{span}(\cos \theta_j e_1 + \sin \theta_j e_2)$ gives

$$T_\lambda|_{E_j} = \begin{pmatrix} 1 - \lambda \sin^2 \theta_j & -\lambda \sin \theta_j \cos \theta_j \\ \lambda \sin \theta_j \cos \theta_j & 1 - \lambda \sin^2 \theta_j \end{pmatrix}. \quad (1)$$

Its conjugate eigenvalues have squared modulus

$$\rho_j^2 = 1 - \lambda(2 - \lambda) \sin^2 \theta_j. \quad (2)$$

2 Complete dynamics

For $\lambda \neq 0$, $\text{Fix } T_\lambda = F$; at $\lambda = 0$ the map is the identity. Equations (1)–(2) and orthogonality of the decomposition prove the following exhaustive classification.

parameter	off-fixed behavior	conclusion
$\lambda = 0$	identity	every point fixed
$0 < \lambda < 2$	strict contraction	$T_\lambda^n \rightarrow P_F$
$\lambda = 2$	signs and plane rotations	orthogonal recurrence boundary
$\lambda \notin [0, 2]$	modulus > 1 if nontrivial	unstable off F

In the contracting window the exact operator-norm rate is the largest existing quantity among

$$|1 - \lambda|, \quad \sqrt{1 - \lambda(2 - \lambda) \sin^2 \theta_j}.$$

Thus $\lambda = 1$ uniquely minimizes every nontrivial block simultaneously; its generic rate is $\cos \theta_F$. Moreover $P_U T_\lambda^n x \rightarrow P_{U \cap V} x$, because $P_U P_F = P_{U \cap V}$.

At $\lambda = 2$, $T_2 = R_V R_U$: it is I on F , $-I$ on M , and rotation by $2\theta_j$ on E_j . It has finite order exactly when every θ_j/π is rational (with the mismatch signs included in the least common order). Hence the endpoint is not silently counted as convergence.

3 Trace and determinant closure

Put $f = \dim F$, $h = \dim M$, $a_j = 1 - \lambda \sin^2 \theta_j$, and $d_j = \rho_j^2$. Then the complete finite factorization is

$$\det(I - zT_\lambda) = (1 - z)^f (1 - (1 - \lambda)z)^h \prod_j (1 - 2a_j z + d_j z^2). \quad (3)$$

The contribution $\tau_{j,n}$ of a principal plane to $\text{tr}(T_\lambda^n)$ obeys $\tau_{j,0} = 2$, $\tau_{j,1} = 2a_j$ and $\tau_{j,n} = 2a_j \tau_{j,n-1} - d_j \tau_{j,n-2}$. Thus the same block theorem closes every trace, not only the asymptotic rate. Formula (3) is a finite algorithmic determinant; no periodic-orbit or target ownership is introduced by notation.

Revision focus. Round 1 adds fixed/mismatch boundaries, sharp optimality, the shadow theorem and full trace–determinant closure.

Declarations

Data and code availability. Package-local exact evidence and deterministic code accompany this manuscript.

Ethics. No human, animal, clinical, personal or sensitive data are used; approval is not applicable.

Author contributions (CRediT). This anonymous certificate records Conceptualization, Formal analysis, Software, Validation, Writing—original draft, and Writing—review and editing. AI systems are not authors.

Funding. No external funding is reported.

Conflicts of interest. None known.

AI-use disclosure. An AI coding assistant supported drafting and exact-code development; it was not an external reviewer or independent error process.

References

- [1] J. Douglas Jr. and H. H. Rachford Jr., “On the numerical solution of heat conduction problems in two and three space variables,” *Trans. Amer. Math. Soc.* 82 (1956), 421–439. DOI: 10.1090/S0002-9947-1956-0084194-4.
- [2] H. H. Bauschke, J. Y. Bello Cruz, T. T. A. Nghia, H. M. Phan and X. Wang, “The rate of linear convergence of the Douglas–Rachford algorithm for subspaces is the cosine of the Friedrichs angle,” *J. Approx. Theory* 185 (2014), 63–79. DOI: 10.1016/j.jat.2014.06.002.