

The Closed SIR Flow for All Positive Parameters: Phase Curves, Final-Size Branches, and the No-Recurrence Boundary

Route-A structural certificate C198

Abstract

We give one complete dynamical atlas for the closed mass-action SIR system at all positive transmission and removal rates. Exact scaling reduces the family to one planar flow; its first integral yields the phase curves, infection peak, time quadrature, final-size sensitivity and the correctly selected Lambert branch. A monotone removed coordinate proves global convergence and excludes every nonconstant recurrent orbit. A Lambert-free high-precision checker independently reconstructs both real final-size intersections. The result is an idealized mathematical certificate, with no clinical data or medical advice. Its monotonicity and absence of intrinsic prime arithmetic force strict Route-A rejection.

Keywords: SIR model; phase portrait; final size; Lambert W; threshold; monotone dynamics; Route-A obstruction.

中文摘要

本文对全部正传播率与移除率的封闭质量作用SIR系统给出统一动力学图谱。精确无量纲化把全参数族化为一个平面流；第一积分同时确定相曲线、感染峰值、时间积分、最终规模灵敏度及正确的Lambert W分支。移除变量严格单调，因而全局收敛并排除全部非常值回归轨道。本文只研究理想化数学模型，不使用临床数据，也不提供医疗建议。
关键词：SIR模型；相图；最终规模；阈值；单调动力学。

1 One phase portrait for all parameters

Fix $\beta, \gamma > 0$ and physical time t in

$$\dot{S} = -\beta SI, \quad \dot{I} = \beta SI - \gamma I, \quad \dot{R} = \gamma I.$$

The model and its threshold/final-size analysis descend from Kermack and McKendrick [1]; Lambert-branch analysis is treated explicitly by Pakes [2]. Put

$$\kappa = \gamma/\beta, \quad x = S/\kappa, \quad y = I/\kappa, \quad z = R/\kappa, \quad \tau = \gamma t.$$

Every parameter pair is thereby conjugate, without changing physical-time orientation, to

$$x' = -xy, \quad y' = y(x-1), \quad z' = y. \quad (1)$$

The nonnegative fixed-population simplex is invariant and compact, so solutions are global. Direct differentiation gives

$$H(x, y) = x + y - \log x = H(x_0, y_0), \quad y = y_0 + x_0 - x + \log(x/x_0). \quad (2)$$

For $y_0 > 0$, x decreases strictly. If $x_0 > 1$, then y grows until the unique crossing $x = 1$ and

$$y_{\max} = y_0 + x_0 - 1 - \log x_0.$$

If $x_0 < 1$, y decreases from the start; at $x_0 = 1$ its first derivative is zero, after which decreasing x makes it negative. Thus the threshold and peak are globally, not locally, classified.

Revision focus. Round 0 freezes the physical model and proves the all-parameter scaling, first integral and complete peak atlas.

Declarations

Data and code availability. Package-local synthetic exact evidence and deterministic code accompany this manuscript; no clinical data are used.

Ethics. No human, animal, clinical, personal or sensitive data are used; approval is not applicable.

Author contributions (CRediT). This anonymous certificate records Conceptualization, Formal analysis, Software, Validation, Writing—original draft, and Writing—review and editing. AI systems are not authors.

Funding. No external funding is reported.

Conflicts of interest. None known.

AI-use disclosure. An AI coding assistant supported drafting and exact-code development; it was not an external reviewer or independent error process.

References

- [1] W. O. Kermack and A. G. McKendrick, “A contribution to the mathematical theory of epidemics,” *Proc. R. Soc. A* 115 (1927), 700–721. DOI: 10.1098/rspa.1927.0118.
- [2] A. G. Pakes, “Lambert’s W meets Kermack–McKendrick epidemics,” *IMA J. Appl. Math.* 80 (2015), 1368–1386. DOI: 10.1093/imamat/hxu057.