

Conductor-Shell p-Adic Fractional Heat Semigroup: Exact Zeta, Scale Oscillation, and Boundary Closure

Route-A source-local certificate HCS-C283

1 September 2026

Abstract

For every fixed prime p and $\alpha > 0$, we define a nonnegative operator on $L^2(\mathbb{Z}_p)$ directly as the Fourier multiplier 0 on constants and $p^{\alpha n}$ on characters of exact conductor n . Its spectrum and a conditional-expectation reconstruction prove a positive conservative Markov semigroup with an exact heat trace. The fixed local prime is not promoted to all-prime target arithmetic.

1 The frozen conductor multiplier

Let p be any fixed rational prime, let $G = \mathbb{Z}_p$ have Haar mass one, and use $\widehat{G} = \mathbb{Q}_p/\mathbb{Z}_p$. A nonzero class ξ has exact conductor $n(\xi) = n$ when χ_ξ is trivial on $p^n\mathbb{Z}_p$ but not on $p^{n-1}\mathbb{Z}_p$. Exactly $p^n - p^{n-1} = (p-1)p^{n-1}$ characters have conductor n .

For $\alpha > 0$ we freeze

$$D_{p,\alpha}1 = 0, \quad D_{p,\alpha}\chi_\xi = p^{\alpha n(\xi)}\chi_\xi, \quad (1)$$

with its maximal diagonal domain. For $\mu \geq 0$, set $A_{p,\alpha,\mu} = D_{p,\alpha} + \mu I$ and $T_t = e^{-tA_{p,\alpha,\mu}}$. Equation (1) is the normalization: the paper does not identify it with every convention called a Vladimirov operator. Pseudodifferential spectra over p-adic fields, p-adic wavelet eigenbases, and ultrametric Markov semigroups form the surrounding literature [1, 2, 3, 5]. More directly, Example 5.1 of Chacón-Cortés-Zúñiga-Galindo [4], at dimension one and exponent α , already gives exactly our positive shell spectrum, multiplicities, geometric zeta, and vertical pole lattice. Those receive zero priority credit. We retain a source-local Markov, scale-boundary, determinant, and finite-reconstruction closure, not a literature-novelty claim.

Let E_n denote conditional expectation onto functions constant on cosets of $p^n\mathbb{Z}_p$, so $E_0 = P_0$ projects onto constants and $E_n \uparrow I$ strongly. Put

$$W_0 = \text{ran } E_0, \quad W_n = \text{ran } E_n \ominus \text{ran } E_{n-1} \quad (n \geq 1).$$

Then $\dim W_n = (p-1)p^{n-1}$ and the character and filtration descriptions give the same orthogonal decomposition.

Theorem 1 (spectrum and Markov reconstruction). *The spectrum of $D_{p,\alpha}$ consists of the simple eigenvalue zero and $p^{\alpha n}$ with multiplicity $(p-1)p^{n-1}$, $n \geq 1$. Its resolvent is compact. The spectrum of $A_{p,\alpha,\mu}$ is the simple eigenvalue μ and $\mu + p^{\alpha n}$ with the same multiplicities. Moreover $e^{-tD_{p,\alpha}}$ is a positive, conservative, self-adjoint contraction semigroup, while T_t is positive sub-Markov and $\|T_t\| = e^{-\mu t}$.*

Proof. Character counting and (1) prove the spectral assertion; the eigenvalues tend to infinity. A second reconstruction is

$$D_{p,\alpha} = \sum_{n \geq 1} p^{\alpha n} P_{W_n} = \sum_{n \geq 0} a_n (I - E_n), \quad (2)$$

where $a_0 = p^\alpha$ and $a_n = p^{\alpha(n+1)} - p^{\alpha n}$ for $n \geq 1$. On W_m , the right side is $\sum_{n < m} a_n = p^{\alpha m}$. For every conditional expectation E ,

$$e^{-sa(I-E)} = E + e^{-sa}(I-E)$$

is positive, constant-preserving, and contractive. The finite factors in (2) commute; their strong limit proves the Markov claims. The scalar killing factor $e^{-\mu t}$ gives the final norm and sub-Markov statement. $\square$

This proof derives positivity from the residue filtration. Nonnegative eigenvalues alone would not have established positivity preservation.

References

- [1] V. S. Vladimirov, *On the spectrum of some pseudodifferential operators over the field of p -adic numbers*, Algebra i Analiz 2:6 (1990), 107–124; Leningrad Math. J. 2:6 (1991), 1261–1278, [MathNet aa223](#).
- [2] S. V. Kozyrev, *Wavelet theory as p -adic spectral analysis*, Izv. Math. 66:2 (2002), 367–376, [DOI record](#).
- [3] A. D. Bendikov, A. A. Grigor'yan, Ch. Pittet, and W. Woess, *Isotropic Markov semigroups on ultra-metric spaces*, Russian Math. Surveys 69:4 (2014), 589–680, [DOI record](#).
- [4] L. F. Chacón-Cortés and W. A. Zúñiga-Galindo, *Heat traces and spectral zeta functions for p -adic Laplacians*, St. Petersburg Math. J. 29:3 (2018), 529–544, [DOI record](#).
- [5] V. S. Vladimirov, I. V. Volovich, and E. I. Zelenov, *p -Adic Analysis and Mathematical Physics*, World Scientific, 1994.