

Exponential Shot-Noise Ornstein–Uhlenbeck Dynamics: Exact Semigroup, Gamma Equilibrium, and Polynomial Filtrations

Route-A source-local certificate HCS-C335

3 September 2026

Abstract

We close the positive Ornstein–Uhlenbeck process driven by Poisson arrivals with exponential marks. Its all-time transition Laplace transform follows directly from the marked-Poisson functional.

1 Frozen process and semigroup

Fix $\gamma, \kappa, \beta > 0$. Let N_t be Poisson of rate κ , let T_j be its jump times, and let the independent marks Y_j have exponential rate β . On $[0, \infty)$ consider

$$dX_t = -\gamma X_t dt + dJ_t, \quad J_t = \sum_{T_j \leq t} Y_j, \quad X_0 = x \geq 0. \quad (1)$$

Theorem 1 (Exact affine Markov closure). *For $t, s \geq 0$, with $\alpha = \kappa/\gamma$,*

$$X_t = e^{-\gamma t} x + \sum_{T_j \leq t} e^{-\gamma(t-T_j)} Y_j, \quad (2)$$

$$\mathbb{E}_x e^{-sX_t} = e^{-se^{-\gamma t} x} \left(\frac{\beta + se^{-\gamma t}}{\beta + s} \right)^\alpha. \quad (3)$$

The unique invariant probability is Gamma with shape α and rate β . For every $p \geq 1$,

$$W_p(P_t(x, \cdot), P_t(y, \cdot)) = e^{-\gamma t} |x - y|. \quad (4)$$

In stationarity,

$$\mathbb{E} X^n = \frac{(\alpha)_n}{\beta^n}, \quad \text{cum}_n(X) = \frac{\alpha(n-1)!}{\beta^n}, \quad \text{Cov}(X_t, X_0) = \frac{\alpha}{\beta^2} e^{-\gamma|t|}. \quad (5)$$

Finally, the generator preserves $\mathcal{P}_m = \text{span}\{1, x, \dots, x^m\}$, and its restriction has exactly the simple eigenvalues $0, -\gamma, \dots, -m\gamma$. This last assertion concerns each fixed finite-dimensional filtration only; it is not a statement about the full L^2 spectrum, completeness, normality, or reversibility.

Proof of the path and transform. Variation of constants between successive jumps proves (2). The Laplace functional of a marked Poisson process gives

$$\begin{aligned} \log \mathbb{E} \exp \left[-s \sum_{T_j \leq t} e^{-\gamma(t-T_j)} Y_j \right] &= \kappa \int_0^t \left(\frac{\beta}{\beta + se^{-\gamma u}} - 1 \right) du \\ &= \frac{\kappa}{\gamma} \log \frac{\beta + se^{-\gamma t}}{\beta + s}. \end{aligned}$$

Multiplication by the deterministic initial term proves (3). Factoring the right side first at decay $r_1 = e^{-\gamma t_1}$ and then at $r_2 = e^{-\gamma t_2}$ cancels the intermediate denominator and leaves $r_1 r_2$, proving the Chapman–Kolmogorov law directly. $\square$

The phrase *shot-noise Laplace semigroup owner* is the round-zero certificate. Formula (3), not a simulation, owns the transition law.

References

- [1] G. Kotler, V. Kontorovich, V. Lyandres, and S. Primak, *On the Markov model of shot noise*, Signal Processing (1999), [doi:10.1016/S0165-1684\(98\)00226-6](https://doi.org/10.1016/S0165-1684(98)00226-6).
- [2] K. Sato and M. Yamazato, *Operator-selfdecomposable distributions as limit distributions of processes of Ornstein–Uhlenbeck type*, Stochastic Processes Appl. 17 (1984), 73–100, [doi:10.1016/0304-4149\(84\)90312-0](https://doi.org/10.1016/0304-4149(84)90312-0).
- [3] O. E. Barndorff-Nielsen and N. Shephard, *Non-Gaussian Ornstein–Uhlenbeck-based models and some of their uses in financial economics*, J. Roy. Statist. Soc. B 63 (2001), 167–241, [doi:10.1111/1467-9868.00282](https://doi.org/10.1111/1467-9868.00282).