

Exact Functional Graphs of Finite Linear Maps from Rational Canonical Data

HCS Research Program

27 August 2026 (revision 2)

Abstract

For every linear endomorphism of a finite-dimensional vector space over a finite field, we derive its complete functional-graph census from its invariant factors. The result simultaneously covers nilpotent, nonsemisimple, and inseparable cases. It gives every fixed count, the periodic subspace and maximal preperiod, exact periods, the Artin–Mazur zeta function, and the characteristic polynomial of composition on the full function space. Exact enumeration, independent rank calculations, and symbolic controls verify eight structurally distinct witnesses, including genuine $\mathbb{F}_4$ arithmetic.

1 Invariant-factor fixed counts

Let $A \in \text{End}_{\mathbb{F}_q}(V)$ have invariant factors $f_1 \mid \cdots \mid f_r$, so that $V \simeq \bigoplus_i \mathbb{F}_q[X]/(f_i)$ and A acts as multiplication by X . Put

$$F_n = q^{\sum_i \deg \gcd(f_i, X^n - 1)}.$$

Theorem 1 (complete finite dynamics). *For every $n \geq 1$, $|\text{Fix}(A^n)| = F_n$. Write $f_i = X^{e_i} g_i$ with $g_i(0) \neq 0$. Then $\text{Per}(A)$ is a vector subspace of dimension $\sum_i \deg g_i$, and the largest preperiod is $\max_i e_i$. The numbers of least-period points and cycles are*

$$P_n = \sum_{d|n} \mu(n/d) F_d, \quad C_n = P_n/n,$$

and $\zeta_A(z) = \prod_{n \geq 1} (1 - z^n)^{-C_n}$. For $U_A \phi = \phi \circ A$ on $\mathbb{C}^V$,

$$\chi_{U_A}(T) = T^{|V| - |\text{Per}(A)|} \prod_{n \geq 1} (T^n - 1)^{C_n}.$$

Proof. On $\mathbb{F}_q[X]/(f_i)$, the kernel of multiplication by $X^n - 1$ has dimension $\deg \gcd(f_i, X^n - 1)$. Direct sums prove the fixed formula; no square-free hypothesis on $X^n - 1$ is used. Primary decomposition separates the nilpotent X -part, of height e_i , from the invertible g_i -part. This proves the periodic-subspace and tail assertions. Möbius inversion gives P_n , orbit division gives C_n , and exponentiating $\sum_{n \geq 1} F_n z^n / n$ gives the zeta product. Each functional-graph component consists of a directed cycle and incoming trees. Ordering the cycle before tree vertices makes the composition matrix block triangular. A d -cycle contributes $T^d - 1$ and every noncycle vertex contributes T , proving the last formula. $\square$

2 Transient and inseparable structure

The theorem distinguishes two phenomena sometimes obscured by fixed-count tables. A factor X^e contributes no nonzero periodic coordinate but fixes the exact maximal tail e . Conversely, repeated factors coprime to X can produce nonsemisimple periodic dynamics. For example, $(X + 1)^4$ over $\mathbb{F}_2$ is inseparable and has restriction order four, while $(X + \alpha)^2$ over $\mathbb{F}_4$, with $\alpha^2 + \alpha + 1 = 0$, has order six. The latter uses field arithmetic, not arithmetic in $\mathbb{Z}/4\mathbb{Z}$.

Lemma 1. *The theorem remains valid when the characteristic divides n and $X^n - 1$ is inseparable.*

Proof. The cyclic-module kernel calculation uses the ideal generated by $\gcd(f_i, X^n - 1)$ with its full multiplicity. It never replaces the gcd by a set of distinct roots. $\square$

3 Exact audit

Eight controls range from 8 to 625 states. All 144 formula cells agree with exhaustive graphs and independent nullities of $A^n - I$. SymPy checks 198 polynomial gcd cells and six full-function characteristic polynomials. The largest transient control has 625 states, five periodic states, tail height three, and therefore Koopman zero multiplicity 620. A byte replay is exact. Seventeen repaired-hash semantic attacks and one separate stale-hash attack are rejected, including substitution of $\mathbb{Z}/4\mathbb{Z}$ for $\mathbb{F}_4$.

control	$ V $	$ \text{Per} $	tail	0-mult.
$\mathbb{F}_2$, nilpotent index 3	8	1	3	7
$\mathbb{F}_2$, nilpotent $\times$ order 3	16	4	2	12
$\mathbb{F}_5$, nilpotent $\times$ order 4	625	5	3	620
$\mathbb{F}_4$, nonsemisimple order 6	16	16	0	0

4 Scope and Route A

These are complete finite-field dynamical invariants, not target local Euler factors. No root number, automorphy statement, target cohomological determinant, or Hilbert–Pólya operator is claimed. The registered verdict is

(A0_FAIL, A1_WEAK, A2_FAIL, A3_FAIL, A4_FORMAL_HINT).

The exact record is `overall=ROUTE_A_REJECTED` and `route_b_invocation_allowed=false`.

Source note

Historical finite-linear context appears in Elspas [1] and Xu–Zou [2]. We make no priority claim.

References

- [1] B. Elspas, The theory of autonomous linear sequential networks, *IRE Trans. Circuit Theory* 6 (1959), 45–60, doi:10.1109/TCT.1959.1086506.
- [2] G. Xu and Y. M. Zou, Linear dynamical systems over finite rings, *J. Algebra* 321 (2009), 2149–2155, doi:10.1016/j.jalgebra.2008.09.029.

Declarations

Scope. The exact registered literal is `NO_BAD_EULER_OR_ROOT_NUMBER`. **Data and code.** All ledgers, independent checks, and build instructions are supplied in the HCS-C204 release package. **Competing interests.** None declared. **AI-use disclosure.** Generative AI tools assisted drafting and code generation. Every theorem statement, source record, exact output, and scope claim was checked through the released internal artifact audit chain; this is not external peer review or an independent error process.