

A Rank-Three Haar Phase Law for a Four-Symbol Open Walsh Gate

Route-A structural certificate C168

Abstract

We study the natural four-symbol open Walsh gate $A = F_4^* \text{diag}(1, 0, 1, 1)$ at the complete register clock $C_k = A^{\otimes k}$. Its characteristic polynomial is $x(x-1)(x^2 + ix/2 - 1/2)$, so the nonzero secular degree is exactly 3^k and the generalized zero space has dimension $4^k - 3^k$ for every register length. The two damped eigenvalues have normalized phases $u_{\pm}$ whose ratio r satisfies $r + r^{-1} = -3/2$. Since this nonintegral rational cannot be the algebraic-integer trace of a root of unity, r is non-torsion. The multiplicity-weighted phase measure is a three-step multiplicative walk, and its m th Fourier coefficient is $((1 + u_+^m + u_-^m)/3)^k$. Every fixed nonzero mode contracts strictly; hence the phase measures converge weakly to Haar measure on the circle. The result is source-side and does not construct a self-adjoint or target-spectrum operator.

Keywords: open quantum maps; Walsh dynamics; tensor spectra; phase equidistribution; Haar measure; algebraic integers.

中文摘要

本文研究自然四符号开放 Walsh 门 $A = F_4^* \text{diag}(1, 0, 1, 1)$ 在完整寄存器时钟 $C_k = A^{\otimes k}$ 下的精确谱结构。单点特征多项式为 $x(x-1)(x^2 + ix/2 - 1/2)$ ，因此对任意寄存器长度，非零世俗多项式次数恰为 3^k ，广义零特征空间维数为 $4^k - 3^k$ 。两个衰减特征值的单位相位记为 $u_{\pm}$ ，其比值 r 满足 $r + r^{-1} = -3/2$ 。若 r 是单位根，则该迹应为代数整数；但非整数有理数 $-3/2$ 不可能如此，故相位比非扭结。按代数重数加权后，相位测度成为三步乘法游走，其第 m 个傅里叶系数精确等于 $((1 + u_+^m + u_-^m)/3)^k$ 。每个固定非零模态均严格收缩，从而相位测度弱收敛到圆周上的 Haar 测度。结论只属于源侧开放动力学，不构造自伴算子或目标谱。

关键词：开放量子映射；Walsh 动力学；张量谱；相位等分布；Haar 测度；代数整数。

1 Frozen gate and one-site certificate

Let $F_4 = (i^{j\ell}/2)_{0 \leq j, \ell < 4}$, $P_1 = \text{diag}(1, 0, 1, 1)$, and $A = F_4^* P_1$. One application of the cyclic register map B_k is one tick; one complete cycle of exactly k ticks is $C_k = B_k^k = A^{\otimes k}$.

Lemma 1. A is diagonalizable with eigenvalues

$$0, \quad 1, \quad \lambda_+ = \frac{\sqrt{7} - i}{4}, \quad \lambda_- = \frac{-\sqrt{7} - i}{4}. \quad (1)$$

For $u_{\pm} = \sqrt{2}\lambda_{\pm}$, the ratio $r = u_+/u_- = (-3 + i\sqrt{7})/4$ is not a root of unity.

Proof. Direct determinant expansion gives

$$\chi_A(x) = x(x-1)(x^2 + ix/2 - 1/2), \quad (2)$$

with four distinct roots (1), and $|\lambda_{\pm}| = 1/\sqrt{2}$. Exact radical arithmetic gives $r + r^{-1} = -3/2$. Were r a root of unity, this sum would be an algebraic integer. A rational algebraic integer is an integer, a contradiction. $\square$

2 All-length secular and phase laws

For $a + b + c = k$, write $M_k(a, b, c) = k!/(a!b!c!)$.

Theorem 2. For every $k \geq 1$,

$$\det(I - zC_k) = \prod_{a+b+c=k} (1 - z\lambda_+^a \lambda_-^c)^{M_k(a, b, c)}. \quad (3)$$

The polynomial in (3) has degree 3^k , and the generalized zero eigenspace has dimension $4^k - 3^k$.

Proof. Tensor products of the four one-site eigenvectors form an eigenbasis. Nonzero products use the three nonzero roots; grouping words by their three contents gives (3) and the multinomial total 3^k . Every other tensor vector has a zero factor. $\square$

Let μ_k be the algebraic-multiplicity-weighted phase law of the nonzero spectrum. The same word count yields

$$\mu_k = 3^{-k} \sum_{a+b+c=k} M_k(a, b, c) \delta_{u_+^a u_-^c},$$

$$\hat{\mu}_k(m) = \left(\frac{1 + u_+^m + u_-^m}{3} \right)^k. \quad (4)$$

Theorem 3. μ_k converges weakly to normalized Haar measure on $\mathbb{T}$.

Proof. For $m \neq 0$, equality in the triangle inequality for the average in (4) would force $u_+^m = u_-^m = 1$, hence $r^m = 1$, contradicting Lemma 1. Every fixed nonzero Fourier mode therefore tends to zero. Density of trigonometric polynomials proves the claim. $\square$

The labels in (3) retain their multiplicities; distinctness across all triples is not asserted. Nor is a Fourier gap uniform in all modes claimed. At every finite k , μ_k is atomic, so its total-variation distance from continuous Haar measure is exactly one.

3 Evidence and boundary

The release freezes exact ledgers through $k = 24$ and modes $m = 24$. Independent strict Python and SymPy reconstructions, byte replay, and hostile semantic mutations test the implementation; the finite ledgers do not prove the all- k statements.

The Route-A tuple is (A1_WEAK, A2_FAIL, A3_FAIL, A4_UNITARY_OR_SCATTERING_CANDIDATE). No target divisor, functional equation, counting law, prime correspondence, arithmetic local data, Euler factor, root number, automorphy, Hilbert–Polya operator, or Route-B authorization is claimed. The scope is NO_BAD_EULER_OR_ROOT_NUMBER.

Declarations. All package data, exact code, and validators are included. No human or animal subjects, private data, or external dataset are involved. The anonymous contribution record covers conceptualization, formal analysis, software, validation, and writing. No funding or competing interest is declared. AI assistance was used for drafting and code generation; all claim-bearing formulas were deterministically reconstructed, and no external reviewer or acceptance score is represented.