

The Hemispherical Higgs Oscillator: Exact Actions, Friedrichs Spectrum, and Identity Revivals

HCS-C373 theorem package

4 September 2026

Abstract

We close the classical and Friedrichs quantum mechanics of the isotropic Higgs oscillator on one open hemisphere, while keeping its singular boundary explicit. For positive classical coupling, a turning-point integral gives the exact radial action $I_r = (J - |L| - \omega R^2)/2$, where $J = (2R^2 E + \omega^2 R^4)^{1/2}$. The Hamiltonian in actions has an exact 2:1 frequency lock, so every regular trajectory closes; circular, meridional, equilibrium, and zero-coupling faces are separated. For the quantum Friedrichs realization with nonnegative coupling, Jacobi modes have $E_N = \hbar^2(N+1)(N+1+2\nu)/(2R^2)$ and multiplicity $N+1$. Consecutive spectral gaps then prove that the propagator is the identity precisely when $\tau = \pi M$ and $M(3+2\nu)$ is even. Thus a positive identity revival exists if and only if 2ν is rational, with an explicit least time. Exact finite enumeration is supplied only as a reproducibility receipt.

1 Positive-coupling classical system

Fix $R > 0, \omega > 0$, and polar coordinates $0 \leq \theta < \pi/2, \phi \in \mathbb{R}/(2\pi\mathbb{Z})$. With $L = p_\phi$, consider

$$H = \frac{p_\theta^2}{2R^2} + \frac{L^2}{2R^2 \sin^2 \theta} + \frac{\omega^2 R^2}{2} \tan^2 \theta. \quad (1)$$

The positive coupling is part of the theorem: its divergent equatorial barrier confines every finite-energy trajectory away from the missing equator. Put

$$J = (2R^2 E + \omega^2 R^4)^{1/2}, \quad x = \sin^2 \theta.$$

Theorem 1 (Turning points, action, and inversion). *For a regular trajectory with $L \neq 0$, the two turning points $0 < x_- < x_+ < 1$ are the roots of*

$$J^2 x^2 - (2R^2 E + L^2)x + L^2 = 0. \quad (2)$$

The radial action and the Hamiltonian in actions are

$$I_r = \frac{1}{2\pi} \oint p_\theta d\theta = \frac{J - |L| - \omega R^2}{2}, \quad (3)$$

$$H(I_r, L) = \frac{(2I_r + |L| + \omega R^2)^2 - \omega^2 R^4}{2R^2}. \quad (4)$$

The allowed chamber is $E \geq \omega|L| + L^2/(2R^2)$, with equality exactly on the circular face.

Proof. At energy E , direct rearrangement of (1) gives

$$p_\theta^2 = 2R^2 E - \frac{L^2}{x} - \frac{\omega^2 R^4 x}{1-x} = \frac{J^2(x-x_-)(x_+-x)}{x(1-x)}.$$

Because $d\theta = dx/(2\sqrt{x(1-x)})$, the two branches of the closed radial loop give

$$I_r = \frac{J}{2\pi} \int_{x_-}^{x_+} \frac{\sqrt{(x-x_-)(x_+-x)}}{x(1-x)} dx. \quad (5)$$

For $0 < a < b < 1$, substitute $x = a + (b-a)\sin^2 u$ and split $1/[x(1-x)] = 1/x + 1/(1-x)$. The two elementary integrals yield

$$\int_a^b \frac{\sqrt{(x-a)(b-x)}}{x(1-x)} dx = \pi \{1 - \sqrt{ab} - \sqrt{(1-a)(1-b)}\}. \quad (6)$$

Vieta's relations for (2) are

$$x_- x_+ = \frac{L^2}{J^2}, \quad (1 - x_-)(1 - x_+) = \frac{\omega^2 R^4}{J^2}.$$

Equations (5) and (6) prove (3); solving for J proves (4). The two roots coalesce exactly at the displayed energy threshold. $\square$

Theorem 2 (Frequency lock and periods). *On either signed chamber $L > 0$ or $L < 0$,*

$$\Omega_r = \frac{\partial H}{\partial I_r} = \frac{2J}{R^2}, \quad \Omega_\phi = \frac{\partial H}{\partial L} = \text{sgn}(L) \frac{J}{R^2}.$$

Every regular trajectory has radial period $\pi R^2/J$ and primitive phase-space period $2\pi R^2/J$.

Proof. Differentiate (4). During one radial return the azimuthal angle advances by $\pi \text{sgn}(L)$, so it does not return to the same phase-space point. Two radial returns give angular increments 4π and $2\pi \text{sgn}(L)$ in the two action angles, proving primitivity. $\square$

2 Classical boundary atlas

The action formula has four distinct boundary interpretations.

face	exact behavior
circular	$I_r = 0, L \neq 0$; the double root is $\sin^2 \theta_c = L /(L + \omega R^2)$ and the angular period is $2\pi R^2/J$.
meridional	$L = 0, I_r > 0$; the roots are 0 and $1 - (\omega R^2/J)^2$. A Cartesian tangent chart continues the orbit through the polar-coordinate singularity.
north equilibrium	$I_r = L = 0$, hence $E = 0$ and the orbit is fixed.
zero coupling	$\omega = 0$ removes the equatorial barrier. The open hemisphere is then incomplete for crossing geodesics, so it is excluded from the classical periodic-flow theorem.

This separation prevents a formula valid on regular Liouville tori from being silently extended through collapsed or incomplete faces.

References

- [1] P. W. Higgs, "Dynamical symmetries in a spherical geometry. I," *J. Phys. A* 12 (1979), doi:10.1088/0305-4470/12/3/006.
- [2] H. I. Leemon, "Dynamical symmetries in a spherical geometry. II," *J. Phys. A* 12 (1979), doi:10.1088/0305-4470/12/4/009.
- [3] S. Bellucci, A. Nersessian, A. Saghatelian, and V. Yeghikyan, "Quantum ring models and action-angle variables," arXiv:1008.3865.
- [4] Ye. M. Hakobyan and G. S. Pogosyan, "On Interbasis Expansion for Isotropic Oscillator on Two-Dimensional Sphere," arXiv:quant-ph/9803085.