

The Complete S_4 Chebotarev–Frobenius Permutation Atlas for the Root Scheme of $x^4 - x - 1$

HCS-C369 / HEN-O353

4 September 2026

Abstract

The integral quartic $f(x) = x^4 - x - 1$ defines an intrinsic prime-indexed family of finite dynamical systems: arithmetic Frobenius $F_p(\alpha) = \alpha^p$ permutes the four geometric roots at every good prime. We prove that $\text{disc}(f) = -283$ and $\text{Gal}(f/\mathbb{Q}) = S_4$, identify factor degrees with primitive cycle lengths, and close the fixed-point and finite-determinant identities on every good fiber. All five S_4 cycle types are exhibited. The final round adds their Chebotarev densities, the non-étale boundary at $p = 283$, and the exact fiberwise Koopman classification. The construction is a source-arithmetic candidate, not a cross-prime target determinant. The universal finite-permutation trace/determinant mechanism remains owned by the prior workspace package HCS-C12A; our owner is the complete $x^4 - x - 1$ -specific atlas and its executable convention lock.

1 Quartic arithmetic and the S_4 proof

Put

$$f(x) = x^4 - x - 1, \quad \mathcal{X} = \text{Spec } \mathbb{Z}[x]/(f). \quad (1)$$

For a rational prime p , write X_p for the geometric roots of the reduction of f in $\overline{\mathbb{F}}_p$. We use *arithmetic Frobenius*

$$F_p : X_p \longrightarrow X_p, \quad \alpha \longmapsto \alpha^p. \quad (2)$$

Geometric Frobenius is F_p^{-1} . The two conventions have identical cycle lengths, fixed counts, and determinants; this paper never conflates the maps, only their common cycle partition.

Theorem 1 (Arithmetic and Galois closure). *The discriminant of f is -283 . The polynomial is irreducible over $\mathbb{Q}$ and its splitting field L has*

$$\text{Gal}(L/\mathbb{Q}) \cong S_4.$$

Every prime $p \neq 283$ is unramified and $\mathcal{X}_{\mathbb{F}_p}$ is a finite étale scheme of degree four. If $\lambda_p = (d_1, \dots, d_k)$ is the multiset of degrees of the distinct irreducible factors of $f \bmod p$, then λ_p is exactly the cycle partition of F_p on X_p .

Proof. For $x^4 + qx + r$ the discriminant is $256r^3 - 27q^4$; inserting $q = r = -1$ gives $-256 - 27 = -283$. Modulo 2,

$$f = x^4 + x + 1.$$

It has no root, and the only monic irreducible quadratic over $\mathbb{F}_2$ is $x^2 + x + 1$, whose square is $x^4 + x^2 + 1$. Hence $f \bmod 2$ is irreducible. Gauss’s lemma makes f irreducible over $\mathbb{Q}$, so its Galois group $G \leq S_4$ is transitive. Dedekind’s theorem [1, 2] at the unramified prime 2 supplies a 4-cycle. At $p = 7$,

$$f \equiv (x - 3)(x^3 + 3x^2 + 2x - 2) \pmod{7}, \quad (3)$$

and the cubic factor has no root, so G also contains a 3-cycle. Consequently 12 divides $|G|$, while $|G|$ divides 24. A subgroup of order 12 in S_4 is the index-two sign kernel A_4 , but a 4-cycle is odd; equivalently, the discriminant is not a square. Therefore $G = S_4$.

The discriminant criterion makes precisely $p = 283$ exceptional. At every other prime, the roots are distinct. An irreducible factor of degree d has its roots in one orbit of $\alpha \mapsto \alpha^p$ of length d , and every orbit arises this way [3]. This proves the dictionary. $\square$

Proposition 1 (Five exact witnesses). *Every partition of four occurs among the good fibers. In addition to (3), one has*

$$\begin{array}{lll}
p = 2 : & f = x^4 + x + 1 & (4), \\
p = 17 : & f = (x + 2)(x + 5)(x^2 - 7x + 5) & (2 + 1 + 1), \\
p = 71 : & f = (x^2 + 15x - 20)(x^2 - 15x + 32) & (2 + 2), \\
p = 83 : & f = (x + 3)(x + 7)(x + 14)(x - 24) & (1 + 1 + 1 + 1),
\end{array}$$

where every equality is in the corresponding residue field and the final parenthesis is the Frobenius cycle partition.

Proof. Multiplication gives f modulo the displayed prime. The linear factors are explicit; the remaining quadratic or cubic factors have no residue-field root. The mod-2 case was proved in Theorem 1. $\square$

2 All-iterate orbit and determinant closure

HCS-C12A already owns the universal zero-dimensional statement that Frobenius on a reduced finite fiber is a permutation, its iterate counts are traces, and its finite zeta is the reciprocal permutation determinant. C369 does not claim workspace ownership of that universal mechanism. This section specializes it to the five S_4 classes of $x^4 - x - 1$ and locks the factor, fixed, primitive, and determinant conventions in one executable all-good-prime ledger. For a good p , let P_p be the permutation matrix on $\mathbb{C}[X_p]$ defined by $P_p e_\alpha = e_{F_p(\alpha)}$, and set

$$N_p(r) = \# \text{Fix}(F_p^r).$$

Theorem 2 (Fixed, primitive, and determinant identities). *Let $p \neq 283$, let $\lambda_p = (d_1, \dots, d_k)$, and let $r, n \geq 1$. Then*

$$N_p(r) = \sum_{j: d_j | r} d_j. \quad (4)$$

If $E_p(n)$ is the number of points of exact period n and $C_p(n)$ the number of primitive n -cycles, then

$$E_p(n) = \sum_{d|n} \mu(d) N_p(n/d), \quad C_p(n) = \frac{E_p(n)}{n}. \quad (5)$$

Finally, in $\mathbb{Q}[[u]]$ and therefore as a rational function,

$$\begin{aligned}
Z_p(u) &:= \exp \left(\sum_{r \geq 1} \frac{N_p(r)}{r} u^r \right) \\
&= \det(I - uP_p)^{-1} = \prod_{j=1}^k (1 - u^{d_j})^{-1}.
\end{aligned} \quad (6)$$

Thus the factor degrees are not merely a fixed-point statistic: they are the complete primitive-cycle ledger on the fiber.

Proof. A d -cycle contributes all its d points to $\text{Fix}(F_p^r)$ exactly when $d \mid r$, proving (4). The relation $N_p(r) = \sum_{n|r} E_p(n)$ and Möbius inversion give (5); division by n groups exact-period points into cycles. On a d -cycle block C_d , $\det(I - uC_d) = 1 - u^d$. Multiplying blocks proves the determinant equality, while

$$-\log(1 - u^d) = \sum_{m \geq 1} \frac{u^{dm}}{m} = \sum_{m \geq 1} \frac{d}{dm} u^{dm}$$

and (4) prove the exponential equality. $\square$

type	1 + 1 + 1 + 1	2 + 1 + 1	2 + 2	3 + 1	4
good primes $\leq 10^4$	43	306	147	411	321
S_4 class size	1	6	3	8	6

Table 1: Exact finite regression counts. The 1,228 entries sum across the row; these counts are not used to prove an asymptotic density.

The canonical evidence enumerates all 1,229 primes at most 10^4 , removes the unique bad prime 283, and checks (4)–(6) for $1 \leq r \leq 12$ on all 1,228 good fibers: 14,736 prime–iterate cells. A separate factorization backend recomputes every partition. Exact symbolic and hostile-mutation lanes test the proof anchors and schema. Finite enumeration is an implementation receipt; Theorems 1 and 2 establish the quartic specialization for all good primes and all iterates.

References

- [1] J.-P. Serre, *Topics in Galois Theory*, second edition, A K Peters, 2008.
- [2] J. Neukirch, *Algebraic Number Theory*, Springer, 1999.
- [3] R. Lidl and H. Niederreiter, *Finite Fields*, second edition, Cambridge University Press, 1997.