

The Five Lagrange Equilibria in the Planar CR3BP: Exact Linear Types and the Defective Routh Boundary

Route-A source-local certificate HCS-C290

2 September 2026

Abstract

For the normalized planar circular restricted three-body problem with $0 < \mu \leq 1/2$, we derive all five equilibria and their complete linear spectral atlas. Each collinear point is uniquely located in its collision-free interval and is saddle-times-center. At the equilateral points the characteristic polynomial is $\lambda^4 + \lambda^2 + 27\mu(1 - \mu)/4$. Writing $\mu_R = (1 - \sqrt{23/27})/2$, the linear flow is bounded elliptic below μ_R and has an unstable Hamiltonian quartet above it. At equality each of $\pm i/\sqrt{2}$ has algebraic multiplicity two and geometric multiplicity one: the linearization is defective, has linearly growing solutions, and is *not* stable.

1 Frozen normalization and claim level

The primaries of masses $1 - \mu$ and μ occupy $(-\mu, 0)$ and $(1 - \mu, 0)$ in the synodic frame. For

$$r_1^2 = (x + \mu)^2 + y^2, \quad r_2^2 = (x - 1 + \mu)^2 + y^2,$$

set

$$\Omega(x, y) = \frac{x^2 + y^2}{2} + \frac{1 - \mu}{r_1} + \frac{\mu}{r_2}. \quad (1)$$

The equations, including our Coriolis convention, are

$$\ddot{x} - 2\dot{y} = \Omega_x, \quad \ddot{y} + 2\dot{x} = \Omega_y. \quad (2)$$

Both collision points are removed. “Linearly stable” below means that all solutions of the constant-coefficient variational equation are bounded, including at mass ratios where the two linear frequencies are resonant. It does not assert nonlinear stability, control resonance bifurcations, or invoke KAM theory.

The equilateral configuration goes back to Lagrange [1]. Gascheau’s 1843 thesis is the first historical owner of the Newtonian triangular stability criterion [2]; Routh’s 1874 paper gives a subsequent treatment and inverse-power-law generalization [3]. A modern Hamiltonian treatment is given by Meyer, Hall, and Offin [4]. Neither the five points nor the threshold is claimed as literature originality; all formulas needed for our convention are rederived here.

Theorem 1 (Equilibrium and linear-type atlas). *For every $0 < \mu \leq 1/2$, the collision-free plane has exactly five equilibria.*

1. *There is exactly one collinear equilibrium in each interval $(-\infty, -\mu)$, $(-\mu, 1 - \mu)$, and $(1 - \mu, \infty)$. Every one is of saddle-times-center type and is linearly unstable.*
2. *The two remaining equilibria are*

$$L_{4,5} = \left(\frac{1}{2} - \mu, \pm \frac{\sqrt{3}}{2} \right).$$

Their characteristic polynomial is

$$p_\mu(\lambda) = \lambda^4 + \lambda^2 + \frac{27}{4}\mu(1 - \mu). \quad (3)$$

With $\mu_R = (1 - \sqrt{23/27})/2$, their linearized flow is bounded for $0 < \mu < \mu_R$, defective with linear growth (hence not stable) for $\mu = \mu_R$, and exponentially unstable for $\mu_R < \mu \leq 1/2$.

At equality, each eigenvalue $\pm i/\sqrt{2}$ has algebraic multiplicity two and geometric multiplicity one.

2 Why there are exactly five points

Define

$$S(x, y) = \frac{1 - \mu}{r_1^3} + \frac{\mu}{r_2^3}. \quad (4)$$

Direct differentiation gives $\Omega_y = y(1 - S)$. Thus an equilibrium either has $y = 0$ or satisfies $S = 1$.

On the axis, put

$$f_\mu(x) = x - (1 - \mu) \frac{x + \mu}{|x + \mu|^3} - \mu \frac{x - 1 + \mu}{|x - 1 + \mu|^3}. \quad (5)$$

Away from collisions,

$$f'_\mu(x) = 1 + 2 \left(\frac{1 - \mu}{|x + \mu|^3} + \frac{\mu}{|x - 1 + \mu|^3} \right) > 0. \quad (6)$$

On $(-\infty, -\mu)$ its limiting signs are negative and positive; on $(-\mu, 1 - \mu)$ they are negative and positive; and on $(1 - \mu, \infty)$ they are negative and positive. The intermediate value theorem and (6) give precisely three roots.

If $y \neq 0$, then $S = 1$. Writing $u = r_1^{-3}$ and $w = r_2^{-3}$, a direct rearrangement yields

$$\Omega_x - x(1 - S) = \mu(1 - \mu)(w - u). \quad (7)$$

Because $0 < \mu < 1$, equilibrium implies $r_1 = r_2$, hence $x = 1/2 - \mu$. Now $S = 1$ becomes $r_1 = r_2 = 1$, so $y^2 = 3/4$. These are exactly L_4 and L_5 , proving the count.

3 The raw rotating-frame determinant

At any equilibrium let $a = \Omega_{xx}$, $c = \Omega_{yy}$, and $d = \Omega_{xy}$. In coordinates $(\xi, \eta, u, w) = (\delta x, \delta y, \delta \dot{x}, \delta \dot{y})$ the raw variational matrix is

$$M = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ a & d & 0 & 2 \\ d & c & -2 & 0 \end{pmatrix}. \quad (8)$$

Expanding, without appealing to a stability label,

$$\det(\lambda I - M) = \lambda^4 + (4 - a - c)\lambda^2 + ac - d^2. \quad (9)$$

At a collinear root, $d = 0$ and

$$a = 1 + 2S, \quad c = 1 - S, \quad (10)$$

where now S is (4) at $y = 0$. We verify $S > 1$ on all three branches. Between the primaries, $r_1 + r_2 = 1$, so $r_1, r_2 < 1$ and $S > (1 - \mu) + \mu = 1$. To the right, write $d = x - 1 + \mu > 0$. At $d = \mu^{1/3}$, equation (5) becomes

$$f_\mu = (1 - \mu)\{1 - (1 + d)^{-2}\} > 0.$$

Since f_μ is increasing in x and tends to $-\infty$ at the right side of the collision, its root has $d < \mu^{1/3}$; hence $S > \mu/d^3 > 1$. To the left, put $d = -x - \mu > 0$. At $d = (1 - \mu)^{1/3}$ one obtains

$$f_\mu = -\mu\{1 - (1 + d)^{-2}\} < 0.$$

Here increasing d decreases x , so the root has $d < (1 - \mu)^{1/3}$ and $S > (1 - \mu)/d^3 > 1$. Substitution in (9) gives

$$\lambda^4 + (2 - S)\lambda^2 + (1 + 2S)(1 - S). \quad (11)$$

Its constant term is negative. The two roots in $z = \lambda^2$ have opposite signs, so there is one real pair and one imaginary pair. This proves the collinear assertion.

At either triangular point direct differentiation of (1) gives

$$a = \frac{3}{4}, \quad c = \frac{9}{4}, \quad d = \pm \frac{3\sqrt{3}}{4}(1 - 2\mu). \quad (12)$$

Thus $a + c = 3$ and $ac - d^2 = 27\mu(1 - \mu)/4$, proving (3) for both L_4 and L_5 .

4 Open chambers and the critical Jordan structure

Set $z = \lambda^2$. The two roots of (3) are

$$z_{\pm} = \frac{-1 \pm \sqrt{1 - 27\mu(1 - \mu)}}{2}. \quad (13)$$

On $(0, 1/2]$, the expression $27\mu(1 - \mu)$ is strictly increasing until the included endpoint and equals one precisely at μ_R . Below μ_R , both $z_{\pm}$ are distinct and negative, giving four distinct imaginary eigenvalues and a bounded diagonalizable linear flow. Above it, the z roots are nonreal conjugates; their square roots form a quartet $\{\pm\alpha \pm i\beta\}$ with $\alpha, \beta > 0$, giving exponential growth.

At equality,

$$p_{\mu_R}(\lambda) = (\lambda^2 + \tfrac{1}{2})^2. \quad (14)$$

Algebraic multiplicity is not enough to decide boundedness. For an eigenvalue λ , the position components solve

$$\begin{pmatrix} \lambda^2 - a & -2\lambda - d \\ 2\lambda - d & \lambda^2 - c \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix} = 0. \quad (15)$$

At $\lambda = \pm i/\sqrt{2}$ the determinant vanishes. The matrix cannot be zero: its two off-diagonal entries would imply simultaneously $2\lambda = -d$ and $2\lambda = d$, hence $\lambda = 0$. Therefore it has rank one and a one-dimensional kernel. Velocities are then uniquely $(u, w) = \lambda(\xi, \eta)$, so each eigenspace of M has dimension one. Each double eigenvalue consequently has a nontrivial size-two Jordan block. The exponential contains $te^{\pm it/\sqrt{2}}$ terms and is unbounded. This proves the critical statement and, in particular, excludes linear stability.

References

- [1] J.-L. Lagrange, *Essai sur le Problème des trois Corps*, Prix de l'Académie royale des sciences de Paris, tome IX (1772); Oeuvres, tome VI, 229–331. [digitized transcription](#).
- [2] G. Gascheau, *Mouvements relatifs d'un système de corps*, thèse de mécanique, Faculté des sciences de Paris, Bachelier, 1843, 36 pp. and plate. [BnF Gallica ark:/12148/bpt6k5789653w](#).
- [3] E. J. Routh, *On Laplace's Three Particles, with a Supplement on the Stability of Steady Motion*, Proc. London Math. Soc. s1-6 (1874), 86–97. [doi:10.1112/plms/s1-6.1.86](#).
- [4] K. R. Meyer, G. R. Hall, and D. Offin, *Introduction to Hamiltonian Dynamical Systems and the N-Body Problem*, 2nd ed., Springer, 2009. [doi:10.1007/978-0-387-09724-4](#).