

RECORDED-TRANSVERSAL LAWS ON RATE-WEIGHTED SUNFLOWER FORESTS

ANONYMOUS

ABSTRACT. Consider the owned random-edge/random-vertex covering process, restricted to vertex-disjoint heterogeneous sunflowers with fixed positive edge rates. We derive the complete law of the full recorded transversal, without reducing it to a minimal transversal. For one sunflower, an inclusion–exclusion integral gives every selected-petal mask at arbitrary rates, and independent uniform marks refine that law to every actual vertex set. At unit rates, elementary symmetric polynomials give the complete stopping probability generating function, mean, and variance; the maximal-time atom is resolved into its all-petal and final-core events. For a disjoint sunflower forest, the marked stopped laws factor and the stopping clocks convolve. The covering algorithm, sunflower carrier, exponential/size-biased ordering, and independence of disjoint restrictions are prior or generic machinery and are not claimed as contributions. A finite exact-arithmetic program checks the formulas on a frozen bounded family.

1. POSITIONING AND CONTRIBUTION

Sunflowers, also called Delta-systems, are classical [Erdős and Rado, 1960]. Their transversals belong to a broad literature that includes extremal and approximation questions [Chvátal and McDiarmid, 1992]. More importantly for the present dynamics, the random covering algorithm is not new. Pitt gave its graph ancestor [Pitt, 1985], and Bar-Yehuda [2000, Section 5.1 and Theorem 6] directly treats the Pitt rule for hypergraph vertex cover. In the unweighted setting, its vertex choice in the selected hyperedge is uniform. We therefore assign zero credit to the covering algorithm, its validity, and every approximation guarantee.

The ordering device is also owned background. Rate-proportional sampling without replacement is a Plackett–Luce/size-biased order [Plackett, 1975]; positive-rate independent exponentials and dissociation of restrictions to disjoint item sets are treated explicitly by Gnedin [2026]. We assign zero credit to the exponential representation, memorylessness, finite ranking probabilities, and generic forest independence. The same applies to inclusion–exclusion, beta integrals, elementary symmetric polynomials, tail-sum identities, and products of PGFs.

After this subtraction, the contribution is deliberately narrow: for the owned covering process restricted to vertex-disjoint heterogeneous sunflowers with fixed edge rates, we derive the complete recorded-transversal law and, at unit rates, the complete stopping PGF and first two moments. The point is the closed conjunction of aggregate endpoint, actual-vertex refinement, stopping law, and marked forest factorization for this carrier. We claim neither a new algorithm nor a general theory of random greedy hitting sets.

2. MODEL AND MARKED ORDER

Fix positive integers

$$m, c, p_1, \dots, p_m$$

and positive real numbers $\lambda_1, \dots, \lambda_m$. A sunflower has a core C of size c , pairwise disjoint petals P_i of size p_i , and edges

$$E_i = C \cup P_i, \quad i \in [m].$$

At each step, an active edge E_i is chosen with probability proportional to its fixed rate λ_i . A vertex is then chosen uniformly from E_i and added to the recorded set. A petal choice deletes only E_i ; a core choice hits all remaining edges and stops the component. If every edge receives a petal choice, the process also stops. The endpoint is the full recorded set: previously selected petal vertices are retained, so the endpoint need not be a minimal transversal.

Write

$$r_i = \frac{p_i}{c + p_i}, \quad q_i = \frac{c}{c + p_i} = 1 - r_i, \quad \Lambda(A) = \sum_{i \in A} \lambda_i.$$

Let $X_i \sim \text{Exp}(\lambda_i)$ independently. Independently of all clocks and one another, give edge i a uniform mark $U_i \in E_i$. Sorting the X_i and reading marks until the first core mark gives the original process. Indeed, the minimum clock has the required rate-proportional law, and after a petal deletion the residual clocks have the same law by memorylessness. The fixed-rate and independent-mark hypotheses are essential for this coupling. This representation is used only as owned machinery.

3. THE COMPLETE WEIGHTED ENDPOINT LAW

For a proper subset $A \subsetneq [m]$, let $\pi(A)$ be the probability that exactly the edges in A are recorded through petal vertices before a core vertex stops the component. Define

$$(1) \quad I(A) = \sum_{B \subseteq A} \frac{(-1)^{|B|}}{\Lambda([m] \setminus A) + \Lambda(B)}.$$

Every denominator is positive because A is proper.

Theorem 3.1 (weighted aggregate endpoint). *For every proper $A \subsetneq [m]$,*

$$(2) \quad \pi(A) = \left(\prod_{i \in A} r_i \right) \left(\sum_{j \notin A} q_j \lambda_j \right) I(A).$$

The remaining, all-petal endpoint has probability

$$(3) \quad \pi([m]) = \prod_{i=1}^m r_i.$$

These 2^m masses are positive and sum to one.

Proof. Fix $A \subsetneq [m]$ and expose the time t of the stopping core mark. For each $i \in A$, edge i must carry a petal mark and satisfy $X_i < t$. For each edge outside A , no clock may ring before t , except that one such edge j rings at t and carries a core mark. Summing the latter density over $j \notin A$ gives

$$(4) \quad \pi(A) = \left(\prod_{i \in A} r_i \right) \left(\sum_{j \notin A} q_j \lambda_j \right) \int_0^\infty e^{-\Lambda([m] \setminus A)t} \prod_{i \in A} (1 - e^{-\lambda_i t}) dt.$$

Expanding the product in (4) and integrating each exponential gives (1) and (2). The only way to terminate without a core mark is for every mark to lie in its petal, proving (3). The marked-clock construction partitions its probability space into these events, which proves normalization; the integral form proves positivity. $\square$

The aggregate law also resolves every actual vertex set, including at unequal rates.

Corollary 3.2 (actual-vertex law). *Fix $A \subsetneq [m]$, vertices $x_i \in P_i$ for $i \in A$, and $y \in C$. Then*

$$(5) \quad \mathbb{P}(R = \{y\} \cup \{x_i : i \in A\}) = \frac{\pi(A)}{c \prod_{i \in A} p_i}.$$

For $x_i \in P_i$ for every i ,

$$(6) \quad \mathbb{P}(R = \{x_1, \dots, x_m\}) = \prod_{i=1}^m \frac{1}{c + p_i}.$$

Proof. Conditional on its category, every petal mark is uniform on its own P_i and the stopping core mark is uniform on C . These conditional marks are independent of the clocks and of one another. Dividing the aggregate mass by $c \prod_{i \in A} p_i$ proves (5). In the all-petal event, edge i selects a specified petal vertex with probability $1/(c + p_i)$; independence gives (6). $\square$

4. UNIT RATES: STOPPING PGF AND MOMENTS

Assume throughout this section that $\lambda_i = 1$. Let $T \in \{1, \dots, m\}$ be the number of recorded vertices at absorption, let $e_t(r_1, \dots, r_m)$ be the elementary symmetric polynomial, and put

$$(7) \quad s_t = \frac{e_t(r_1, \dots, r_m)}{\binom{m}{t}}, \quad 0 \leq t \leq m,$$

with $s_0 = 1$.

Theorem 4.1 (complete stopping law). *For $0 \leq t < m$,*

$$(8) \quad \mathbb{P}(T > t) = s_t.$$

Consequently,

$$(9) \quad \mathbb{P}(T = t) = s_{t-1} - s_t \quad (1 \leq t < m), \quad \mathbb{P}(T = m) = s_{m-1}.$$

The maximal-time atom is the disjoint union of two mechanisms:

$$(10) \quad \mathbb{P}(T = m) = \prod_{i=1}^m r_i + \frac{1}{m} \sum_{j=1}^m q_j \prod_{i \neq j} r_i = \frac{e_{m-1}(r_1, \dots, r_m)}{m}.$$

In particular, its first term is the all-petal endpoint, while its second term is a final core choice after $m-1$ petal choices.

Proof. At unit rates the clock order is a uniform random permutation of $[m]$, independent of the marks. The event $T > t$ says that the first t distinct edges in this order all have petal marks. Averaging $\prod_{i \in A} r_i$ over the $\binom{m}{t}$ possible first-edge sets gives (8), and differencing tails gives (9) below the top atom. For $T = m$, either all m marks are petals, or a unique edge j is last in the uniform order, its mark is core, and all other marks are petals. This gives the first equality in (10). Substituting $q_j = 1 - r_j$ reduces it to $m^{-1} \sum_j \prod_{i \neq j} r_i = e_{m-1}/m = s_{m-1}$. $\square$

Corollary 4.2 (PGF, mean, and variance). *The probability generating function and first two moments are*

$$(11) \quad G(z) = \sum_{t=1}^{m-1} (s_{t-1} - s_t) z^t + s_{m-1} z^m,$$

$$(12) \quad \mathbb{E}T = \sum_{t=0}^{m-1} s_t,$$

$$(13) \quad \mathbb{E}T^2 = \sum_{t=0}^{m-1} (2t+1) s_t,$$

$$(14) \quad \text{Var}(T) = \sum_{t=0}^{m-1} (2t+1) s_t - \left(\sum_{t=0}^{m-1} s_t \right)^2.$$

Proof. Equation (11) is the mass formula (9). For every positive integer-valued $T \leq m$, $T = \sum_{t=0}^{m-1} \mathbf{1}_{\{T > t\}}$ and $T^2 = \sum_{t=0}^{m-1} (2t+1) \mathbf{1}_{\{T > t\}}$. Taking expectations and using (8) proves (12)–(14). $\square$

The heterogeneity is visible in every e_t : replacing the r_i by their mean generally changes all terms of degree at least two and does not preserve the endpoint law.

5. DISJOINT SUNFLOWER FORESTS

Let $H_1, \dots, H_d$ be vertex-disjoint sunflower components, each with its own positive parameters and fixed edge rates. The global scheduler chooses among all active edges with probability proportional to their rates. Write R_a, T_a , and G_a for the recorded endpoint, number of choices, and PGF in component a ; write R and T for the forest quantities.

Theorem 5.1 (marked stopped factorization). *For arbitrary positive fixed rates,*

$$(15) \quad \mathcal{L}(R) = \bigotimes_{a=1}^d \mathcal{L}(R_a), \quad T = \sum_{a=1}^d T_a, \quad G_T(z) = \prod_{a=1}^d G_a(z).$$

Thus the endpoint mass of a forest endpoint is the product of the component masses from Theorem 3.1 and Corollary 3.2; at unit rates each G_a is given by (11).

Proof. Assign independent clocks and marks to every edge. Each local stopped marked history (R_a, T_a) is a function only of the clocks and marks in H_a ; hence these pairs are independent. Sorting all clock rings across components gives the global rate-proportional scheduler. Once a core mark stops one component, suppressing its later clocks does not change any local history. The forest records exactly the union of the local records and makes exactly the sum of their choices. Tensorization and multiplication of PGFs follow. $\square$

The theorem’s content is only the fully marked stopped consequence for this process. Independence/dissociation of disjoint exponential-order restrictions is generic and receives no novelty credit.

6. EXACT CONTROLS AND LIMITATIONS

The accompanying self-contained Python program uses only integers and `fractions.Fraction`. It performs no sampling or floating-point calculation. It recursively enumerates the process and compares every mass with the displayed formulas. The frozen scope is:

lane	parameter range	inputs
unit-rate aggregate	$c \leq 3, m \leq 5, p_i \leq 4$	4092
weighted aggregate	$c \leq 2, m \leq 3, p_i, \lambda_i \leq 3$	1638
unit-rate actual vertices	$c \leq 2, m \leq 3, p_i \leq 3$	78
two-component forests	three unit-rate and one weighted control	4

These are falsification controls, not proofs. In particular, the program does not exhaust weighted vertex-resolved endpoints or arbitrary forests; those claims follow from the all-parameter arguments above. The bounded owner search located the direct process and ordering owners already subtracted, but did not locate this complete conjunction. That non-hit is not a novelty certificate, and a direct package owner would supersede the present positioning.

7. CONCLUSION

On a heterogeneous sunflower, the marked size-biased order separates the recorded-transversal problem into a weighted endpoint integral, independent actual marks, and—at unit rates—an elementary-symmetric stopping law. On vertex-disjoint components, these marked stopped laws tensorize. The result is a complete exact atlas for a sharply restricted owned process, not a new covering algorithm, ordering method, or independence principle.

REFERENCES

- Reuven Bar-Yehuda. One for the price of two: A unified approach for approximating covering problems. *Algorithmica*, 27(2):131–144, 2000. doi: 10.1007/s004530010009.
- Vášek Chvátal and Colin McDiarmid. Small transversals in hypergraphs. *Combinatorica*, 12(1):19–26, 1992. doi: 10.1007/BF01191201.
- Paul Erdős and Richard Rado. Intersection theorems for systems of sets. *Journal of the London Mathematical Society*, s1-35(1):85–90, 1960. doi: 10.1112/jlms/s1-35.1.85.
- Alexander Gnedin. Infinite size-biased orders. *Order*, 43:29, 2026. doi: 10.1007/s11083-026-09743-2.
- Leonard Pitt. A simple probabilistic approximation algorithm for vertex cover. Technical Report YaleU/DCS/TR-404, Department of Computer Science, Yale University, 1985.
- Robin L. Plackett. The analysis of permutations. *Journal of the Royal Statistical Society. Series C (Applied Statistics)*, 24(2):193–202, 1975. doi: 10.2307/2346567.