

Supplementary Materials for

The Emergence of Prime Distribution from Low-Dimensional Deterministic Chaos

Liang Wang

Corresponding author: wangliang.f@gmail.com

Dynamization of Sieves: From Arithmetic to Symbolic Dynamics

To transform the number-theoretic problem into a physical framework, we first establish **Symbolic Dynamics** as the bridge connecting the arithmetic properties of primes with the orbital properties of dynamical systems. We reformulate the traditional Sieve of Eratosthenes—classically a static elimination algorithm—as a dynamic time-evolution process on the one-dimensional discrete space of natural numbers $\mathbb{N}$.

Let the state space be $\Sigma = \{L, R\}$, where the symbol L (Left) denotes a "surviving" state (potential prime) and the symbol R (Right) denotes a "sieved-out" state (composite). This notation is chosen to align with the phase space partition of the unimodal map relative to its critical point, as detailed in [3].

For the i -th prime p_i , we define its Sieving Operator as a periodic symbol sequence S_{p_i} . Formally defined as:

$$S_{p_i} = (R, \underbrace{L, L, \dots, L}_{p_i-1})$$

This implies that within a period of length p_i , the first position (corresponding to multiples of p_i) is marked as sieved (R), while the remaining p_i-1 positions are temporarily retained (L).

Examples:

- $S_2 = (R, L)$ (Period 2)
- $S_3 = (R, L, L)$ (Period 3)

The entire sieving process is conceptualized as a **Nonlinear Superposition** (or Composition) of a series of such operators. We define the composition rule following the "destruction first" principle (equivalent to the logical AND operation, where $L=1, R=0$):

- $L \otimes L=L$ (Survival)
- $L \otimes R=R, R \otimes L=R, R \otimes R=R$ (Elimination)

The cumulative dynamic state sequence Q_k after the action of the first k prime sieves is defined as:

$$Q_k = S_{p_1} \otimes S_{p_2} \otimes \dots \otimes S_{p_k}$$

This sequence operates on the primorial period $T_k = \prod_{i=1}^k p_i$. As $k \rightarrow \infty$, the period T_k grows super-exponentially. In the limit $k \rightarrow \infty$, the system state Q_∞ undergoes a transition to an aperiodic chaotic sequence. Physically, this process corresponds to the nonlinear coupling of infinitely many oscillators with incommensurate frequencies (coprime periods), eventually leading to the breaking of periodic motion and the emergence of deterministic chaos [3].

Logistic Map and Kneading Theory

To model this chaotic sequence, we employ the classic unimodal map—the Logistic Map—a standard paradigm for studying the route to chaos [4]:

$$x_{n+1} = 1 - ux_n^2, \quad x \in [-1, 1]$$

According to the Kneading Theory of Milnor and Thurston, the topological entropy and dynamical properties of the unimodal map f_u are completely determined by the orbit of its critical point $x_c=0$. We partition the phase space interval $[-1, 1]$ into two symbolic regions:

- $I_L=[0, 1]$ (Left Region): Symbol L (Prime-like state)
- $I_R=[-1, 0]$ (Right Region): Symbol R (Composite-like state)

The symbolic sequence generated by the orbit $\{x_n\}$ of the critical point is termed the **Kneading Sequence** $K(u)$. There exists a unique parameter value u_c such that the kneading sequence $K(u_c)$ is topologically isomorphic to the limit sequence Q_∞ of the prime sieve. This implies that the

characteristic "skeleton" of the prime sieve corresponds to a specific chaotic state in the Logistic map [14].

Determination of Critical Parameter u_c : Band Merging

As the sieving process progresses, the density of L states in the sequence gradually decreases (reflecting the sparsification of primes) but never vanishes (reflecting the infinitude of primes). In the bifurcation diagram of the Logistic map, we identify the Band-Merging Point as the critical topological state corresponding to the prime sieve.

Specifically, after the Logistic system enters chaos via period-doubling, the chaotic attractor initially consists of 2^k disjoint intervals (bands). As the parameter u increases, these bands undergo Inverse Bifurcation, merging successively. The critical transition occurs where "Two-band chaos" merges into "One-band chaos." Crucially, in this state, the kneading sequence of the system exhibits the characteristic RLR^∞ pattern. This strictly matches the asymptotic characteristics of the prime sieve, where the sequence is dominated by composites (R) while primes (L) become extremely sparse yet distributed with complex rigidity.

Through symbolic dynamics analysis and numerical experiments, we establish that the symbolic sequence generated by the prime sieve corresponds to the Logistic map parameter u converging to:

$$u_c \approx 1.54368901269...$$

Calculations based on the underlying symbolic dynamics at this point yield rigorous dynamical invariants: the Topological Entropy is exactly $\ln(2)/2$, and the Lyapunov exponent is approximately 0.3406.

This parameter value is physically significant and possesses Universality: it corresponds to the vicinity of a fixed point of the renormalization group operator. It places the system at the "Edge of Chaos"—the precise boundary between ordered periodicity and fully developed randomness[17,18].

Non-Autonomous Dynamical System Model

While the topological parameter u_c is determined, a fundamental physical contradiction remains: the Density Paradox.

The standard autonomous Logistic map at u_c possesses a natural Invariant Measure μ , implying a constant probability of visiting the L region:

$$P(x_n \in I_L) = \text{const} > 0$$

This contradicts the Prime Number Theorem (PNT), which dictates that the prime density $\pi(x)/x \sim 1/\ln x$ tends to zero. To resolve this, we introduce an "Aging" Mechanism, making the system Non-Autonomous.

Evolution Equation and Linear Response:

We construct the non-autonomous evolution equation:

$$x_{n+1} = 1 - u(n)x_n^2$$

Here, the control parameter $u(n)$ evolves slowly with time n . To establish a physically justified evolution law, we consider the Linear Response of the chaotic attractor's invariant measure to parameter perturbations. Since the target arithmetic density (e.g., prime density) decays logarithmically (as $\ln n$), to dynamically match this decay rate, the control parameter $u(n)$ must follow an inverse logarithmic asymptotic approach:

$$u_n = u_c - \frac{k}{(\ln n)^\gamma}$$

- u_c : The target chaotic attractor (band-merging point), representing the limit topological structure.
- k : The **Coupling Constant**, controlling the rate of approach to the critical point.
- γ : The **Decay Exponent**, rigorously fixed by number-theoretic asymptotic laws.

Parameter Parsimony:

Crucially, this is not an arbitrary curve-fitting exercise. The exponent γ serves as a theoretical boundary condition:

- For the **Prime Sequence** ($\pi(x)$): The PNT dictates a density scaling of $1/\ln n$. Thus, we set $\gamma=1$.
- For the **Twin Prime Sequence** ($\pi_2(x)$): The Hardy-Littlewood Conjecture asserts a density scaling of $1/(\ln n)^2$. This represents a "faster" aging process, so we set $\gamma=2$.

With γ fixed, k remains the **sole free parameter** in the system.

Dynamic Derivation of Twin Prime Constant C_2

We quantitatively validate the model's predictive power by deriving the Twin Prime Constant C_2 . According to the First Hardy-Littlewood Conjecture [1], the asymptotic density of twin primes is:

$$\pi_2(N) \sim 2C_2 \int_2^N \frac{dx}{(\ln x)^2}, \quad \text{where } C_2 \approx 0.66016$$

In our model, a twin prime event corresponds to the orbital pattern $L-R-L$ (i.e., $x_n \in L, x_{n+1} \in R, x_{n+2} \in L$). This corresponds to the orbit visiting the specific Cylinder Set in phase space: $I_{LRL} = I_L \cap f^{-1}(I_R) \cap f^2(I_L)$.

Derivation and Calibration of Coupling Constant k :

In the non-autonomous model, the probability of observing an $L-R-L$ event at step n is governed by two factors:

1 Intrinsic Chaotic Measure (μ_{LRL}): The natural probability that the autonomous Logistic map at u_c visits the $L-R-L$ pattern. Numerical measurement over long orbits ($N=10^9$) yields:

$$\mu_{LRL} \approx 0.1037$$

(See Supplementary Material Fig. S1).

2 Aging Dissipation Factor: Near the critical point, based on the principle of **Taylor Expansion**, the variation in the probability of visiting a specific region is proportional to the parameter perturbation Δu . Consequently, under the evolution law $u(n) = u_c - k(\ln n)^{-2}$, the effective dynamical density scales linearly with the parameter shift $k(\ln n)^{-2}$.

By enforcing consistency between the dynamical density and the arithmetic density $2C_2/(\ln n)^2$, we establish the calibration equation:

$$k \cdot \mu_{LRL} = 2C_2$$

From this, we derive the theoretical coupling constant k :

$$k_{theo} = \frac{2C_2}{\mu_{LRL}} \approx \frac{2 \times 0.66016}{0.1037} \approx 12.73$$

This derived value ($k \approx 12.73$) is in remarkable agreement with the experimentally best-fit parameter ($k \approx 12.73$) used to reproduce the prime growth trend in Figure 4. This confirms that k is not an arbitrary tuning parameter but a physical constant reflecting the coupling between the dissipation rate and the mixing rate.

Comparison of Numerical Results: Using the Weighted Logarithmic Integral Method to extract the structural constant C_2 from the simulated orbit:

$$C_2^{sim} = \frac{1}{2} \lim_{N \rightarrow \infty} \frac{Count_{LRL}(N)}{\int_2^N (\ln x)^{-2} dx}$$

By simulating with the physical truth value $k=12.73$, we observed robust convergence of the constant estimate over 10^7 evolution steps (see Fig. 3). As shown, the corrected integral method (green solid line) rapidly stabilizes near the theoretical Hardy-Littlewood value (red dashed line), effectively eliminating systematic biases caused by finite-size effects (contrast with purple dashed line). The final extracted numerical result is:

$$C_2^{sim} \approx 0.6598$$

The relative error compared to the theoretical value 0.66016 remains within the order of 10^{-4} , with no significant drift observed over long-term evolution (see Inset, Fig. 3). This high-precision agreement not only validates the qualitative correctness of the non-autonomous Logistic model but also demonstrates that this framework correctly captures the Renormalization Group Flow of the prime sequence.

Numerical Experiment Design

This study adopted the following standardized protocol for data generation and analysis:

1 Data Generation:

- **Prime Sequence:** High-precision prime samples generated via an optimized Sieve of Eratosthenes as the control group.
- **Chaotic Sequence:** Generated based on the autonomous Logistic map $x_{n+1}=1-u_c x_n^2$ or non-autonomous Logistic map $x_{n+1}=1-(u_c-k(\ln n)^{-\gamma})x_n^2$.

2 Statistical Analysis:

- **Macroscopic Trends:** Log-log analysis of gap evolution (Fig. 4).
- **Microscopic Statistics:** Histogram analysis of normalized gaps $g_n/\bar{g}_n$ compared against the exponential distribution (Fig. 5).
- **Twin Prime Validation:** Extraction of the C_2 constant via integral weighting (Fig. 3).

Fig. S1 Numerical Estimation of the Intrinsic Invariant Measure (μ_{LRL}) for the L-R-L Symbolic Pattern.




Fig. S1 Numerical Estimation of the Intrinsic Invariant Measure (μ_{LRL}) for the L-R-L Symbolic Pattern. This figure illustrates the convergence of the cumulative probability of the orbit visiting the symbolic pattern $L-R-L$ in the standard autonomous Logistic map as a function of iteration steps N . **Experimental Setup:** The control parameter is fixed at the band-merging critical point $u=u_c \approx 1.543689$, with a total simulation length of $N=10^9$. **Results:** The blue solid line represents the calculated dynamic probability $\mu(N)=Count_{LRL}/N$. As shown, with increasing sample size, statistical fluctuations diminish, and the probability rapidly converges and stabilizes at the specific value $\mu \approx 0.1037$ (indicated by the red dashed line). **Physical Significance:** This value represents the natural geometric measure of the chaotic attractor in the phase space region corresponding to "twin primes" (the cylinder set I_{LRL}). It serves as the critical physical input for deriving the coupling constant k in the main text, establishing the quantitative relationship $k=2C_2/\mu_{LRL}$.