

A Source-Locked Census for the Production Phase Registry: Fixed- $h_0 = 2$ Data, Missing Named-Atom Values, and Determinant-Two Packet Coordinates

Liang Wang
School of Artificial Intelligence and Automation,
Huazhong University of Science and Technology, Wuhan 430074, P.R. China
wangliang.f@gmail.com

July 2026

Abstract

The TPC-170 metric corridor controls all prefixes of every packet in a prescribed determinant-two schedule for Lebesgue-almost every fixed phase. To consume that theorem at the physical endpoint one first needs a value-bearing production phase registry: a source-located named atom, its exact phase modulo one, and its map into the precise packet coordinates on which the metric theorem acts. We audit the frozen TPC-157–172 phase-interface artifacts under canonical source locks. They prove that the relevant actual core has fixed $h_0 = 2$, and they specify the type of the missing H9 phase record. They do not contain a value-bearing named atom, a phase-value locator, a production packet schedule, or any physical-atom-to-packet row.

We give the minimal typed registry contract, including canonical Bézout representative and covariant-multiplier fields, and return `NOT_TESTABLE` with the first missing object equal to the named atom together with its phase-value source locator. The result is a scoped L1 source-interface obstruction. It is not a global nonexistence theorem, does not synthesize phase zero or any other phase, and supplies no decay exponent.

Claim boundary. The fixed- $h_0 = 2$ fact and the phase registry are separate data objects. An H9 registry has decay coordinate `NONE`. Neither an obligation label, a source hash, a phase-average theorem, nor a numerical fixture supplies a named physical phase. No fixed-atom estimate, program-positive L2, strict $1/400$, prime-pair lower bound, or twin-prime theorem is proved.

1 The exact question

TPC-171 records the missing physical phase leaf as

$$\text{H9.phase_cell_registry.} \tag{1}$$

Its six-axis signature is

axis	required value	
carrier	PHYSICAL_PHASE_REGISTRY	
phase	NAMED_FIXED_ATOM	
endpoint	NOT_APPLICABLE	
scale	DETERMINISTIC_ALL_SCALE	
decay	NONE	
support	ACTUAL_ACTIVE_SUPPORT.	(2)

The node has no arithmetic parent. This direction is deliberate: arithmetic theorems may consume a physical phase record, but cannot manufacture one [4, 9].

Definition 1.1 (Value-bearing production phase record). A production phase record contains at least:

- (i) a stable named physical atom identifier;
- (ii) its exact value $\alpha_\star \in \mathbb{T}$;
- (iii) a theorem-usable source path and locator for that value;
- (iv) a separate source lock for $h_0 = 2$;
- (v) the exact production packet schedule to which the phase is mapped; and
- (vi) source-located rows realizing the map into the packet coordinates of TPC-170.

An object that contains only the words “named phase”, only a target node such as (1), or only a schema allowing these fields is not a record in the sense of [definition 1.1](#).

2 Fixed $h_0 = 2$ is present and is not a phase

TPC-157 and TPC-159 both export the determinant-two actual-core field

$$h_0 = 2 \tag{3}$$

in their machine-readable theorem records [1, 3]. Endpoint Ledger V4 in TPC-171 likewise marks the fixed physical h_0 gate as proved on the relevant actual core [9]. These three locators are retained independently in the census.

Proposition 2.1 (Data-axis separation). *The source-backed fact (3) does not instantiate any of the first three fields of [definition 1.1](#).*

Proof. The integer h_0 is the prescribed physical shift, while $\alpha_\star$ is a character coordinate in $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. Their types and source fields differ. The cited records contain the integer 2 in a `fixed_h0` or `fixed-physical-h0` field. None gives a named-atom identifier, a phase value modulo one, or a locator for such a value. Reusing the integer 2, reducing it modulo one, or silently declaring $\alpha_\star = 0$ would be a type-changing invention. $\square$

This proposition does not weaken the fixed- h_0 result. It only prevents that genuine data fact from filling a different missing field.

3 Packet coordinates required by TPC-170

For a determinant-two packet, TPC-170 uses

$$su - ad = 2, \quad q = as, \quad t_d(z) = ad + qz, \tag{4}$$

and the coefficient sequence

$$c_{d,u}(z) = \mu(d + sz)\mu(u + az). \tag{5}$$

At ambient scale X_n , a prescribed finite list $\mathcal{P}_n$ contains a canonical fiber, a terminal scale $T_{n,p}$, and a multiplier $\rho_{n,p}$. Its ordered coordinates $z_{n,p,j}$ enter

$$S_{n,p,k}(\alpha) = \sum_{j \leq k} c_{d,u}(z_{n,p,j}) \rho_{n,p}(z_{n,p,j}) e(-\alpha z_{n,p,j}). \tag{6}$$

Thus a production crosswalk must give actual values for

$$(X_n, p, a, s, d, u, q, T_{n,p}, \rho_{n,p}, z_{n,p,j}), \tag{7}$$

the physical occurrence locator, and the common $\alpha_\star$.

The formula (6) defines a consumer interface; it is not a production row.

There is also a representative condition. For fixed (a, s) , all solutions of (4) are

$$(d, u) \mapsto (d + sr, u + ar), \quad r \in \mathbb{Z}. \quad (8)$$

TPC-170 proves that this translates the coefficient packet. The multiplier must be translated covariantly; then the twist changes by only a unit scalar. A production registry must therefore select one canonical representative or record the translation index and the covariant multiplier map [8]. The theorem (8) does not select the actual production representative by itself.

4 Frozen source census

The executable census locks the phase-interface outputs of TPC-157–161 and TPC-167–172. Its relevant findings are summarized in table 1.

Object	Frozen value	Exact meaning
fixed physical h_0	2, source-backed	One literal data gate is present.
TPC-158 production phase cell	NOT-TESTABLE	The major/minor interface has no production phase value.
TPC-161 phase artifact	MISSING	The target description is present; the required artifact is not.
TPC-171 H9 phase node	NOT-TESTABLE, evidence null	The six-axis obligation is typed but uninstantiated.
TPC-170 phase mode	Lebesgue-a.e. fixed phase	A metric quantifier, not a named atom.
named atom/value/locator rows	0/0/0	No value-bearing production phase record occurs in the seven explicitly mapped fields.
production packet-coordinate rows	0	The mapped theorem/status fields supply no row of (7).

Table 1: The fixed- h_0 fact survives the census; the production phase values do not appear.

TPC-167 proves a continuous phase- L^2 identity, TPC-168 a separated-registry density theorem, TPC-169 a maximal prefix theorem, and TPC-170 an almost-every-fixed-phase corridor [5–8]. Their claim boundaries all reject a named production phase. They are positive arithmetic inputs in their declared norms, not value-bearing entries for (1).

Theorem 4.1 (Scoped registry census). *In the frozen source-locked TPC-157–172 phase-interface corpus:*

- (i) $h_0 = 2$ is source-backed on the relevant actual core;
- (ii) the target type (2) is source-backed;
- (iii) the representative rule (8) is proved;
- (iv) none of the seven explicitly mapped fields is a value-bearing named physical atom or phase-value locator; and
- (v) those mapped fields contain no production row realizing (7).

Consequently the production phase registry has status NOT-TESTABLE.

Proof. The first three statements are exact field checks against the source-locked TPC-157, TPC-159, TPC-170 and TPC-171 outputs. The last two statements are an explicit mapped-field census: three fixed- h_0 records and four phase-obligation records. It is not a generic scanner, and newly added source fields require an explicit mapping update. The machine audit rejects any snapshot that fills a null phase value, locator, schedule, or packet row without a new source. By definition 1.1, the mapped evidence does not construct a production registry. $\square$

Remark 4.2 (Scope). [Theorem 4.1](#) is not a theorem that no physical phase description exists mathematically or in every possible archive. It says that the explicitly mapped fields in the declared frozen corpus do not provide the values needed to instantiate [definition 1.1](#). A larger corpus or a new field must be named, mapped, locked, and audited before it can change the result.

5 Decision and typed progress

The exact machine result is

$$\begin{array}{l} \text{Verdict} = \text{NOT_TESTABLE}, \\ \text{ProductionPhaseRegistryConstructed} = \text{false}, \\ \text{FirstMissing} = \text{named_physical_atom_id_and_} \\ \quad \text{phase_value_source_locator}. \end{array} \tag{9}$$

Independent missing fields include the production schedule locator, the packet-coordinate rows, and the actual representative/covariant multiplier map.

The progress levels are:

- L0: deterministic hashes, schemas, field census, and mutation diagnostics;
- L1: the scoped source-interface conclusion [theorem 4.1](#); and
- L2: none.

The fixed- $h_0 = 2$ field is a proved literal data fact, but it creates no cancellation exponent. The H9 phase leaf retains decay axis NONE.

6 Reproducible artifacts and next gate

The generator writes

```
experiments/tpc180_phase_registry_census.json
experiments/tpc180_phase_registry_census_audit.json
```

and checks both strict schemas. Source hashes have integrity semantics only. Mutation regressions reject a target obligation used as a phase value, an unsourced locator, a synthetic packet row, H9 decay promotion, a scoped census promoted to global nonexistence, and the fixed- h_0 fact promoted to L2.

The next arithmetic object remains

$$\text{H2.metric_fixed_atom_crosswalk}. \tag{10}$$

Even a future positive phase registry would supply data only. TPC-181 must separately ask whether a theorem places its named atom in the TPC-170 good set. No such theorem is inferred here.

7 Conclusion

The phase question is now split at the correct seam. Fixed $h_0 = 2$ and the determinant-two Fourier interface are present; the value-bearing named phase and its production packet crosswalk are not. The only source-compatible TPC-180 output is therefore the contract and scoped missing-value census (9). This is useful L1 localization, not fixed-atom arithmetic progress.

References

- [1] Liang Wang. Periodic approximation of literal multipliers on determinant-two möbius fibers. TPC-157 manuscript, 2026.
- [2] Liang Wang. An exact major-arc gate for additive phases: Periodic projection and a minor-arc route obstruction. TPC-158 manuscript, 2026.
- [3] Liang Wang. Dyadic-shadow lifting from good-scale möbius correlations to almost-endpoint prefixes. TPC-159 manuscript, 2026.
- [4] Liang Wang. Source-locked occurrence-to-return integration. TPC-161 manuscript, 2026.
- [5] Liang Wang. A parseval corridor for direct additive twists on the determinant-two möbius core. TPC-167 manuscript, 2026.
- [6] Liang Wang. Separated phase registries for direct core twists. TPC-168 manuscript, 2026.
- [7] Liang Wang. One phase-exceptional set for every atomic prefix. TPC-169 manuscript, 2026.
- [8] Liang Wang. Metric all-prefix return on prescribed determinant-two packet corridors. TPC-170 manuscript, 2026.
- [9] Liang Wang. Source-locked occurrence–phase return integration. TPC-171 manuscript, 2026.
- [10] Liang Wang. MVP7 on the occurrence–phase–atomic corridor. TPC-172 manuscript, 2026.