

Parity-Resolved Dynamics at a Quadratic Band-Merging Parameter: Peripheral Spectrum, Periodograms, and Two-Step Sequential Stability

Liang Wang

School of Artificial Intelligence and Automation,
Huazhong University of Science and Technology, Wuhan 430074, P.R. China
wangliang.f@gmail.com

July 2026

Abstract

Let $f_u(x) = 1 - ux^2$ and let u_c be the first band-merging parameter, characterized by

$$u_c^3 - 2u_c^2 + 2u_c - 2 = 0, \quad u_c = 1.543689012692\dots$$

The invariant interval of f_{u_c} has a genuine period-two decomposition $X = C \cup H$ with $f_{u_c}(C) = H$ and $f_{u_c}(H) = C$ modulo their common endpoint. Consequently the Koopman operator has the eigenfunction $s = \mathbf{1}_C - \mathbf{1}_H$ with eigenvalue -1 , and the raw map is ergodic but not mixing. Under an explicit two-step component spectral-gap hypothesis on an adapted density space, the Perron–Frobenius peripheral spectrum is exactly $\{1, -1\}$.

We give the exact parity-resolved operator decomposition under that hypothesis, for $n \geq 1$:

$$P^n = \Pi_+ + (-1)^n \Pi_- + R^n, \quad \|R^n\| \leq K \vartheta^n,$$

where $\Pi_+ g = h \int g$ and $\Pi_- g = sh \int sg$, with $s = \mathbf{1}_C - \mathbf{1}_H$. Hence every sufficiently regular real observable has covariance

$$\text{Cov}_\mu(\phi, \phi \circ f_{u_c}^n) = a_\phi^2 (-1)^n + O(\vartheta^n), \quad a_\phi = \int \phi s d\mu.$$

Its observable spectral measure contains an atom of mass a_ϕ^2 at frequency $1/2$ (angular frequency π), in addition to the mean atom at zero and an analytic absolutely continuous remainder. We also prove that finite empirical discrete periodograms along almost every orbit converge weakly to this measure.

For non-autonomous dynamics, the correct primitive object is the two-step block operator $Q_n = P_{2n} P_{2n-1}$. We prove an abstract adiabatic tracking theorem for a moving rank-two projector bundle and a parity-preserving sequential Birkhoff theorem under explicit uniform block memory-loss and paired-mean assumptions. Pairing consecutive iterates cancels the -1 mode and yields L^2 and almost-everywhere convergence without imposing an incompatible one-mode estimate. The exact geometry, parity eigenmode, ordinary Birkhoff theorem, and pathwise periodogram limit for each fixed bounded real observable are unconditional. Exponential operator remainders are stated under the component-gap hypothesis, and application of the sequential theorem to a prescribed nearby quadratic schedule remains conditional on verifying a common or moving two-phase bundle and uniform product estimates.

Keywords: quadratic map; band merging; Perron–Frobenius operator; Koopman spectrum; peripheral eigenvalue; periodogram; sequential dynamical system; adiabatic tracking.

MSC 2020: 37E05; 37A30; 37M25; 47A35; 47B33.

1 Introduction

At a mixing parameter, exponential memory loss is naturally expressed by a transfer-operator decomposition with a single peripheral eigenvalue 1. At a band-merging parameter the geometry is different: two cyclic components are exchanged by one iterate and mixed only by the square of the map. On an adapted density space the Perron–Frobenius operator then has peripheral eigenvalues 1 and -1 , while the Koopman operator has the corresponding eigenfunctions $\mathbf{1}$ and s . No claim is made that the full Koopman operator spectrum consists only of these points. Dropping the -1 eigenmode changes correlations, single-time density limits, and spectral measures.

This distinction refines a proposed sequential Birkhoff theorem for logarithmically drifting quadratic maps [12]. That manuscript records that all first-return times at the limiting parameter are even, whereas a later one-mode raw estimate suppresses the resulting parity coordinate. At the constant schedule $u_n \equiv u_c$, the parity observable alternates forever, so the estimate must be replaced by a two-mode formulation. The Birkhoff conclusion may nevertheless survive because Cesàro averaging cancels the alternating term.

The broader prime-dynamics program was introduced in Wang [9], and the exact band-merging geometry was isolated in the rigorous reformulation of the prime-sieve correspondence [11]. The present paper develops the operator and spectral consequences of that geometry. It is logically independent of finite-stage sieve admissibility and of the inverse prime-kneading parameter studied in Wang [10]. No arithmetic or Riemann-zero assertion is used.

Periodic decompositions and root-of-unity peripheral modes for quasi-compact transfer operators are classical [2]; so are the Herglotz representation and the spectral measure of a stationary observable [4]. Sequential memory loss and limit theorems have been developed for several classes of time-dependent interval maps [1, 3], while strong–weak stability of isolated operator clusters is governed by perturbation results such as Keller and Liverani [6]. The contribution here is not a new general cyclic spectral theorem. It is the exact diagnosis of the period-two mode at this algebraic band-merging parameter, its explicit consequence for orbit periodograms, and a two-step sequential architecture which separates contraction transverse to a rank-two bundle from the unresolved dynamics inside that bundle.

Main results

Let P be the Perron–Frobenius operator of f_{u_c} relative to Lebesgue measure, let h be the invariant density, and let $U\phi = \phi \circ f_{u_c}$ be the Koopman isometry on $L^2(\mu)$, $d\mu = h \, dx$. The results are:

R1. The band-merging map has a two-cycle partition C, H . Unconditionally, with $s = \mathbf{1}_C - \mathbf{1}_H$,

$$Us = -s, \quad P(sh) = -sh.$$

Under [Assumption 3.1](#), the Perron–Frobenius peripheral spectrum is exactly $\{1, -1\}$ and the rest contracts exponentially.

R2. Under the same component-gap hypothesis, every regular real observable admits the orthogonal decomposition

$$\phi = m_\phi + a_\phi s + \psi,$$

and

$$\text{Cov}_\mu(\phi, \phi \circ f_{u_c}^n) = a_\phi^2 (-1)^n + O(v^n).$$

Thus its spectral measure is

$$\sigma_\phi = m_\phi^2 \delta_0 + a_\phi^2 \delta_{1/2} + w_\psi(t) \, dt,$$

where w_ψ is nonnegative and real analytic.

- R3.** For every fixed bounded real observable and μ -almost every initial point, the finite DFT periodogram measures converge weakly to σ_ϕ . The parity atom, when its mass is nonzero, is therefore a pathwise spectral feature, not merely an operator-level formalism.
- R4.** For a sequence of parity-preserving maps, uniform exponential memory loss is imposed on the two-step block products, rather than on the raw one-step maps. Under a common paired-mean hypothesis, pairing one observation from each phase cancels the alternating mode, and the combined Birkhoff average converges in L^2 and almost everywhere to that common limit. At the constant schedule the limit is the equal component average.
- R5.** More generally, if $Q_n = P_{2n}P_{2n-1}$ is equipped with a moving rank-two projector bundle $E_n = \text{Ran } \Pi_n$ and the complementary products contract, then the non-autonomous density tracks E_n with an explicit convolution bound. A logarithmic schedule has summable parameter increments, but its internal two-dimensional cocycle must still be analyzed: convergence $Q_n \rightarrow P_{u_c}^2$ alone does not decide whether the parity mode persists or decays.

The exact Markov geometry, invariant component measures, parity eigenmode, Birkhoff limit, and pathwise periodogram theorem for each fixed bounded real observable apply unconditionally to the algebraic Misiurewicz map by classical interval-map theory [2, 5, 8, 13]. The strong operator remainder and analytic spectral-density statements are conditional on [Assumption 3.1](#); a tower construction is a standard route to that hypothesis, but the present paper does not build the required interval density space from scratch. The sequential results are likewise abstract theorems with explicit hypotheses. We do not assert that an arbitrary one-sided logarithmic schedule lies in a single uniform tower-supporting parameter set.

2 Exact band-merging geometry

Let

$$f_u(x) = 1 - ux^2, \quad 0 < u \leq 2. \quad (1)$$

The following proposition is recalled from Wang [11]; its short calculation is reproduced for self-containment.

Proposition 2.1 (Algebraic parameter and Markov partition (recalled)). *The polynomial*

$$F(u) = u^3 - 2u^2 + 2u - 2 \quad (2)$$

has a unique zero u_c in $(1, 2)$. Put $b = u_c - 1$. Then

$$0 \mapsto 1 \mapsto -b \mapsto b \mapsto b. \quad (3)$$

On $X = [-b, 1]$, define

$$J_L = [-b, 0], \quad J_0 = [0, b], \quad J_1 = [b, 1].$$

Up to common endpoints,

$$f_{u_c}(J_L) = J_1, \quad f_{u_c}(J_0) = J_1, \quad f_{u_c}(J_1) = J_L \cup J_0. \quad (4)$$

The transition matrix is

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}, \quad \text{spec}(A) = \{0, \sqrt{2}, -\sqrt{2}\}. \quad (5)$$

Proof. One has $F(1) = -1$, $F(2) = 2$, and

$$F'(u) = 3 \left(u - \frac{2}{3} \right)^2 + \frac{2}{3} > 0.$$

Thus the zero is unique. Equation $F(u_c) = 0$ is equivalent to $1 - u_c b^2 = b$, while $1 - u_c = -b$, proving (3). On $[-b, 0]$ the map is increasing from b to 1; on $[0, b]$ it is decreasing from 1 to b ; and on $[b, 1]$ it is decreasing from b to $-b$. This proves (4). Finally,

$$\det(\lambda I - A) = \lambda(\lambda^2 - 2),$$

which gives (5). □

Define, modulo the common endpoint b ,

$$C = [-b, b], \quad H = [b, 1]. \quad (6)$$

Corollary 2.2 (Exact period two). *The map satisfies*

$$f_{u_c}(C) = H, \quad f_{u_c}(H) = C. \quad (7)$$

Moreover $f_{u_c}^2$ is topologically mixing on each component with respect to its Markov partition. Thus the raw Markov system has period exactly two.

Proof. The image relations follow from (4). Squaring A separates the state J_1 from the two states J_L, J_0 ; the restriction to $\{J_L, J_0\}$ is the positive matrix with every entry equal to 1. More explicitly,

$$f_{u_c}^2(J_L) = f_{u_c}^2(J_0) = C.$$

On H , put $c_H = u_c^{-1/2}$. Since $u_c b^2 = 1 - b < 1$ and $u_c > 1$, one has $b < c_H < 1$. Split

$$H_0 = [b, c_H], \quad H_1 = [c_H, 1].$$

Then $f_{u_c}^2(H_0) = f_{u_c}^2(H_1) = H$. Thus both two-step component graphs have full branches and are aperiodic and mixing; see Lind and Marcus [7, Chapter 4]. □

The critical orbit is finite and lands on the repelling fixed point b : indeed $F(3/2) < 0$ gives $u_c > 3/2$, hence $b > 1/2$ and $|f'_{u_c}(b)| = 2u_c b > 3/2$. The component critical orbits also terminate there:

$$0 \xrightarrow{f_{u_c}^2} -b \xrightarrow{f_{u_c}^2} b, \quad c_H \xrightarrow{f_{u_c}^2} 1 \xrightarrow{f_{u_c}^2} b.$$

Thus the two component maps are nonflat Misiurewicz maps. Classical results give an absolutely continuous invariant probability measure and exponential correlation estimates on each mixing component of the square for standard Hölder or tower-regular observable classes [5, 8, 13]. An operator-norm formulation requires an adapted tower or weighted space which accommodates the postcritical singularities of the invariant density; it need not be ordinary $BV(X)$.

Proposition 2.3 (Invariant component measures). *The map f_{u_c} has a unique absolutely continuous invariant probability measure μ . It is ergodic and not mixing. One has*

$$\mu(C) = \mu(H) = \frac{1}{2}.$$

The probability measures

$$\mu_C = 2\mu|_C, \quad \mu_H = 2\mu|_H \quad (8)$$

are the two mixing absolutely continuous invariant measures of $f_{u_c}^2$, and

$$(f_{u_c})_* \mu_C = \mu_H, \quad (f_{u_c})_* \mu_H = \mu_C.$$

Proof. The Misiurewicz results just cited give unique mixing absolutely continuous invariant probabilities ν_C and ν_H for the two primitive components of $f_{u_c}^2$. Component interchange implies that $(f_{u_c})_*\nu_C$ is an absolutely continuous $f_{u_c}^2$ -invariant probability on H , hence equals ν_H by uniqueness; similarly $(f_{u_c})_*\nu_H = \nu_C$. Therefore $\mu = (\nu_C + \nu_H)/2$ is f_{u_c} -invariant. Conversely, any absolutely continuous f_{u_c} -invariant probability restricts, after normalization, to ν_C and ν_H , and the interchange forces equal weights. Thus μ is unique, $\mu_C = \nu_C$, $\mu_H = \nu_H$, and (8)–(7) follow. A measurable f_{u_c} -invariant set has square-map-invariant intersections with both components. Component ergodicity and the interchange then force its measure to be 0 or 1, proving ergodicity. Nonmixing follows from the nontrivial eigenfunction constructed below. $\square$

3 The Perron–Frobenius peripheral spectrum is $\{1, -1\}$

Let m denote Lebesgue measure on X , write $d\mu = h dm$, and let P be the Perron–Frobenius operator of f_{u_c} relative to m :

$$\int_X \chi P g dm = \int_X (\chi \circ f_{u_c}) g dm.$$

We isolate the exact operator input needed below.

Assumption 3.1 (Two-step component spectral gap). There is a Banach space $\mathcal{B} \hookrightarrow L^1(m)$ of signed densities such that multiplication by $\mathbf{1}_C, \mathbf{1}_H$ and the operator P are bounded on $\mathcal{B}$, and $h \in \mathcal{B}$. Put

$$h_C = 2h\mathbf{1}_C, \quad h_H = 2h\mathbf{1}_H.$$

There exist $M < \infty$ and $0 < \rho < 1$ such that, for $D \in \{C, H\}$ and every $g \in \mathcal{B}$ supported in D ,

$$\left\| (P^2)^n g - h_D \int_D g dm \right\|_{\mathcal{B}} \leq M \rho^n \|g\|_{\mathcal{B}}. \quad (9)$$

A mixing exponential-tail tower for each component is a standard route toward [Assumption 3.1](#) [2, 13]. Passing from a tower operator to the precise interval-density norm in the assumption requires a specified quotient or weighted-space construction. We therefore retain [Assumption 3.1](#) explicitly instead of identifying it with ordinary BV decay.

All spectral statements below are understood on the complexification $\mathcal{B}_{\mathbb{C}}$ of the signed-density space.

Set

$$s = \mathbf{1}_C - \mathbf{1}_H. \quad (10)$$

The value at b is irrelevant because μ is nonatomic.

Theorem 3.2 (Parity-resolved Perron–Frobenius decomposition). Under [Assumption 3.1](#), define

$$\Pi_+ g = h \int_X g dm, \quad \Pi_- g = sh \int_X sg dm. \quad (11)$$

Then Π_+ and Π_- are mutually annihilating rank-one spectral projectors,

$$Ph = h, \quad P(sh) = -sh, \quad (12)$$

and, for every $n \geq 1$,

$$P^n = \Pi_+ + (-1)^n \Pi_- + R^n, \quad R = P(I - \Pi_+ - \Pi_-). \quad (13)$$

There exist $K < \infty$ and $0 < \vartheta < 1$ such that

$$\|R^n\|_{\mathcal{B} \rightarrow \mathcal{B}} \leq K \vartheta^n. \quad (14)$$

In particular, the peripheral spectrum of P is exactly $\{1, -1\}$, and both eigenvalues are algebraically simple.

Proof. The component interchange and invariance give

$$P(h\mathbf{1}_C) = h\mathbf{1}_H, \quad P(h\mathbf{1}_H) = h\mathbf{1}_C.$$

Adding and subtracting yields (12). The corresponding left eigenfunctionals are

$$\ell_+(g) = \int g \, dm, \quad \ell_-(g) = \int sg \, dm,$$

because $s \circ f_{u_c} = -s$ almost everywhere. Since

$$\int sh \, dm = \mu(C) - \mu(H) = 0, \quad \int s^2 h \, dm = 1,$$

the formulas in (11) are normalized projectors and $\Pi_+ \Pi_- = \Pi_- \Pi_+ = 0$.

For $g = g_C + g_H$, (9) gives

$$(P^2)^n g = h_C \int_C g \, dm + h_H \int_H g \, dm + O_{\mathcal{B}}(\rho^n \|g\|_{\mathcal{B}}).$$

The main term equals $(\Pi_+ + \Pi_-)g$. One further application of the bounded operator P changes it to $(\Pi_+ - \Pi_-)g$. Thus the even and odd remainders decay geometrically. Because P commutes with the spectral projectors, while $R\Pi_{\pm} = \Pi_{\pm}R = 0$, this is precisely (13)–(14), after adjusting the constant and taking $\sqrt{\rho} \leq \vartheta < 1$. Finally, the 1-eigenspaces of P^2 on the two mixing components are spanned by h_C, h_H . On their span, P is the component-swap matrix, whose eigenvalues 1, -1 are simple. No other peripheral eigenvalue remains. $\square$

Remark 3.3 (Koopman and Perron–Frobenius eigenvectors). For the Koopman isometry $U\phi = \phi \circ f_{u_c}$ on $L^2(\mu)$,

$$U\mathbf{1} = \mathbf{1}, \quad Us = -s.$$

For the Lebesgue-reference Perron–Frobenius operator, the right eigenvectors are h and sh , not generally $\mathbf{1}$ and s . The statement $Ps = -s$ is therefore usually false; it becomes valid only for the transfer operator normalized relative to μ .

4 Parity-resolved correlations and observable spectra

The following formula is the correct replacement for a one-step raw mixing estimate.

Theorem 4.1 (Parity-resolved correlation formula). *Let ϕ, χ be real observables such that $\phi h \in \mathcal{B}$ and $\chi \in L^\infty(\mu)$. Then, for every $n \geq 0$,*

$$\begin{aligned} \int_X \phi(\chi \circ f_{u_c}^n) \, d\mu &= \left(\int_X \phi \, d\mu \right) \left(\int_X \chi \, d\mu \right) \\ &\quad + (-1)^n \left(\int_X \phi s \, d\mu \right) \left(\int_X \chi s \, d\mu \right) + O(\vartheta^n \|\phi h\|_{\mathcal{B}} \|\chi\|_{\infty}). \end{aligned} \tag{15}$$

Proof. By transfer duality and theorem 3.2,

$$\int_X \phi(\chi \circ f_{u_c}^n) \, d\mu = \int_X \chi P^n(\phi h) \, dm.$$

For $n \geq 1$, the Π_+ term is the product of the ordinary means, the Π_- term is the alternating product of the two parity means, and the R^n term is bounded by (14) and the L^1 embedding of $\mathcal{B}$. The case $n = 0$ follows directly after enlarging the implicit constant. $\square$

Taking $\phi = \chi = s$ gives

$$\int_X s(s \circ f_{u_c}^n) d\mu = (-1)^n. \quad (16)$$

Thus no estimate of the form

$$\left| \int \phi(\chi \circ f_{u_c}^n) d\mu - \int \phi d\mu \int \chi d\mu \right| \leq C\vartheta^n$$

can hold for all regular observables.

For a real observable $\phi \in L^2(\mu)$, set

$$m_\phi = \int_X \phi d\mu, \quad a_\phi = \int_X \phi s d\mu, \quad \psi = \phi - m_\phi - a_\phi s. \quad (17)$$

Since $\mathbf{1}$ and s are orthonormal in $L^2(\mu)$, this is the orthogonal decomposition

$$\phi = m_\phi \mathbf{1} + a_\phi s + \psi, \quad \langle \psi, \mathbf{1} \rangle = \langle \psi, s \rangle = 0. \quad (18)$$

Corollary 4.2 (Covariance decomposition). *If $\phi h \in \mathcal{B}$ and $\phi \in L^\infty(\mu)$, then*

$$\text{Cov}_\mu(\phi, \phi \circ f_{u_c}^n) = a_\phi^2(-1)^n + r_\phi(n), \quad |r_\phi(n)| \leq K_\phi \vartheta^n, \quad (19)$$

where

$$r_\phi(n) = \int_X \psi(\psi \circ f_{u_c}^n) d\mu.$$

Proof. Apply [theorem 4.1](#) to ψ . The cross terms with $\mathbf{1}$ vanish by centering. The cross terms with s vanish exactly: for example,

$$\int_X s(\psi \circ f_{u_c}^n) d\mu = \int_X \psi P^n(sh) dm = (-1)^n \int_X \psi s d\mu = 0.$$

The remaining parity term is $a_\phi^2(-1)^n$. □

4.1 The observable spectral measure

We use the frequency circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ and normalized Haar measure dt . Although the Koopman operator of a noninvertible interval map is an isometry rather than a unitary operator, the stationary correlation sequence is positive definite. Herglotz' theorem, equivalently the unitary natural extension, therefore defines a unique finite positive measure σ_ϕ by

$$\hat{\sigma}_\phi(n) := \int_{\mathbb{T}} e^{2\pi i n t} d\sigma_\phi(t) = \int_X \phi(\phi \circ f_{u_c}^n) d\mu, \quad n \geq 0, \quad (20)$$

with negative coefficients given by conjugation.

Theorem 4.3 (Zero and parity atoms). *Under the hypotheses of [corollary 4.2](#),*

$$\sigma_\phi = m_\phi^2 \delta_0 + a_\phi^2 \delta_{1/2} + w_\psi(t) dt, \quad (21)$$

where

$$w_\psi(t) = \sum_{n \in \mathbb{Z}} r_\phi(n) e^{-2\pi i n t}, \quad r_\phi(-n) = \overline{r_\phi(n)}. \quad (22)$$

The function w_ψ is nonnegative and real analytic. In angular frequency, the atom at $1/2$ is the atom at π .

Proof. The Fourier coefficients of $m_\phi^2 \delta_0$ are m_ϕ^2 , and those of $a_\phi^2 \delta_{1/2}$ are $a_\phi^2 (-1)^n$. The remainder coefficients in (19) decay exponentially in both directions, so the Fourier series (22) converges absolutely in a complex strip and defines a real-analytic function. It is nonnegative because it is the Radon–Nikodym density of the positive spectral measure of the orthogonal component ψ . $\square$

Exponential decay is stronger than is needed for absolute continuity. Summability of $r_\phi(n)$ already gives a continuous density; the tower gap upgrades it to analyticity.

Corollary 4.4 (The left-symbol observable). *Let $\ell = \mathbf{1}_{J_L}$, assume $\ell h \in \mathcal{B}$ as in the Markov-adapted tower construction, and put $q = \mu(J_L)$. Then*

$$m_\ell = q, \quad a_\ell = q,$$

and therefore

$$\sigma_\ell = q^2 \delta_0 + q^2 \delta_{1/2} + w_\ell(t) dt, \quad w_\ell = w_{\ell-q\delta_s}. \quad (23)$$

Centering ℓ removes the zero-frequency atom but leaves the parity atom unchanged. For the parity observable itself, $\sigma_s = \delta_{1/2}$ exactly.

Proof. The set J_L lies in C , so $s = 1$ on J_L modulo endpoints. Hence $\int \ell s d\mu = \int \ell d\mu = q$. The last assertion follows from $s \circ f_{u_c} = -s$. $\square$

5 Pathwise convergence of finite periodograms

The spectral measure above is recovered from almost every single orbit. Let $T = f_{u_c}$, let $\phi \in L^\infty(\mu)$ be real-valued, and define the unitary DFT of the first N observations by

$$\widehat{\phi}_{N,x}(j) = \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} \phi(T^n x) e^{-2\pi i j n / N}, \quad 0 \leq j < N. \quad (24)$$

The associated finite measure on $\mathbb{T}$ is

$$\mathcal{I}_{N,x} = \frac{1}{N} \sum_{j=0}^{N-1} |\widehat{\phi}_{N,x}(j)|^2 \delta_{j/N}. \quad (25)$$

Discrete Parseval gives

$$\mathcal{I}_{N,x}(\mathbb{T}) = \frac{1}{N} \sum_{n=0}^{N-1} |\phi(T^n x)|^2. \quad (26)$$

Theorem 5.1 (Pathwise periodogram limit). *For μ -almost every x ,*

$$\mathcal{I}_{N,x} \Longrightarrow \sigma_\phi. \quad (27)$$

If $\|\phi\|_{L^2(\mu)} > 0$, then for almost every x the mass $\mathcal{I}_{N,x}(\mathbb{T})$ is positive for all sufficiently large N , and the corresponding normalized probability measures converge weakly to $\sigma_\phi / \|\phi\|_2^2$.

Proof. For every fixed integer $r \geq 0$, discrete Wiener–Khinchin gives

$$\widehat{\mathcal{I}_{N,x}}(r) = \frac{1}{N} \sum_{n=0}^{N-1} \phi(T^n x) \phi(T^{n+r \bmod N} x). \quad (28)$$

At most r summands wrap around the endpoint. Since ϕ is bounded, their total contribution is $O_\phi(r/N)$. The remaining average differs by $o(1)$ from the Birkhoff average of the integrable observable

$$x \longmapsto \phi(x) \phi(T^r x).$$

The map T is ergodic by [proposition 2.3](#), so for almost every x the right-hand side of (28) converges to $\hat{\sigma}_\phi(r)$. Intersecting the full-measure sets over the countably many r handles all nonnegative coefficients; the negative coefficients follow by conjugation. Equation (26) and Birkhoff give convergence of the total masses. Density of trigonometric polynomials in $C(\mathbb{T})$ now proves weak convergence. $\square$

When N is even, the parity mode lies exactly in the bin $j = N/2$. When N is odd it leaks across the grid, with its largest concentration near the closest bins, but [theorem 5.1](#) shows that the leaked mass still converges weakly to the atom at $1/2$. The theorem concerns measure convergence; it does not claim pointwise consistency of individual unsmoothed periodogram ordinates.

6 Why Birkhoff averages survive the -1 mode

The missing parity mode invalidates one-step mixing and one-step convergence of general densities, but it does not invalidate Cesàro convergence.

Proposition 6.1 (Alternating single-time expectations). *Let $g \in \mathcal{B}$ be a probability density and let ϕ be bounded. For every $n \geq 0$,*

$$\int_X \phi \circ f_{u_c}^n g \, dm = m_\phi + (-1)^n a_\phi \int_X s g \, dm + O(\vartheta^n \|g\|_{\mathcal{B}} \|\phi\|_\infty). \quad (29)$$

Unless $a_\phi = 0$ or the initial phase masses are balanced, the expectation has two alternating limits rather than one.

Proof. For $n \geq 1$, insert (13) into

$$\int \phi \circ f_{u_c}^n g \, dm = \int \phi P^n g \, dm.$$

The case $n = 0$ follows directly after enlarging the implicit constant. $\square$

Proposition 6.2 (Paired autonomous Birkhoff theorem). *For every $\phi \in L^1(\mu)$, for μ -almost every x ,*

$$\begin{aligned} \frac{1}{2N} \sum_{n=0}^{2N-1} \phi(f_{u_c}^n x) &= \frac{1}{2} \left(\frac{1}{N} \sum_{j=0}^{N-1} \phi(f_{u_c}^{2j} x) + \frac{1}{N} \sum_{j=0}^{N-1} \phi(f_{u_c}^{2j+1} x) \right) \\ &\longrightarrow \frac{1}{2} \left(\int \phi \, d\mu_C + \int \phi \, d\mu_H \right) = \int \phi \, d\mu. \end{aligned} \quad (30)$$

If $\phi \in L^\infty(\mu)$ and $\phi h \in \mathcal{B}$, the $L^2(\mu)$ variance is $O_\phi(N^{-1})$.

Proof. If $x \in C$, the even iterates are governed by the mixing system $(C, f_{u_c}^2, \mu_C)$ and the odd iterates by its transported copy on H ; for $x \in H$ the roles reverse. Birkhoff's theorem on each component proves the almost-everywhere limit. For the variance, summing (19) over a block of $2N$ observations gives a parity contribution

$$\frac{a_\phi^2}{(2N)^2} \left| \sum_{n=0}^{2N-1} (-1)^n \right|^2 = 0,$$

while the exponentially summable remainder contributes $O_\phi(N^{-1})$. $\square$

Thus the correct logic is

two-step component mixing $\implies$ paired Cesàro convergence,

not one-step raw mixing. The next sections formulate the same mechanism for non-autonomous operators.

7 A moving rank-two bundle for two-step operators

Let $P_n = P_{u_n}$ be Markov transfer operators and group them into

$$Q_n = P_{2n}P_{2n-1}. \quad (31)$$

At the limiting parameter, $Q_* = P_{u_c}^2$ has a two-dimensional 1-eigenspace, even though the eigenvalues 1 and -1 of P_{u_c} are individually simple. This rank-two cluster is the stable object for perturbation theory.

We state a strong-weak abstract theorem. Let

$$\mathcal{B}_s \hookrightarrow \mathcal{B}_w \hookrightarrow L^1(m), \quad |g|_w \leq C\|g\|_s.$$

For $n \geq 0$, let ξ_n be a block parameter and let $\Pi_n \in \mathcal{L}(\mathcal{B}_s) \cap \mathcal{L}(\mathcal{B}_w)$ be a rank-two projector, with both operator norms bounded uniformly in n . For $n \geq 1$, assume that $Q_n \in \mathcal{L}(\mathcal{B}_s)$ commutes with Π_n , and put

$$E_n = \text{Ran } \Pi_n, \quad R_n = Q_n(I - \Pi_n). \quad (32)$$

Assumption 7.1 (Sequential rank-two gap). The following bounds hold with constants independent of the allowed sequence.

(B1) Every orbit $g_n = Q_n \cdots Q_1 g_0$ with $g_0 \in \mathcal{B}_s$ satisfies

$$\sup_{n \geq 0} \|g_n\|_s \leq C\|g_0\|_s.$$

(B2) For some $0 < \rho < 1$, all $1 \leq j \leq n$ and $v \in \mathcal{B}_s$,

$$|R_n R_{n-1} \cdots R_j v|_w \leq C \min\{|v|_w, \rho^{n-j+1}\|v\|_s\}. \quad (33)$$

(B3) For the block parameters ξ_n in a metric space, there is $0 < \eta \leq 1$ such that

$$|(\Pi_n - \Pi_{n-1})v|_w \leq C\delta_n^\eta \|v\|_s, \quad \delta_n = d(\xi_n, \xi_{n-1}), \quad (34)$$

while $\|\Pi_n - \Pi_{n-1}\|_{\mathcal{B}_s \rightarrow \mathcal{B}_s}$ is uniformly bounded.

The product estimate (33) is a genuine cocycle hypothesis. It does not follow merely because each autonomous Q_n has a spectral gap. In applications it must be proved from a matched inducing construction, a cone/minorization argument, or a sequential Lasota–Yorke and covering theorem.

Theorem 7.2 (Rank-two adiabatic tracking). Under [Assumption 7.1](#), let

$$g_n = Q_n \cdots Q_1 g_0, \quad z_n = (I - \Pi_n)g_n.$$

Then

$$|z_n|_w \leq C\rho^n \|z_0\|_s + C\|g_0\|_s \sum_{j=1}^n \min\{\delta_j^\eta, \rho^{n-j+1}\}. \quad (35)$$

If, in addition,

$$\|(\Pi_n - \Pi_{n-1})v\|_s \leq C\delta_n^\eta \|v\|_s, \quad (36)$$

then the sharper bound

$$|z_n|_w \leq C\rho^n \|z_0\|_s + C\|g_0\|_s \sum_{j=1}^n \rho^{n-j+1} \delta_j^\eta \quad (37)$$

holds.

Proof. Since Q_n commutes with Π_n ,

$$\begin{aligned} z_n &= (I - \Pi_n)Q_n g_{n-1} \\ &= R_n z_{n-1} + R_n(\Pi_{n-1} - \Pi_n)g_{n-1}. \end{aligned}$$

Iterating this identity gives the exact formula

$$z_n = R_n \cdots R_1 z_0 + \sum_{j=1}^n R_n \cdots R_j (\Pi_{j-1} - \Pi_j) g_{j-1}.$$

The strong norms of the g_{j-1} are uniformly bounded by (B1). For each summand, the first entry in the minimum in (33), together with (34), gives $C\delta_j^\eta \|g_0\|_s$; the second gives $C\rho^{n-j+1} \|g_0\|_s$ using the uniform strong projector bound. Taking the better of the two proves (35). Under (36), apply the exponentially contracting branch of (33) directly to obtain $C\rho^{n-j+1} \delta_j^\eta \|g_0\|_s$, which proves (37). $\square$

7.1 Logarithmic schedules

For the rest of this subsection, take the product metric

$$d((a, b), (a', b')) = |a - a'| + |b - b'|.$$

Suppose the block parameter is

$$\xi_n = (u_{2n-1}, u_{2n})$$

and the schedule is genuinely specified by

$$u_n = u_* - c [\log(n + n_0)]^{-\beta}, \quad \beta > 0, \quad n_0 > 1. \quad (38)$$

The offset and the increment have different scales:

$$|u_n - u_*| = O((\log n)^{-\beta}), \quad |u_{n+1} - u_n| = O\left(\frac{1}{n(\log n)^{\beta+1}}\right). \quad (39)$$

An envelope bound $|u_n - u_*| \leq C(\log n)^{-\beta}$ alone does not imply the increment estimate.

Corollary 7.3 (Adiabatic rate for logarithmic drift). *Define*

$$\text{dist}_w(g, E) = \inf_{e \in E} |g - e|_w.$$

For the explicit schedule (38),

$$\text{dist}_w(g_n, E_n) = O\left(n^{-\eta} (\log n)^{1-\eta(\beta+1)}\right) \quad (40)$$

under (35). Under the strong estimate (37),

$$\text{dist}_w(g_n, E_n) = O\left(n^{-\eta} (\log n)^{-\eta(\beta+1)}\right). \quad (41)$$

If, in addition,

$$\|\Pi(\xi) - \Pi(\xi_*)\|_{s \rightarrow w} \leq C d(\xi, \xi_*)^\eta,$$

where $\Pi_n = \Pi(\xi_n)$, $\Pi_* = \Pi(\xi_*)$, and $E_* = \text{Ran } \Pi_*$, then

$$\text{dist}_w(g_n, E_*) = O((\log n)^{-\beta\eta}). \quad (42)$$

Proof. The mean-value theorem applied to (38) gives $\delta_n = O(n^{-1}(\log n)^{-\beta-1})$. In (35), split the sum at $n - L_n$ with L_n proportional to $\log n$. The old terms are exponentially small, while the last L_n terms are bounded by $L_n \delta_n^\eta$, giving (40). A geometric convolution with a regularly varying sequence is of the order of its final term, proving (41). Finally,

$$\begin{aligned} \text{dist}_w(g_n, E_*) &\leq |(I - \Pi_*)g_n|_w \\ &\leq |(I - \Pi_n)g_n|_w + |(\Pi_n - \Pi_*)g_n|_w, \end{aligned}$$

and (B1) plus projector continuity gives the rate in (42). $\square$

The theorem deliberately stops at the rank-two bundle. The frozen restriction $\Pi_n Q_n|_{E_n}$ is not itself the non-autonomous internal cocycle. The relevant transport is

$$B_n = \Pi_n Q_n \Pi_{n-1}|_{E_{n-1}} : E_{n-1} \longrightarrow E_n. \quad (43)$$

In moving frames $V_n : \mathbb{C}^2 \rightarrow E_n$, one must analyze the products of

$$M_n = V_n^{-1} \Pi_n Q_n V_{n-1}.$$

Only when the parity direction forms a covariant one-dimensional subbundle does this reduce to scalar multipliers ζ_n ; in that special case persistence may be governed by products $\prod_n \zeta_n$ and conditions such as summability of $1 - |\zeta_n|$. In general the matrices need not commute. Operator convergence $Q_n \rightarrow Q_*$ does not decide the internal cocycle.

8 A parity-preserving sequential Birkhoff theorem

We now give a clean situation in which the rank-two bundle has fixed phase coordinates and a full Birkhoff theorem follows. Let $X = X^0 \sqcup X^1$ modulo the reference measure. Let $T_n : X \rightarrow X$ be nonsingular maps with transfer operators P_n , and put

$$\Phi_n = T_n \circ \cdots \circ T_1, \quad \Phi_0 = \text{id}.$$

Assumption 8.1 (Uniform parity-preserving block memory loss). The following conditions hold.

(S1) Every map swaps the phases:

$$T_n(X^i) \subset X^{1-i} \quad \text{modulo } m, \quad i \in \{0, 1\}.$$

Hence $Q_n = P_{2n} P_{2n-1}$ preserves densities supported in each X^i .

(S2) There are Banach spaces $\mathcal{B}_i$ of signed densities supported in X^i , continuously embedded in L^1 , with $Q_n : \mathcal{B}_i \rightarrow \mathcal{B}_i$. For every allowed normalized initial density $g_{0,i} \in \mathcal{B}_i$,

$$\sup_{n \geq 0} \|Q_n \cdots Q_1 g_{0,i}\|_{\mathcal{B}_i} \leq C \|g_{0,i}\|_{\mathcal{B}_i}. \quad (44)$$

(S3) For every zero-mass $v \in \mathcal{B}_i$ and all $n \geq j$,

$$\|Q_n Q_{n-1} \cdots Q_j v\|_1 \leq C \rho^{n-j+1} \|v\|_{\mathcal{B}_i}, \quad 0 < \rho < 1. \quad (45)$$

(S4) The block observables used below satisfy

$$\sup_n \|G_n\|_\infty \leq \|\phi\|_\infty, \quad \sup_n \|M_{G_n}\|_{\mathcal{B}_i \rightarrow \mathcal{B}_i} \leq C_\phi, \quad (46)$$

where $M_{G_n} v = G_n v$.

(S5) For each allowed initial density conditioned on phase i , and each observable ϕ under consideration, the conditional paired means $c_{n,i}$ defined in (51) converge to a common value $m_*(\phi)$, independent of i and of that initial density.

Here is a concrete sufficient condition for the last clause. Let $g_{n-1,i}$ be the conditional density at the beginning of block n . Suppose there are limiting densities h_*^0, h_*^1 such that

$$\|g_{n-1,i} - h_*^i\|_1 + \|P_{2n-1}g_{n-1,i} - h_*^{1-i}\|_1 \longrightarrow 0. \quad (47)$$

Then, for bounded ϕ , transfer duality gives (S5) with

$$m_*(\phi) = \frac{1}{2} \left(\int_X \phi h_*^0 dm + \int_X \phi h_*^1 dm \right). \quad (48)$$

For y at the beginning of block n , define

$$G_n(y) = \frac{1}{2} [\phi(T_{2n-1}y) + \phi(T_{2n}T_{2n-1}y)]. \quad (49)$$

If $Y_{n-1} = \Phi_{2n-2}(x)$, then

$$\frac{1}{2M} \sum_{r=1}^{2M} \phi(\Phi_r(x)) = \frac{1}{M} \sum_{n=1}^M G_n(Y_{n-1}). \quad (50)$$

For an initial law conditioned on phase i , let

$$c_{n,i} = \mathbb{E}_i[G_n(Y_{n-1})]. \quad (51)$$

The dependence of $c_{n,i}$ on the fixed conditioned initial density is suppressed in the notation.

Theorem 8.2 (Parity-preserving sequential Birkhoff theorem). *Under [Assumption 8.1](#), for every allowed bounded observable ϕ and every initial probability density in $\mathcal{B}_0 \oplus \mathcal{B}_1$,*

$$\frac{1}{N} \sum_{r=1}^N \phi(\Phi_r) \longrightarrow m_*(\phi) \quad (52)$$

in $L^2(\nu_{g_0})$ and ν_{g_0} -almost everywhere, where $d\nu_{g_0} = g_0 dm$. Quantitatively, for even $N = 2M$ and either initial phase,

$$\left\| \frac{1}{2M} \sum_{r=1}^{2M} \phi(\Phi_r) - m_*(\phi) \right\|_{L^2} \leq \frac{C_{\phi, g_0}}{\sqrt{M}} + \frac{1}{M} \sum_{n=1}^M |c_{n,i} - m_*(\phi)|. \quad (53)$$

If $|c_{n,i} - m_*(\phi)| = O((\log(n+1))^{-\kappa})$, then the deterministic bias in (53) is $O((\log(M+1))^{-\kappa})$.

Proof. Condition on an initial phase i . Let $g_{m-1,i}$ be the density of Y_{m-1} in that phase. The signed density

$$v_{m,i} = (G_m - c_{m,i})g_{m-1,i}$$

has zero mass and uniformly bounded $\mathcal{B}_i$ norm by [equations \(44\) and \(46\)](#). More explicitly, transfer duality gives

$$\begin{aligned} & \text{Cov}_i(G_m(Y_{m-1}), G_n(Y_{n-1})) \\ &= \int_X (G_n - c_{n,i}) Q_{n-1} \cdots Q_m [(G_m - c_{m,i}) g_{m-1,i}], dm. \end{aligned}$$

Thus (45) gives, for $n > m$,

$$|\text{Cov}_i(G_m(Y_{m-1}), G_n(Y_{n-1}))| \leq C_{\phi, g_0} \rho^{n-m}. \quad (54)$$

Summing the diagonal terms and the geometric covariance tail yields

$$\text{Var}_i \left(\frac{1}{M} \sum_{n=1}^M G_n(Y_{n-1}) \right) \leq \frac{C_{\phi, g_0}}{M}. \quad (55)$$

The triangle inequality between the random average, its expectation, and $m_*(\phi)$ proves (53). Clause (S5) and Cesàro convergence prove the L^2 limit.

For almost-everywhere convergence, take $M_j = j^2$. Chebyshev's inequality, (55), and Borel–Cantelli show that the paired average minus its expectation converges to zero along M_j almost surely. Since $M_{j+1}/M_j \rightarrow 1$ and G_n is uniformly bounded, the gaps between successive squares are filled by the standard interpolation estimate. The conditional expectations converge to $m_*(\phi)$, proving the full paired limit. The two initial phases cover a general density, and an unpaired final observation changes an odd-length average by $O_\phi(N^{-1})$. $\square$

If, beyond [Assumption 8.1](#), the phase-coordinate analysis supplies the quantitative paired-mean estimate

$$|c_{n,i} - m_*(\phi)| = O((\log(n+1))^{-\beta\zeta}), \quad \zeta > 0, \quad (56)$$

so (53) becomes

$$\left\| \frac{1}{N} \sum_{r=1}^N \phi(\Phi_r) - m_*(\phi) \right\|_{L^2} = O\left(N^{-1/2} + (\log N)^{-\beta\zeta}\right). \quad (57)$$

For any prescribed $\beta > 0$, if the corresponding schedule satisfies [Assumption 8.1](#) and (56), the argument gives almost-everywhere convergence; the probabilistic step itself imposes no threshold $\beta > 1$. This is not an assertion that every logarithmic quadratic schedule satisfies those hypotheses. A threshold can arise from a weaker covariance estimate with a nonsummable drift error, but it is not caused by the parity mode itself.

9 What is, and is not, proved for nearby quadratic schedules

At the constant schedule $u_n \equiv u_c$, the phase swap (S1) and the common paired limit (S5) follow from the exact component interchange and mixing. If [Assumption 3.1](#) is realized on component Banach spaces stable under the multipliers in (S4), it also supplies (S2)–(S3) and the exponential covariance bound, so [theorem 8.2](#) applies. Independently, [proposition 6.2](#) gives the autonomous almost-everywhere Birkhoff limit without the component-gap assumption; its quantitative $O(N^{-1})$ variance estimate uses that assumption. For a nonconstant schedule in the quadratic family, parameter convergence by itself is insufficient. A concrete application of [theorems 7.2](#) and [8.2](#) requires all of the following.

- (1) The actual parameters must lie in a common matched inducing or tower set with uniform expansion, distortion, and tail data. The existence of a positive-measure Benedicks–Carleson-type set accumulating at u_c does not imply that a prescribed sequence lies in it.
- (2) Mixed blocks $P_b P_a$, not only autonomous squares P_u^2 , must possess a uniformly isolated rank-two cluster. Keller–Liverani stability [6] is a natural tool once the strong–weak estimates and branch matching have actually been established.
- (3) Arbitrary products of the complementary block operators must satisfy (33) or (45). Separate spectral gaps for each frozen block do not imply this cocycle estimate.
- (4) A fixed or covariantly moving phase label must be constructed. For $u \neq u_c$, the exact sets C, H in (6) need not be swapped, and their boundaries may move or cease to define two invariant phases.

- (5) The internal 2×2 bundle cocycle must be analyzed. If a covariant parity line exists, its scalar multiplier may persist or decay; in the general case the full matrix product controls the outcome.

These are substantive verification tasks. They are the hypotheses not captured by a one-mode raw estimate. The present results identify the required operator architecture and prove the conclusions once that architecture is available; they do not claim that every logarithmically cooled quadratic schedule satisfies it.

10 Conclusion

The first quadratic band-merging parameter is a period-two, not a mixing, raw dynamical system. Unconditionally, its Koopman dynamics contain the parity eigenvalue -1 , producing an atom at frequency $1/2$ for every observable with $a_\phi \neq 0$, and in particular $\sigma_s = \delta_{1/2}$. Under the explicit two-step component spectral-gap hypothesis, the Perron–Frobenius decomposition contains exactly the two peripheral modes $1, -1$; after removing them, the remaining correlations decay exponentially and give an analytic absolutely continuous spectral density. For each fixed bounded real observable, finite orbit periodograms converge weakly to the observable spectral measure without requiring that gap.

The same observation clarifies the sequential logic. Consecutive iterates must be paired, or equivalently the cocycle must be studied through $Q_n = P_{2n}P_{2n-1}$. The two-step complement can mix while the rank-two projector bundle retains phase information. Under uniform block memory loss and the common paired-mean hypothesis, paired Birkhoff averages converge in L^2 and almost everywhere; under the moving-projector hypotheses, the adiabatic lag is controlled by adjacent parameter increments, while comparison with the limiting statistics is controlled by the larger parameter offset.

The main theoretical boundary is now explicit. The exact geometry, parity eigenmode, ergodicity, Birkhoff limit, and pathwise periodogram limit at u_c for each fixed bounded real observable are unconditional. Exponential operator-norm remainders require the stated component gap. A non-autonomous quadratic application further requires a verified matched tower, a rank-two block bundle, sequential complement contraction, and an analysis of the internal two-dimensional cocycle. Until those ingredients are supplied, a numerical cooling schedule cannot be promoted to an operator-spectral theorem merely because its parameters approach u_c .

Data and code availability

All results are analytic and no numerical data are used. The manuscript source and verified PDF are available at https://github.com/maris205/prime_dynamics_theory/tree/main/papers/RH-3-parity-resolved-band-merging-spectrum.

Acknowledgements

The author thanks colleagues whose comments motivated a separate treatment of the period-two mode. AI-assisted tools were used for auxiliary typesetting and proof auditing; all mathematical statements and conclusions are the responsibility of the author.

Disclosure statement

The author reports no competing interests.

References

- [1] Romain Aimino, Huyi Hu, Matthew Nicol, Andrei Török, and Sandro Vaienti. Polynomial loss of memory for maps of the interval with a neutral fixed point. *Discrete and Continuous Dynamical Systems*, 35(3):793–806, 2015. doi: 10.3934/dcds.2015.35.793.
- [2] Viviane Baladi. *Positive Transfer Operators and Decay of Correlations*, volume 16 of *Advanced Series in Nonlinear Dynamics*. World Scientific, Singapore, 2000. doi: 10.1142/9789812813633.
- [3] Jean-Pierre Conze and Albert Raugi. Limit theorems for sequential expanding dynamical systems on $[0, 1]$. In *Ergodic Theory and Related Fields*, volume 430 of *Contemporary Mathematics*, pages 89–121. American Mathematical Society, Providence, RI, 2007. doi: 10.1090/conm/430/08253.
- [4] I. P. Cornfeld, S. V. Fomin, and Ya. G. Sinai. *Ergodic Theory*, volume 245 of *Grundlehren der mathematischen Wissenschaften*. Springer, New York, 1982. doi: 10.1007/978-1-4615-6927-5.
- [5] Welington de Melo and Sebastian van Strien. *One-Dimensional Dynamics*, volume 25 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer, Berlin, 1993. doi: 10.1007/978-3-642-78043-1.
- [6] Gerhard Keller and Carlangelo Liverani. Stability of the spectrum for transfer operators. *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze*, 28(1):141–152, 1999. URL https://www.numdam.org/item/ASNSP_1999_4_28_1_141_0/.
- [7] Douglas Lind and Brian Marcus. *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press, Cambridge, 1995. doi: 10.1017/CBO9780511626302.
- [8] Michał Misiurewicz. Absolutely continuous measures for certain maps of an interval. *Publications Mathématiques de l’IHÉS*, 53:17–51, 1981. doi: 10.1007/BF02698687.
- [9] Liang Wang. The emergence of prime distribution from low-dimensional deterministic chaos. *Research in Mathematics*, 13(1):2684334, 2026. doi: 10.1080/27684830.2026.2684334.
- [10] Liang Wang. An exact prime kneading orbit: Sparse admissible repair and fourier-spectral stability of sieve words. Preprint, 2026. URL https://github.com/maris205/prime_dynamics_theory/tree/main/papers/RH-2-exact-prime-kneading-spectral-stability.
- [11] Liang Wang. From sieve words to kneading coordinates: A rigorous reformulation of the one-dimensional prime-sieve correspondence. Preprint, 2026. URL https://github.com/maris205/prime_dynamics_theory/tree/main/papers/RH-1-rigorous-sieve-kneading-reformulation.
- [12] Liang Wang. A sequential Birkhoff theorem for slow logarithmic drift in non-uniformly expanding unimodal maps. Manuscript, 2026.
- [13] Lai-Sang Young. Decay of correlations for certain quadratic maps. *Communications in Mathematical Physics*, 146(1):123–138, 1992. doi: 10.1007/BF02099209.