

Extended Deterministic Numerator Anchor Atlas

Bin Wang

July 2026

Abstract

The analytic-tail theorem of RH-252 is exact but does not identify the finite target coefficients beyond the order-12 table of RH-243. We evaluate the same deterministic periodic-point formula through order 28. This adds 16 orders and enumerates 32767 physical fixed points at the largest order. The new order block contributes 0.0021942543 to the unit-disk logarithmic norm, and its descriptive log-linear root rate is 0.7009986. These are finite reproducible facts. They do not prove a uniform trace envelope, a cloud coefficient bridge, or any statement about Riemann zeros.

1 Exact target dictionary

Let P_n denote the deterministic flat periodic trace from the quadratic band-merging map. The RH-243 dictionary is

$$a_n = r_H^{-n} \left[P_n - 1 - (-1)^n + 2\mathbf{1}_{2|n} \lambda^{-n/2} \right], \quad r_H = 0.85, \quad (1)$$

with $\lambda = 1.6785735104283177 \dots$ [2]. The periodic point and multiplier formulas used to evaluate P_n are the exact finite formulas archived with the Collet–Eckmann completion [1].

Proposition 1.1 (Finite extension). *For every integer $2 \leq n \leq 28$, equation (1) defines an exact finite deterministic target value. The order-28 computation enumerates 32767 distinct physical fixed points before summing their flat weights.*

Proof. The inverse-branch construction in RH-11 enumerates all cyclic Markov words, removes the duplicated shared endpoint, and verifies the physical fixed-point count. Applying the displayed finite sum for each order and then the explicit algebraic correction in (1) gives the claim. $\square$

2 The new order block

The following values are representative rows from the 16 new orders. The last column is the Hardy-scaled target, not a noisy trace.

n	P_n	unscaled g_n	a_n
13	0.00118921	0.00118921	0.00983586
16	1.96851809	0.000250789	0.00337758
20	1.98876937	0.0000316550	0.000816703
24	1.99600689	0.00000399130	0.000197269
28	1.99858189	0.000000502810	0.0000476073

For the unit disk and the logarithmic norm

$$J_I(1) := \sum_{n \in I} \frac{|a_n|}{n}, \quad (2)$$

the archived computation gives

$$J_{2:12}(1) = 0.49450543569144195, \quad (3)$$

$$J_{13:28}(1) = 0.0021942543215719553, \quad (4)$$

$$J_{2:28}(1) = 0.496699690013014. \quad (5)$$

At radius 0.8, the new block contributes only $8.7150803572 \times 10^{-5}$.

3 Finite slope diagnostic and its boundary

Fitting $\log |a_n|$ linearly against n on the new block gives

$$\hat{q}_{13:28} = 0.7009986349256669, \quad \hat{q}_{\text{odd}} = 0.7009206511866494, \quad \hat{q}_{\text{even}} = 0.7011096165507584. \quad (6)$$

The agreement between parity subsequences is a useful finite consistency check with the RH-252 radius budget [3]. It is not a proof of $|a_n| \leq Cq^n$ for all n : a finite fit has no control over orders 29 and beyond, nor over the constant needed in the Cauchy tail theorem.

Remark 3.1 (Scope). The atlas is deterministic and finite. It does not select a noisy spectral cloud, certify a quotient block, or identify the coefficients of a moving operator. The target is an anchor, not an observed eigenvalue list.

4 Route decision

RH-252 provides an all-order analytic existence statement for the target tail, while this paper supplies a longer finite target table. The next genuinely new input must come from the spectral side: an expanded resolved candidate window or an invariant quotient selector. Reusing the frozen RH-248 shell window would not be progress. Gates A–E remain false/open; no Hilbert–Polya operator, zeta-divisor equality, Riemann-zero identification, or RH implication is asserted.

References

- [1] Bin Wang. Collet–eckmann flat-trace completion. *Preprint*, 2026. RH-11 preprint.
- [2] Bin Wang. A deterministic numerator coefficient-anchor dictionary. *Preprint*, 2026. RH-243 preprint.
- [3] Bin Wang. Deterministic-numerator analytic-tail certificate. *Preprint*, 2026. RH-252 preprint.