

# Taeniasis and Cysticercosis:

a substantial public health and economic challenge

Marisa Wong & Angela Shen

# Taeniasis? Cysticercosis?

## Taeniasis

Refers to intestinal infection with adult tapeworms

3 parasite species cause taeniasis

- ***Taenia solium* (pork tapeworm)**
- *Taenia saginata* (beef tapeworm)
- *Taenia asiatica* (Asian tapeworm)

Humans are the only definitive hosts for these adult *Taenia* tapeworms

Only *T. solium* causes major health problems

## Cysticercosis

Infection caused by the larva form (cysticerci) of *T. solium*

Humans and pigs can act as intermediary hosts: harbor the larval stage

Human cysticercosis can result in devastating health effects

- Nodules can develop across the body
- **Neurocysticercosis (NCC):**  
**cysticerci lodge/develop in the brain or CNS**



# Life Cycle

Swine are the usual intermediate host for *T. solium*

- Humans, the usual definitive host, can serve as accidental intermediate hosts following ingestion of infectious eggs


**Cysticercosis** acquired only from the **fecal-oral route (ingestion of eggs)**, not via the **ingestion of cysticerci in undercooked pork**, which is associated with intestinal **taeniasis**

Tapeworm carrier can infect him- or herself with tapeworm eggs: autoinfection



*Taenia solium*

4DPDx




# Symptoms & Health Effects

Taeniasis is usually characterized by **mild and non-specific** symptoms:

- Abdominal pain
- Nausea
- Diarrhoea or constipation

Symptoms present when tapeworms become fully developed in the intestine, ~ 8 weeks after ingestion of meat containing cysticerci

Continue until the tapeworm dies, ~2-3 years without intervention



Symptoms of Cysticercosis depend on the location of cysticerci

Extraneural cysticercosis

- No major symptoms
- Visible, palpable nodules under the skin, which can become tender

Ophthalmic cysticercosis

- Rarest form, cysts in retro-ocular space can cause proptosis or vision loss

Neurocysticercosis symptoms depend on the number, size, and topography of cysts

- **Epileptic seizures (50-80% of patients)**
- Chronic headaches, Hydrocephalus, Meningitis, Blindness

**Most frequent preventable cause of epilepsy** worldwide

- 30% of all epilepsy cases in endemic settings
- Specific communities 70% of all epilepsy cases

# Global Burden of Disease



## Acquisition

- *T. saginata* or *T. asiatica*: consume undercooked infected beef meat or pig liver [Taeniasis]
- ***T. solium***: consume raw or undercooked, infected pork [Taeniasis]
  - *T. solium* eggs: consume contaminated food or water (fecal-oral route) [Cysticercosis]

Disproportionately found in **rural areas of developing countries**: poor sanitation, pigs roam freely and eat human feces

- *Subsistence farmers*


Rare among persons who live in countries where pigs are not raised and where pigs do not have contact with human feces.

*T. solium*: ~ 2.8 million DALYs in 2015

NCC: ~2.56–8.30 million people

- Massive economic burden
  - Treatment and productivity losses consume 54% of the annual minimum wage
  - Symptoms (seizure and epilepsy) cause two thirds of employees to lose their jobs

Endemicity of *Taenia solium*, 2022



The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement. © WHO 2022. All rights reserved.

Data Source: World Health Organization  
Map Production: Control of Neglected  
Tropical Diseases (NTD)  
World Health Organization



# Treatment

## Stop Parasite Transmission Cycle & Treat Mild Symptoms

- Single dose of 10 mg/kg praziquantel, single dose of 1-2g niclosamide, or 400 mg of albendazole for 3 days
  - Preventive Chemotherapy for *T. solium*
  - Individual basis

## Neurocysticercosis

- Specialized treatment with high doses of praziquantel or albendazole for long durations
- Dosage and duration of treatment varies based on number, size, location, development stage of cysts, and severity of symptoms

# Prevention

160 F




145 F



# Control



## Core “rapid impact” Interventions



## Supporting measures



## Societal Changes





# Impact of Climate Change



## Humidity

High humidity improving  
egg survival



## Temperature

Moderate temperature  
(5-25 C) favors egg survival



## Rural Animal Husbandry

Increased zoonotic  
potential & cross-species  
jump of the parasite

# References



- CDC - Cysticercosis - Biology
- WHO - Taeniasis and Cysticercosis Fact Sheet
- CDC Parasite - Taeniasis
- Jansen, F. et al. (2021). The survival and dispersal of Taenia eggs in the environment: what are the implications for transmission? A systematic review
- O'Hara, K. et al. (2019). Evaluating the association between climatic events and sheep condemnations in the United States using cluster analysis and spatio-temporal modeling.

# Thanks

## Any Questions?

[angela.shen2@emory.edu](mailto:angela.shen2@emory.edu)

[marisa.wong@emory.edu](mailto:marisa.wong@emory.edu)

