


Junior problems

- J91. The squares in the figure below are labeled 1 through 16 such that the sum of the numbers in each row and each column is the same. The positions of 1, 5, and 13 are given.



Prove that there is only one possibility for the number in the darkened square and find this number.

Proposed by Ivan Borsenco, Massachusetts Institute of Technology, USA

- J92. Find all primes q_1, q_2, \dots, q_6 such that $q_1^2 = q_2^2 + \dots + q_6^2$.

Proposed by Titu Andreescu, University of Texas at Dallas, USA

- J93. Let a and b be positive real numbers. Prove that

$$\frac{a^6 + b^6}{a^4 + b^4} \geq \frac{a^4 + b^4}{a^3 + b^3} \cdot \frac{a^2 + b^2}{a + b}.$$

Proposed by Arkady Alt, San Jose, California, USA

- J94. Prove that the equation $x^3 + y^3 + z^3 + w^3 = 2008$ has infinitely many solutions in integers.

Proposed by Titu Andreescu, University of Texas at Dallas, USA

- J95. Let ABC be a triangle and let I_a, I_b, I_c be its excenters. Denote by O_a, O_b, O_c the circumcenters of triangles I_aBC, I_bAC, I_cAB . Prove that the area of triangle $I_aI_bI_c$ is twice the area of hexagon $O_aCO_bAO_cB$.

Proposed by Mehmet Sahin, Ankara, Turkey

- J96. Let n be an integer. Find all integers m such that $a^m + b^m \geq a^n + b^n$ for all positive real numbers a and b with $a + b = 2$.

Proposed by Oleg Mushkarov, Bulgarian Academy of Sciences, Sofia, Bulgaria

Senior problems

- S91. Find all triples (n, k, p) , where n and k are positive integers and p is a prime, satisfying the equation

$$n^5 + n^4 + 1 = p^k.$$

Proposed by Titu Andreescu, University of Texas at Dallas, USA

- S92. Let ABC be a triangle with altitudes BE and CF and let M be a point on its circumcircle. Denote by P the intersection of MB and CF and by Q the intersection of MC and BE . Prove that EF bisects the segment PQ .

Proposed by Son Ta Hong, Ha Noi University, Vietnam

- S93. Let n be an integer greater than 1 and let x_1, x_2, \dots, x_n be nonnegative real numbers whose sum is $\sqrt{2}$. Determine the maximum, as a function of n , of

$$\frac{x_1^2}{1+x_1^2} + \frac{x_2^2}{1+x_2^2} + \cdots + \frac{x_n^2}{1+x_n^2}.$$

Proposed by Alex Anderson, Washington University in St. Louis, USA

- S94. Consider a quadrilateral that is inscribed in a circle and circumscribed about a circle. Prove that the product of its diagonals is a constant.

Proposed by Ivan Borsenco, MIT, USA

- S95. There are 16 bad boys in a neighborhood. During one year 69 quarrels have occurred among the members of the neighborhood (each quarrel involves exactly two persons). At the end of the year a local wrestling club wants to organize a match between two teams of three people. The boys will fight each other if and only if each member of one team has quarreled with each member of the other team. Prove that the club can always organize such a fight.

Proposed by Iurie Boreico, Harvard University and Ivan Borsenco, MIT, USA

- S96. Let n be an integer greater than 2. Prove that $\binom{n-1}{k} \equiv (-1)^k \pmod{n}$ for each $k = 1, 2, \dots, n-1$, if and only if n is a prime.

Proposed by Dorin Andrica, "Babes-Bolyai" University, and Mihai Piticari, "Dragos Voda" National College, Romania

Undergraduate problems

U91. Prove that there are no polynomials $P, Q \in \mathbb{R}[x]$ such that

$$\int_0^{\log n} \frac{P(x)}{Q(x)} dx = \frac{n}{\pi(n)},$$

for all $n \geq 1$, where $\pi(n)$ is the prime counting function.

Proposed by Cezar Lupu, University of Bucharest, Romania

U92. Find the maximum value of $F(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \min \left\{ \frac{\|\mathbf{y} - \mathbf{z}\|}{\|\mathbf{x}\|}, \frac{\|\mathbf{z} - \mathbf{x}\|}{\|\mathbf{y}\|}, \frac{\|\mathbf{x} - \mathbf{y}\|}{\|\mathbf{z}\|} \right\}$

where $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are arbitrary nonzero vectors in $\mathbb{R}^n, n \geq 2$.

Proposed by Arkady Alt, San Jose, California, USA

U93. Let $x_0 \in (0, 1]$ and $x_{n+1} = x_n - \arcsin(\sin^3 x_n), n \geq 0$. Evaluate $\lim_{n \rightarrow \infty} \sqrt{n}x_n$.

Proposed by Titu Andreescu, University of Texas at Dallas, USA

U94. Let Δ be the plane domain consisting of all interior and boundary points of a rectangle $ABCD$, whose sides have lengths a and b . Define $f : \Delta \rightarrow \mathbb{R}$, $f(P) = PA + PB + PC + PD$. Find the range of f .

Proposed by Mircea Becheanu, University of Bucharest, Romania

U95. Find all monic polynomials P and Q , with real coefficients, such that

$$P(1) + P(2) + \cdots + P(n) = Q(1 + 2 + 3 + \cdots + n),$$

for all $n \geq 1$.

Proposed by Ovidiu Furdui, Cluj-Napoca, Romania

U96. Let $f : (0, \infty) \rightarrow [0, \infty)$ be a bounded function. Prove that if

$$\lim_{x \rightarrow 0} \left(f(x) - \frac{1}{2} \sqrt{f\left(\frac{x}{2}\right)} \right) = 0 \quad \text{and} \quad \lim_{x \rightarrow 0} (f(x) - 2f(2x)^2) = 0,$$

then $\lim_{x \rightarrow 0} f(x) = 0$.

*Proposed by Dorin Andrica, "Babes-Bolyai" University, Romania and
Mihai Piticari, "Dragos Voda" National College, Romania*

Olympiad problems

O91. Let ABC be an acute triangle. Prove that

$$\tan A + \tan B + \tan C \geq \frac{s}{r},$$

where s and r are semiperimeter and inradius of triangle ABC , respectively.

Proposed by Mircea Becheanu, University of Bucharest, Romania

O92. Let n be a positive integer. Prove that

- a) there are infinitely many triples (a, b, c) of distinct integers such that $\min(a, b, c) \geq n$ and $abc + 1$ divides one of the numbers $(a - b)^2, (b - c)^2, (c - a)^2$.
- b) there is no triple (a, b, c) of distinct positive integers such that $abc + 1$ divides more than one of the numbers $(a - b)^2, (b - c)^2, (c - a)^2$.

Proposed by Titu Andreescu, University of Texas at Dallas, USA

O93. Let k be a positive integer. Find all functions $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $f(x) + f(y)$ divides $x^k + y^k$ for all $x, y \in \mathbb{N}$.

Proposed by Nguyen Tho Tung, Hanoi University of Education, Vietnam

O94. Let ω be a circle with center O and let A be a fixed point outside ω . Choose points B and C on ω , with $AB \neq AC$, such that AO is a symmedian, but not a median, in triangle ABC . Prove that the circumcircle of triangle ABC passes through a second fixed point.

Proposed by Alex Anderson, Washington University in St. Louis, USA

O95. Prove that there is a sequence x_1, x_2, \dots of integers such that

- For each $n \in \mathbb{Z}$ there exists i such that $x_i = n$.

- $\prod_{d|n} d^{\frac{n}{d}} = \sum_{i=1}^n x_i.$

Proposed by Juan Ignacio Restrepo, Universidad de Los Andes, Colombia

O96. Let p and q be primes, $q \geq p$. Prove that pq divides $\binom{p+q}{p} - \binom{q}{p} - 1$.

Proposed by Dorin Andrica, "Babes-Bolyai" University, Romania