

Negative binomial regression


Mark Andrews

Psychology Department, Nottingham Trent University

✉ `mark.andrews@ntu.ac.uk`

The Poisson Distribution

- The mean of a Poisson distribution is equal to its rate parameter λ .
- Its variance is also equal to λ .



As λ increases, so too does the variance.

Means and variances in a Poisson distribution:

- ▶ In a Poisson distribution, the variance of a sample should be approximately the same as the mean of a sample.
- ▶ Example 1:

```
x <- rpois(25, lambda = 5)
c(mean(x), var(x), var(x)/mean(x))
```

```
## [1] 5.4400000 5.4233333 0.9969363
```

- ▶ Example 2:

```
x <- rpois(25, lambda = 5)
c(mean(x), var(x), var(x)/mean(x))
```


```
## [1] 5.4000000 6.416667 1.188272
```

Overdispersion

- ▶ If the variance of a sample is greater than would be expected according to a given theoretical model, then we say the data is *overdispersed*.
- ▶ In count data, if the variance of a sample is much greater than its mean, we say it is overdispersed.
- ▶ Using a Poisson distribution in this situation, this is an example of model mis-specification.
- ▶ It will also usually underestimate the standard errors in the Poisson model.

Overdispersion

- In the bioChemists data set, we have counts of the number of articles published by PhD students in the last three years (publications):



```
var(publications)/mean(publications)
```

```
## [1] 2.191358
```

Overdispersion

- This leads standard errors to be *underestimated* if we use a Poisson model:

```
M <- glm(publications ~ 1, family=poisson)
summary(M)$coefficients
```

##	Estimate	Std. Error	z value	Pr(> z)
## (Intercept)	0.5264408	0.02540804	20.71945	2.312911e-95

Fixing overdispersion using a Quasi-poisson model

- ▶ A *quasi* Poisson model allows us to correct over-dispersion

```
M <- glm(publications ~ 1, family=quasipoisson)
summary(M)$coefficients
```

```
##              Estimate Std. Error  t value    Pr(>|t|)
## (Intercept) 0.5264408 0.03761239 13.99647 1.791686e-40
```


- ▶ It does so by calculating an overdispersion parameter (roughly, the ratio of the variance to the mean) and multiplying the standard error by its square root.
- ▶ In this example, the overdispersion parameter is 2.1913892 and so its square root is 1.4803341.
- ▶ Alternatively, a *negative binomial regression* is an alternative to Poisson regression that can be used with overdispersed count data.

Negative binomial distribution

- ▶ A negative binomial distribution is a distribution over non-negative integers.
- ▶ To understand the negative binomial distribution, we start with the binomial distribution:
- ▶ If, for example, we have a coin whose probability of coming up heads is θ , then the number of Heads in a sequence of n flips will follow a binomial distribution.
- ▶ In this example, an outcome of Heads can be termed a *success*.


Negative binomial distribution

- Here is a binomial distribution where $n = 25$ and $\theta = 0.75$.




Negative binomial distribution

- ▶ A *negative binomial* distribution gives the probability distribution over the number of *successes* (e.g. Heads) before r *failures* (e.g. r Tails).
- ▶ For example, here we have the number of Heads (*successes*) that occur before we observe $r = 2$ Tails (*failures*), when the probability of Heads is $\theta = 0.75$:



Negative binomial distribution

- Here, we have the number of Heads (*successes*) that occur before we observe $r = 3$ Tails (*failures*), when the probability of Heads is $\theta = 0.5$:



Negative binomial distribution

- The probability mass function for the negative binomial distribution is:

$$P(x = k|r, \theta) = \binom{r+k-1}{k} \theta^r (1-\theta)^k$$

or more generally

$$P(x = k|r, \theta) = \frac{\Gamma(r+k)}{\Gamma(r)k!} \theta^r (1-\theta)^k,$$

where $\Gamma()$ is a Gamma function ($\Gamma(n) = (n-1)!$).

- In R, for any k , r , and θ , we can calculate $P(x = k|r, \theta)$ using `dnbinom`, e.g. $P(x = k = 2|r = 3, \theta = 0.75)$ is

```
dnbinom(2, 3, 0.75)
```

```
## [1] 0.1582031
```

Negative binomial distribution


- In the negative binomial distribution, the mean is

$$\mu = \frac{\theta}{1 - \theta} \times r,$$

and we can generally parameterize the distribution by μ and r .

Why use negative binomial distribution?

- A negative binomial distribution is equivalent as weighted sum of Poissons.



- So it is appropriate to use when the data can be seen as arising from a mixture of Poisson distributions, each with different means.

Negative binomial regression

- In negative binomial regression, we have observed counts $y_1, y_2 \dots y_n$, and some predictor variables $x_1, x_2 \dots x_n$, and we assume that

$$y_i \sim \text{NegBinomial}(\mu_i, r),$$

where $\text{NegBinomial}(\mu_i, r)$ is a negative binomial with mean μ_i and a dispersion parameter r , and then

$$\log(\mu_i) = \beta_0 + \beta x_i.$$