

Tutorial: Dependencies

1. Consider the relational schema $R = \{A, B, C, D, E\}$ with the following set of functional dependencies.

$$\Sigma = \{\{A, B\} \rightarrow \{C\}, \{D\} \rightarrow \{D, B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}, \{A, B, D\} \rightarrow \{A, B, C, D\}\}$$

- (a) Compute all the closures of the the sets of attributes that are not equal to themselves, are not super-keys or are candidate keys. What information is not essential and could be removed.

Solution:

$$\{A, B\}^+ = \{A, B, C, D, E\}$$

$$\{A, D\}^+ = \{A, B, C, D, E\}$$

$$\{A, E\}^+ = \{A, B, C, D, E\}$$

$$\{B\}^+ = \{B, D, E\}$$

$$\{D\}^+ = \{B, D, E\}$$

$$\{E\}^+ = \{B, D, E\}$$

$$\{B, D\}^+ = \{B, D, E\}$$

$$\{B, E\}^+ = \{B, D, E\}$$

$$\{D, E\}^+ = \{B, D, E\}$$

$$\{B, C\}^+ = \{B, C, D, E\}$$

$$\{D, C\}^+ = \{B, C, D, E\}$$

$$\{E, C\}^+ = \{B, C, D, E\}$$

$$\{B, D, C\}^+ = \{B, C, D, E\}$$

$$\{B, E, C\}^+ = \{B, C, D, E\}$$

$$\{D, E, C\}^+ = \{B, C, D, E\}$$

We see that the candidate keys are $\{A, B\}$, $\{A, D\}$ and $\{A, E\}$.

Remark: You can even remove from the right-hand side the members of the set on the left-hand side. It also keeps all the essential information (how to prove it?).

$$\{A, B\}^+ = \{C, D, E\}$$

$$\{A, D\}^+ = \{B, C, E\}$$

$$\{A, E\}^+ = \{B, C, D\}$$

$$\{B\}^+ = \{D, E\}$$

$$\{D\}^+ = \{B, E\}$$

$$\{E\}^+ = \{B, D\}$$

$$\{B, D\}^+ = \{E\}$$

$$\{B, E\}^+ = \{D\}$$

$$\{D, E\}^+ = \{B\}$$

$$\{B, C\}^+ = \{D, E\}$$

$$\{D, C\}^+ = \{B, E\}$$

$$\{E, C\}^+ = \{B, D\}$$

$$\{B, D, C\}^+ = \{E\}$$

$$\{B, E, C\}^+ = \{D\}$$

$$\{D, E, C\}^+ = \{B\}$$

Remark: You can even remove an equality if the left-hand side is a superset of another equality left-hand-side and its right-hand side is a subset of the right-hand side. It also keeps all the essential information (how to prove it?).

$$\{A, B\}^+ = \{C, D, E\}$$

$$\{A, D\}^+ = \{B, C, E\}$$

$$\{A, E\}^+ = \{B, C, D\}$$

$$\{B\}^+ = \{D, E\}$$

$$\{D\}^+ = \{B, E\}$$

$$\{E\}^+ = \{B, D\}$$

- (b) What are the candidate keys of R with Σ ?

Solution: The candidate keys are $\{A, B\}$, $\{A, D\}$ and $\{A, E\}$ (see question 1).

- (c) Find a minimal cover of R with Σ that can be reached from Σ using the algorithm from the lecture.

Solution:

Step 1: We construct Σ' , which is equivalent to Σ , by minimizing the right-hand side of every functional dependency in Σ .

$$\Sigma' = \{\{A, B\} \rightarrow \{C\}, \underline{\{D\} \rightarrow \{D\}}, \underline{\{D\} \rightarrow \{B\}}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}, \\ \{A, B, D\} \rightarrow \{A\}, \{A, B, D\} \rightarrow \{B\}, \{A, B, D\} \rightarrow \{C\}, \{A, B, D\} \rightarrow \{D\}\}$$

Step 2: We construct Σ'' , which is equivalent to Σ' (and to Σ), by minimizing the left-hand side of every functional dependency in Σ' . We may have the choice between several possible simplifications. We only need to choose one.

$$\Sigma'' = \{\{A, B\} \rightarrow \{C\}, \{D\} \rightarrow \{D\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}, \\ \underline{\{A\} \rightarrow \{A\}}, \underline{\{B\} \rightarrow \{B\}}, \underline{\{A, B\} \rightarrow \{C\}}, \underline{\{D\} \rightarrow \{D\}}\}$$

or

$$\Sigma'' = \{\{A, B\} \rightarrow \{C\}, \{D\} \rightarrow \{D\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}, \\ \underline{\{A\} \rightarrow \{A\}}, \underline{\{B\} \rightarrow \{B\}}, \underline{\{A, D\} \rightarrow \{C\}}, \underline{\{D\} \rightarrow \{D\}}\}$$

Notice that we can keep all the candidates with a minimal left-hand side. Namely, it suffices, in the example to keep:

$$\Sigma'' = \{\underline{\{A, B\} \rightarrow \{C\}}, \{D\} \rightarrow \{D\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}, \\ \{A\} \rightarrow \{A\}, \{B\} \rightarrow \{B\}, \underline{\{A, D\} \rightarrow \{C\}}, \{D\} \rightarrow \{D\}\}$$

Step 3: We construct Σ''' , which is equivalent to Σ'' (and to Σ), by minimizing the whole set Σ'' . We may have the choice between several possible simplifications. The choice depends on the order in which we consider the functional dependencies.

$$\Sigma''' = \{\{A, B\} \rightarrow \{C\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}\}$$

or

$$\Sigma''' = \{\{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}, \{A, D\} \rightarrow \{C\}\}$$

(d) Find all the minimal covers of R with Σ .

Solution:

$$\Sigma''' = \{\{A, B\} \rightarrow \{C\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}\}$$

$$\{\{A, D\} \rightarrow \{C\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}\}$$

$$\{\{A, E\} \rightarrow \{C\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{D\}\}$$

$$\{\{A, B\} \rightarrow \{C\}, \{D\} \rightarrow \{E\}, \{E\} \rightarrow \{B\}, \{B\} \rightarrow \{D\}\}$$

$$\{\{A, D\} \rightarrow \{C\}, \{D\} \rightarrow \{E\}, \{E\} \rightarrow \{B\}, \{B\} \rightarrow \{D\}\}$$

$$\{\{A, E\} \rightarrow \{C\}, \{D\} \rightarrow \{E\}, \{E\} \rightarrow \{B\}, \{B\} \rightarrow \{D\}\}$$

$$\{\{A, D\} \rightarrow \{C\}, \{B\} \rightarrow \{D\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{B\}\}$$

$$\{\{A, D\} \rightarrow \{C\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{D\}, \{E\} \rightarrow \{D\}, \{D\} \rightarrow \{E\}\}$$

$$\{\{A, D\} \rightarrow \{C\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{B\}, \{E\} \rightarrow \{D\}, \{D\} \rightarrow \{E\}\}$$

$$\{\{A, E\} \rightarrow \{C\}, \{B\} \rightarrow \{D\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{B\}\}$$

$$\{\{A, E\} \rightarrow \{C\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{D\}, \{E\} \rightarrow \{D\}, \{D\} \rightarrow \{E\}\}$$

$$\{\{A, E\} \rightarrow \{C\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{B\}, \{E\} \rightarrow \{D\}, \{D\} \rightarrow \{E\}\}$$

$$\{\{A, B\} \rightarrow \{C\}, \{B\} \rightarrow \{D\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{B\}\}$$

$$\{\{A, B\} \rightarrow \{C\}, \{D\} \rightarrow \{B\}, \{B\} \rightarrow \{D\}, \{E\} \rightarrow \{D\}, \{D\} \rightarrow \{E\}\}$$

$$\{\{A, B\} \rightarrow \{C\}, \{B\} \rightarrow \{E\}, \{E\} \rightarrow \{B\}, \{E\} \rightarrow \{D\}, \{D\} \rightarrow \{E\}\}$$

(e) Prove, using the three Armstrong axioms, that the following set of functional dependencies is equivalent to Σ .

$$\Sigma''' = \{\{A, B\} \rightarrow \{C, D, E\}, \{A, D\} \rightarrow \{B, C, E\}, \{A, E\} \rightarrow \{B, C, D\}, \\ \{B\} \rightarrow \{D, E\}, \{D\} \rightarrow \{B, E\}, \{E\} \rightarrow \{B, D\}\}$$

Solution: We prove that every functional dependency in one of Σ and Σ''' can be obtained from those in the other set.

For example $\{B\} \rightarrow \{D, E\}$ can be obtained from Σ .

1. We know that $\{B\} \rightarrow \{E\}$.
2. We know that $\{E\} \rightarrow \{D\}$.
3. Therefore $\{E\} \rightarrow \{D, E\}$ by Augmentation of (2) with $\{E\}$.
4. Therefore $\{B\} \rightarrow \{D, E\}$, by Transitivity of (1) and (3).

Q.E.D.

Post the other answers on the Luminus Forum (type in Latex and post the PDF).