

30 — LUCY Threshold Geometry and Relational Closure

Professional reading edition of the evolving public MKUFT canon

Mark Charles McLaughlin

ORCID: [0009-0005-7736-1511](https://orcid.org/0009-0005-7736-1511)

Scientific affiliation	Independent Researcher
Object status	Research module reading edition — not a standalone Zenodo publication
Public source	Open canonical public source
Source commit	26e06b6fea52e6b052dbdd540e80d1d926b96b07
MKUFT backbone DOI	10.5281/zenodo.17780566

Status boundary. This PDF is a reader-facing rendering of an evolving GitHub research module. It is not a frozen standalone paper and does not receive a separate DOI. The cited Git commit fixes the exact source identity of this edition.

Rights. Copyright © 2026 Mark Charles McLaughlin. All rights reserved. Historical licences remain controlling only for their exact frozen objects.

A Layer-Before-Law Hypothesis for Local Relational Closure and Emergent Gravitational Behaviour

Author: Mark Charles McLaughlin

Framework: McLaughlin–Kairos Unified Field Theory (MKUFT)

MKUFT DOI: [10.5281/zenodo.17780566](https://doi.org/10.5281/zenodo.17780566)

Citation and provenance: [PROVENANCE_DOI_AND_ATTRIBUTION.md](#)

Status: public working hypothesis; not a completed physical derivation.

Abstract

General relativity already removes gravitation from the ordinary Newtonian category of a force, describing free motion through dynamical spacetime geometry. A substantial research literature also treats spacetime and gravitational dynamics as emergent, induced, thermodynamic, hydrodynamic, collective, or quantum-information phenomena. This paper therefore does not claim novelty for emergent gravity itself.

It develops one downstream MKUFT/LUCY physical branch. In the canonical framework, Local Unified Coherence Yield (LUCY) is the proposed crossing threshold at which an addressed relational or informational configuration becomes consequential at the next layer. For the physics-facing I→P case, the primary claim is the threshold at which a defined relational configuration changes physical transition weighting, admissibility, boundary condition, or effective state beyond the adequate physical baseline.

This paper studies the special case in which that crossing produces a stable interface and then a sustained local domain. In that branch, the useful sequence is **LUCY crossing threshold** → **LUCY-1 membrane** → **LUCY-2 sustained pocket**. The membrane and pocket are therefore downstream physical morphologies, not the definition or compulsory geometry of LUCY itself.

The central hypothesis is that some gravity-like observations could be effective outputs of a selected relational geometry rather than evidence of a separately generated wave, particle, or force.

A conventional Landau–Ginzburg and scalar–tensor style scaffold is used only as a toy realisation. Neutron-rich superheavy nuclei are treated as one candidate class of extreme configuration-sensitive matter because shell effects can stabilise nuclei that would otherwise fission rapidly and astrophysical r-process environments can access nuclei far beyond current terrestrial synthesis. No established result connects such nuclei to controllable gravitational behaviour. The proposed cross-layer coupling may be zero.

The paper states prior art, recovery requirements, discriminating signatures, evidence tiers, and falsifiers.

1. Scientific position

Two points are already established or well represented in existing research:

1. In general relativity, gravitation is not an ordinary force transmitted through a fixed background. Freely falling bodies follow geodesics of a dynamical spacetime geometry.
2. Emergent-gravity research includes induced gravity, thermodynamic derivations, analogue gravity, quantum-many-body space-time, condensate cosmology, matrix models, and scalar-tensor threshold effects.

The proposed MKUFT contribution in this paper is narrower:

A local physical system may admit more than one stable relational configuration. When declared control parameters cross a critical surface, a relational configuration may become physically consequential and the selected physical state may change discontinuously. In the boundary-forming branch studied here, matter trajectories, light propagation, clock relations, and inertial response would then change together because they are outputs of one effective geometry rather than several independently adjusted forces.

This becomes physical science only if it yields quantitative residuals beyond the strongest conventional electromagnetic, thermal, plasma, optical, acoustic, mechanical, scalar-tensor, and general-relativistic explanations.

2. Layer before law

MKUFT uses typed layer addresses,

$$S \rightarrow I \rightarrow P \rightarrow O \rightarrow S,$$

where S denotes substrate or source-potential, I information/relation/constraint/address, P physical expression, and O observer-positioned registration and record.

These are modelling addresses rather than extra spatial dimensions.

The Layer Before Law proposal is that a law observed at P may be the effective regularity of a deeper relational condition rather than the final primitive rule. The ordinary-physics limit remains compulsory.

In this paper, **relational geometry** means the complete constraint structure determining which physical paths, couplings, and transitions are admissible. It is not assumed to be a Lorentzian metric at the deepest level. An effective spacetime metric is one possible P-layer output.

The general LUCY crossing relation is **addressed relational constraint** $\rightarrow$ **thresholded physical consequence**.

The special branch developed here continues as **relational constraints** $\rightarrow$ **consequential I $\rightarrow$ P crossing** $\rightarrow$ **selected stable geometry** $\rightarrow$ **effective matter, light, and clock behaviour**.

3. LUCY crossing and boundary-forming branch

The canonical definition is owned by [LUCY Boundary Threshold Framework](#): LUCY is the threshold at which an addressed relation becomes consequential at the next layer. It is not intrinsically a bubble, circle, membrane, pocket, or other spatial shape.

This paper studies a particular P-layer morphology in which a stable interface follows the crossing:

- **LUCY** is the critical condition at which a candidate relational state becomes physically consequential under a declared coupling.
- **LUCY-1** is a stable interface or membrane when the physical implementation produces one.
- **LUCY-2** is a sustained finite regime when that interface can maintain a distinct local regime.

Coherence is used operationally for mutually consistent constraints that maintain a state against perturbation. **Relational closure** is the stronger condition in which the active constraints form a sufficiently complete loop for the state to persist, correct small deviations, or constrain downstream physical routing.

A physical implementation requires defined state variables and spaces, control parameters, a dynamical or variational rule, order parameter, critical surface, typed carrier into P, ordinary limit, energy and entropy accounting, observables, scale, sign, and failure conditions. Boundary conditions are additionally required for the membrane/pocket branch studied here.

Without these, LUCY remains architecture rather than physics.

4. Prior art and narrow novelty claim

Neighbouring work already includes Sakharov-style induced gravity, thermodynamic derivations of Einstein dynamics, analogue Lorentzian metrics in condensed matter, spacetime as a quantum many-body or condensate phase, scalar-tensor threshold phenomena and spontaneous scalarisation, and Landau–Ginzburg thresholds, metastability, domain walls, and hysteresis.

The MKUFT/LUCY contribution developed here is the conjunction of typed layer discipline, a thresholded I $\rightarrow$ P crossing condition, a boundary-forming three-stage physical branch, geometry as a selected state rather than an amplified force, a cross-channel common-boundary signature, and strongest-fair-null governance with preregistered promotion gates.

This conjunction is a research programme rather than a demonstrated mechanism.

5. Minimal toy scaffold

Let ψ be a candidate order parameter, g an effective metric, and λ declared control parameters. A conventional scalar–tensor style scaffold is

$$S_{\text{eff}}[g, \psi; \lambda] = \int d^4x \sqrt{-g} \left[\frac{M_0^2}{2} R - \frac{1}{2} K(\lambda) \nabla_\mu \psi \nabla^\mu \psi - V(\psi; \lambda) - \frac{1}{2} \xi \psi^2 R + \mathcal{L}_{\text{matter}}(g, \Phi) \right].$$

This is not claimed as an original or final MKUFT action. Related forms are standard in scalar-tensor theory, cosmology, and phase-transition modelling.

A toy potential is

$$V(\psi; \lambda) = \frac{1}{2} a(\lambda) \psi^2 + \frac{1}{4} b \psi^4 + \frac{1}{6} c \psi^6, \quad c > 0.$$

The LUCY critical surface may be defined by equality of relevant effective minima,

$$\Delta F(\lambda) = F[\psi_1, g_1; \lambda] - F[\psi_0, g_0; \lambda] = 0.$$

Below the surface the baseline state is preferred. Across it, a distinct physical state may become preferred. In this morphology-specific branch, LUCY-1 corresponds to a stable interface solution and LUCY-2 requires a finite domain whose bulk gain exceeds interface and dissipation costs.

For a schematic spherical pocket,

$$\Delta E(R) = 4\pi R^2 \sigma - \frac{4\pi}{3} R^3 \Delta \varepsilon + E_{\text{coupling}}(R).$$

In the simplest uncoupled case,

$$R_c = \frac{2\sigma}{\Delta \varepsilon}.$$

The spherical form is only a toy morphology for a bounded-domain branch. It is not the canonical shape of LUCY. Any geometry-coupling term must be dimensionally defined rather than inserted to force stability. The scaffold supplies threshold, boundary, pocket, hysteresis, and critical-radius concepts for this special case; it does not establish modified gravitation.

6. Effect regimes without two gravities

The paper distinguishes regimes by observable effect rather than naming two kinds of gravity.

A **localised high-gradient regime** is confined to nuclear, mesoscopic, material, or resonator scales where shell structure, pairing, deformation, collective modes, and boundary conditions may dominate stability.

An **extended effective-geometry regime** is a finite region in which neutral matter, light, and clocks exhibit a common change describable by one effective geometry.

The proposed relation is not a microscopic gravitational wave amplified into macroscopic gravity. It is one state-selection problem expressed differently at different scales.

A scale bridge requires an explicit coarse-graining map,

$$C_{\ell \rightarrow L} : \mathcal{X}_{\ell} \rightarrow \mathcal{X}_L,$$

that preserves energy, causality, and tested equivalence-principle limits. Until derived, the two regimes remain only a candidate relation.

7. Neutron-rich superheavy nuclei as a candidate seed class

Superheavy nuclear lifetime is highly configuration-sensitive. Liquid-drop trends favour fission, while shell corrections, pairing, and deformation can raise fission barriers and lengthen lifetimes. The location and lifetime of any island of enhanced stability remain model-dependent.

Terrestrial fusion currently reaches neutron-deficient superheavy isotopes. Astrophysical r-process environments can traverse nuclei far from laboratory access, but survival to long-lived trans-actinide species remains uncertain and depends on poorly known masses, separation energies, beta-decay rates, and fission barriers.

The scientifically permissible statement is:

Extreme astrophysical environments may create neutron-rich superheavy configurations unavailable to current terrestrial synthesis, and shell effects may produce lifetimes very different from measured neutron-deficient isotopes.

It is not established that such a nucleus controls gravitation. In LUCY, a nuclear state is at most a candidate seed or control variable; the cross-layer coupling is unknown and may be zero.

A nucleosynthesis exposure may be represented as

$$\Theta_r = \tau_{\text{env}} n_n \langle \sigma v \rangle.$$

This describes neutron-capture exposure and is not itself a LUCY or gravitational coupling.

8. Specific LUCY hypothesis

Let $q(x)$ be a dimensionless relational-closure order parameter, with $q = 0$ the baseline state and $q \neq 0$ a candidate closed state. Let λ collect material, geometric, electromagnetic, thermodynamic, and environmental controls.

The base hypothesis in this paper is that there exists a critical surface Σ_L in control-parameter space such that

$$q = 0 \longrightarrow q = q_*$$

marks a consequential physical crossing and, in the branch under test, produces a stable boundary with a correlated change in more than one independent class of physical observable.

The stronger gravitational hypothesis is that, after known material and field effects are removed, the $q \neq 0$ state modifies an effective geometry experienced by neutral matter and electromagnetic signals in a composition-independent manner consistent with a declared $g_{\text{eff}}(q)$.

The informational or relational variable must earn this physical status prospectively. If it is fully reducible to the adequate measured physical state, then the I-layer description may remain useful but this experiment has not established independent I→P dynamics.

Optical distortion alone is insufficient. Plasma, thermal gradients, nonlinear optics, electromagnetic forces, acoustic fields, vibration, and sensor artefacts can all create boundary-like phenomena. A metric interpretation rises only when one model explains independent matter, light, and clock observables with the same boundary, sign, timing, and threshold.

9. Discriminating predictions

Threshold and hysteresis

The response should be nonlinear in a preregistered control parameter and may show hysteresis. A smooth response proportional to applied power favours ordinary leakage, heating, or known field response.

Independent-content test

The candidate relational or informational variable must improve held-out prediction after the relevant physical state and history are controlled. Relation-preserving re-encoding should not alter the physical prediction unless the re-encoding itself changes a physical or observer variable.

Common boundary

Because this paper specifically studies the boundary-forming branch, optical phase, neutral test-mass motion, clock comparison, and fluid transport should change at one mapped boundary. Separate boundaries or unrelated onset times weaken the unified-geometry claim.

Geometry dependence

After controlling electromagnetic cavity modes, thermal transport, stress, and mass distribution, threshold or residual should depend on normalised boundary geometry under a declared scaling law.

Composition and charge

A metric-like effect should couple approximately universally to neutral test bodies. Strong dependence on charge, conductivity, magnetic susceptibility, dielectric constant, or surface finish favours ordinary material effects.

Wavelength dependence

A metric-like optical-path change should be nearly achromatic in vacuum after dispersion is removed. Strong frequency dependence favours plasma, refractive-index, cavity, or metamaterial mechanisms.

Clock and phase comparison

At least two independent clock or interferometric methods should show a consistent proper-time or phase residual before an altered-geometry interpretation is promoted.

Conservation accounting

Any apparent change in inertia, trajectory, or propagation requires complete source-energy, momentum, radiation, heat, and mechanical-reaction accounting. Boundary language is not an exemption from conservation laws.

Astrophysical signature

If a specific neutron-rich superheavy configuration is required, nuclear-network calculations should predict distinctive fission fragments, decay chains, or abundance residuals in r-process ejecta.

10. Promotion route

-
- **Tier 0 — mathematical consistency:** typed variables, units, cross-layer carrier where claimed, action or update rule, stability, ordinary limit, and causal structure.
 - **Tier 1 — conventional analogue:** threshold and, where applicable, membrane behaviour in a known system without a gravity claim.
 - **Tier 2 — cross-channel anomaly:** blinded reproducible common-boundary residual in at least two independent channels, with any proposed I→P variable adding held-out predictive value beyond the adequate physical baseline.
 - **Tier 3 — metric discrimination:** one effective geometry outpredicts the strongest electromagnetic, thermal, plasma, acoustic, mechanical, and sensor models across neutral matter, light, and clocks.
 - **Tier 4 — independent replication and scaling:** separate-laboratory replication of preregistered threshold, representation-invariance, scaling, hysteresis, composition, and wavelength predictions.

No ontology claim is promoted before Tier 3 and no foundational promotion before Tier 4.

11. Falsifiers

The gravitational branch fails or is materially reduced if known conventional models close the observations; the candidate informational relation reduces to an adequate physical-state description; relation-preserving re-encoding changes the claimed physical result without a physical reason; no dimensionally valid coupling from the order parameter to effective geometry can be constructed; general relativity and ordinary quantum field theory are not recovered in tested regimes; claimed boundaries differ across measurement channels; neutral-matter and clock effects are absent while optical or charged effects remain; the response is smooth where a critical transition was preregistered; scaling depends on uncontrolled apparatus details rather than declared invariants; energy or momentum accounting fails; the nuclear seed is unnecessary, nonspecific, or contradicted by data; parameters or thresholds are selected after the result; or an existing theory reproduces every predicted observable with equal or fewer assumptions.

A null result cannot be moved to an unmeasured layer. The failed branch must be revised or removed.

12. Relationship to canonical LUCY and MKUFT

This paper is downstream of the canonical [LUCY Boundary Threshold Framework](#). The canonical owner defines LUCY as the thresholded crossing at which an addressed relation becomes consequential at the next layer. This paper preserves that definition and studies one physical specialisation in which the crossing yields **LUCY-1 membrane** → **LUCY-2 sustained pocket** and potentially a different effective geometry.

The membrane and pocket are not retroactively imposed on non-boundary LUCY branches.

This paper also connects to [Layer Before Law](#), [Gradient Mechanics and Boundary-Shear Geometry](#), [Typed Traversal and Equation Hygiene](#), [Discriminating Experiments and Promotion Gates](#), and [Falsification Summary](#).

13. Compact rule

Do not amplify “gravity.” Define the addressed relation, define the $I \rightarrow P$ carrier, cross the threshold, and let the physical state decide its morphology. If that morphology is a boundary, map it; if it sustains a domain, test the pocket; if effective geometry is claimed, let independent matter, light, and clock observables decide whether geometry changed.

References

1. A. D. Sakharov, “Vacuum quantum fluctuations in curved space and the theory of gravitation,” *Sov. Phys. Dokl.* 12, 1040–1041 (1968), original 1967.
2. T. Jacobson, “Thermodynamics of Spacetime: The Einstein Equation of State,” *Physical Review Letters* 75, 1260–1263 (1995). DOI: 10.1103/PhysRevLett.75.1260.
3. C. Barceló, S. Liberati, and M. Visser, “Analogue Gravity,” *Living Reviews in Relativity* 14, 3 (2011). DOI: 10.12942/lrr-2011-3.
4. T. Padmanabhan, “Emergent Gravity Paradigm: Recent Progress,” *Modern Physics Letters A* 30, 1540007 (2015). arXiv:1410.6285.
5. D. Oriti, “Spacetime as a quantum many-body system,” arXiv:1710.02807 (2017).
6. J. Ben Achour et al., “Quantum Gravity, Hydrodynamics and Emergent Cosmology: A Collection of Perspectives,” arXiv:2411.12628 (2024).
7. T. Damour and G. Esposito-Farèse, “Nonperturbative Strong-Field Effects in Tensor-Scalar Theories of Gravitation,” *Physical Review Letters* 70, 2220–2223 (1993). DOI: 10.1103/PhysRevLett.70.2220.
8. D. M. Siegel, “r-Process nucleosynthesis in gravitational-wave and other explosive astrophysical events,” *Nature Reviews Physics* 4, 306–318 (2022).