




Dr Jasmeet Soar







Resuscitation Council (UK)



Chain of Survival



Chain of Prevention





The National Early Warning Score (NEWS)


PHYSIOLOGICAL PARAMETERS	3	2	1	0	1	2	3
Respiration Rate	≤8		9 - 11	12 - 20		21 - 24	≥25
Oxygen Saturations	≤91	92 - 93	94 - 95	≥96			
Any Supplemental Oxygen		Yes		No			
Temperature	≤35.0		35.1 - 36.0	36.1 - 38.0	38.1 - 39.0	≥39.1	
Systolic BP	≤90	91 - 100	101 - 110	111 - 219			≥220
Heart Rate	≤40		41 - 50	51 - 90	91 - 110	111 - 130	≥131
Level of Consciousness				A			V, P, or U



Adult Advanced Life Support







Unresponsive and not
breathing normally



To confirm cardiac arrest...

- Patient response
- Open airway
- Check for normal breathing
 - Caution agonal breathing
- Check for signs of life



Unresponsive and not
breathing normally




To confirm cardiac arrest...


- Pulse check if trained to do so
- Take less than 10 seconds for assessment
- Call for help early



Cardiac arrest confirmed



Cardiac arrest confirmed



Chest compression



- 30:2
- Compressions
 - centre of chest
 - 5-6 cm depth
 - 2 per second (100-120 min⁻¹)
- Maintain high quality compressions with minimal interruptions (<5 s)
- Continuous compressions once airway secured
- Switch CPR provider every 2 min cycle to avoid fatigue

Chest compression



- 30:2
- Compressions
 - centre of chest
 - 5-6 cm depth
 - 2 per second (100-120 min⁻¹)
- Maintain high quality compressions with minimal interruptions (<5 s)
- Continuous compressions once airway secured
- Switch CPR provider every 2 min cycle to avoid fatigue

Less than 5 second pause in chest compression for:

- Rhythm check
- Shock delivery
- Tracheal intubation



START

PAUSE

CPR

Assess rhythm

**Shockable
(VF/Pulseless VT)**

**Non-shockable
(PEA/Asystole)**

MINIMISE INTERRUPTIONS IN CHEST COMPRESSIONS



Resuscitation Council (UK)

GUIDELINES
2015

START

PAUSE

CPR

Assess rhythm

**Shockable
(VF/Pulseless VT)**

**Non-shockable
(PEA/Asystole)**


MINIMISE INTERRUPTIONS IN CHEST COMPRESSIONS



Resuscitation Council (UK)

GUIDELINES
2015

**Shockable
(VF/Pulseless VT)**




**IMMEDIATELY
RESTART CPR**



**Shockable
(VF/Pulseless VT)**


IMMEDIATELY
RESTART CPR




MINIMISE INTERRUPTIONS IN CHEST COMPRESSIONS



**Shockable
(VF/Pulseless VT)**





During CPR

- Ensure high quality chest compressions
- Minimise interruptions to compressions
- Give oxygen
- Use waveform capnography
- Continuous compressions when advanced airway in place
- Vascular access (intravenous or intraosseous)
- Give adrenaline every 3-5 min
- Give amiodarone after 3 shocks



Waveform capnography




Information from waveform capnography during CPR

- Tracheal tube placement
- Guide to rate of ventilation
- Quality of chest compressions
- ROSC
- Prognostication



Example end-tidal CO₂ trace



Information from waveform capnography during CPR

- Tracheal tube placement
- Guide to rate of ventilation
- Quality of chest compressions
- ROSC
- Prognostication



Information from waveform capnography during CPR

- **Tracheal tube placement**
- **Guide to rate of ventilation**
- Quality of chest compressions
- **ROSC**
- Prognostication



During CPR

- Ensure high quality chest compressions
- Minimise interruptions to compressions
- Give oxygen
- Use waveform capnography
- Continuous compressions when advanced airway in place
- Vascular access (intravenous or intraosseous)
- Give adrenaline every 3-5 min
- Give amiodarone after 3 shocks



Airway and ventilation during CPR

- Stepwise approach according to rescuer/patient
 - Compression-only
 - Mouth-to-mouth
 - Mouth-to-mask
 - Bag-mask
 - Supraglottic airway device (e.g. LMA, i-gel)
 - Tracheal tube (videolaryngoscopy)



Airway and ventilation during CPR

- Stepwise approach according to rescuer/patient
 - Compression-only
 - Mouth-to-mouth
 - Mouth-to-mask
 - Bag-mask
- Supraglottic airway device
- Tracheal tube
- 10 breaths/minute = continuous compressions



During CPR

- Ensure high quality chest compressions
- Minimise interruptions to compressions
- Give oxygen
- Use waveform capnography
- Continuous compressions when advanced airway in place
- Vascular access (intravenous or intraosseous)
- Give adrenaline every 3-5 min
- Give amiodarone after 3 shocks



Vascular access

- Peripheral versus central veins
- Intraosseous



Vascular access

- Peripheral veins

- Intraosseous

**ROLE OF DRUGS
LIMITED
DURING CPR**




Adrenaline & Amiodarone


- Adrenaline
 - Timings and dose not changed
 - Non-shockable – during CPR
 - Shockable – during third two minutes of CPR
 - Subsequent doses every 3 -5 min
- Amiodarone
 - After three shocks



Treat Reversible Causes

- Hypoxia
- Hypovolaemia
- Hypo-/hyperkalaemia/metabolic
- Hypothermia
- Thrombosis - coronary or pulmonary
- Tension pneumothorax
- Tamponade – cardiac
- Toxins





Consider

- Ultrasound imaging
- Mechanical chest compressions to facilitate transfer/treatment
- Coronary angiography and percutaneous coronary intervention
- Extracorporeal CPR

Assess rhythm



**Return of spontaneous
circulation**



**Immediate post cardiac
arrest treatment**

- Use ABCDE approach
- Aim for SpO₂ of 94-98%
- Aim for normal PaCO₂
- 12-lead ECG
- Treat precipitating cause
- Targeted temperature management



Resuscitation team

- Roles planned in advance
- Identify team leader
- Importance of non-technical skills
 - task management
 - team working
 - situational awareness
 - decision making
- Structured communication
 - SBAR or RSVP



Implementation



Advanced Life Support

Sixth Edition



Immediate Life Support

Third Edition



Advanced Life Support


Seventh Edition



Resuscitation Council (UK)

Immediate Life Support

Fourth Edition



Resuscitation Council (UK)



Resuscitation Council (UK)

GUIDELINES
2015

iResus

[View More by This Developer](#)

By RESUSCITATION COUNCIL (U.K.) TRADING LTD

Open iTunes to buy and download apps.



[View in iTunes](#)

✚ This app is designed for both iPhone and iPad

Free

Category: [Medical](#)

Released: 16 October 2015

Version: 1.0

Size: 4.7 MB

Language: English

Developer: RESUSCITATION COUNCIL (U.K.) TRADING LTD
© RESUSCITATION COUNCIL (U.K.) TRADING LTD

You must be at least 17 years old to download this application.

Infrequent/Mild Alcohol,

Tobacco, or Drug Use or

References

Frequent/Intense

Medical/Treatment Information

Compatibility: Requires iOS 8.0

or later. Compatible with

iPhone, iPad, and iPod touch.

Customer Ratings

Current Version:

★★★★★ 6 Ratings

Description

iResus is a free support tool that has been developed by the Resuscitation Council (UK) and Cranworth Medical Ltd.


iResus enables the user to access the latest Adult, Paediatric and Newborn resuscitation algorithms without the need for of an internet connection at the point of care.

iResus is fully compliant with the Resuscitation Council (UK) Guidelines 2015 and is an essential tool for UK doctors.

[RESUSCITATION COUNCIL \(U.K.\) TRADING LTD Web Site](#) • [iResus Support](#) •

Screenshots

iPhone | iPad



Resuscitation Council (UK)



Summary

- Importance of high quality chest compressions
- Minimise interruptions in chest compressions
- Monitoring during CPR
- Correct reversible causes of cardiac arrest
- Role of resuscitation team

