

# Fundamentals (Numbers, Sets, Words, Functions, and Relations)

---

**Exercise 1.** How many numbers are there between 100 and 1000 that are

- (a) divisible by 3?
- (b) divisible by 5?
- (c) divisible by 15?

**Solution:** Using the formula  $\lfloor \frac{m}{k} \rfloor - \lfloor \frac{n-1}{k} \rfloor$ :

- $\lfloor \frac{1000}{3} \rfloor - \lfloor \frac{99}{3} \rfloor = 300$  numbers divisible by 3 (102, 105, ..., 999);
- $\lfloor \frac{1000}{5} \rfloor - \lfloor \frac{99}{5} \rfloor = 181$  numbers divisible by 5 (100, 105, ..., 1000);
- $\lfloor \frac{1000}{15} \rfloor - \lfloor \frac{99}{15} \rfloor = 60$  numbers divisible by 15 (105, 120, ..., 990).

**Exercise 2.** Let  $\Sigma = \{a, b, c\}$  and  $\Phi = \{a, c, e\}$ .

- (a) How many words are in the set  $\Sigma^2$ ?
- (b) What are the elements of  $\Sigma^2 \setminus \Phi^*$ ?
- (c) Is it true that  $\Sigma^* \setminus \Phi^* = (\Sigma \setminus \Phi)^*$ ? Why?

**Solution:** (a)  $\Sigma^2 = \{aa, ab, ac, ba, \dots, cc\}$ , hence  $|\Sigma^2| = 3 \cdot 3 = 9$ .

(b)  $\Sigma^2 \setminus \Phi^* = \{ab, ba, bb, bc, cb\}$ , that is, all words in  $\Sigma^2$  with the letter  $b$ .

(c) No; for example,  $ab \in \Sigma^*$  and  $ab \notin \Phi^*$ , hence  $ab \in \Sigma^* \setminus \Phi^*$ ; but  $\Sigma \setminus \Phi = \{b\}$ , hence  $ab \notin (\Sigma \setminus \Phi)^*$ .

**Exercise 3.** Prove that  $(A \setminus B) \cup (B \setminus A) = (A \cup B) \setminus (A \cap B)$

**Solution:** We show in both directions that if an element belongs to  $(A \setminus B) \cup (B \setminus A)$  then it also belongs to  $(A \cup B) \setminus (A \cap B)$  and vice versa:

- Suppose that an element  $x \in (A \setminus B) \cup (B \setminus A)$ . Therefore, either  $x \in A \setminus B$  or  $x \in B \setminus A$ . In either case, we conclude that  $x \in A \cup B$  and (by the definition of set difference)  $x \notin A \cap B$ . Therefore,  $x \in (A \cup B) \setminus (A \cap B)$ .
- Suppose that  $x \in (A \cup B) \setminus (A \cap B)$ . This means that  $x \in A \cup B$  (and, therefore, either  $x \in A$  or  $x \in B$ ), but  $x \notin A \cap B$ . If  $x \in A$  and  $x \notin A \cap B$ , then  $x \in A \setminus B$ ; alternatively, if  $x \in B$  and  $x \notin A \cap B$ , then  $x \in B \setminus A$ . In either case, we conclude that  $x \in (A \setminus B) \cup (B \setminus A)$ .

**Exercise 4.** Consider the relation  $R \subseteq \mathbb{R} \times \mathbb{R}$  defined by  $aRb$  if, and only if,  $b + 0.5 \geq a \geq b - 0.5$ . Is  $R$

- (a) reflexive?
- (b) antireflexive?

- (c) symmetric?
- (d) antisymmetric?
- (e) transitive?

**Solution:**

- (a) Yes, since  $a + 0.5 \geq a \geq a - 0.5$  for all  $a \in \mathbb{R}$
- (b) No; see (a)
- (c) Yes, since  $(b + 0.5 \geq a) \wedge (a \geq b - 0.5)$  implies  $(b \geq a - 0.5) \wedge (a + 0.5 \geq b)$ .
- (d) No; e.g.  $(0, 0.1) \in R$  and  $(0.1, 0) \in R$ .
- (e) No; e.g.  $(1.1, 1.5) \in R$  and  $(1.5, 1.9) \in R$  but  $(1.1, 1.9) \notin R$  since  $1.9 - 0.5 > 1.1$

**Exercise 5.** For each of the following statements, provide a valid proof if it is true for all sets  $S$  and all relations  $R_1 \subseteq S \times S$  and  $R_2 \subseteq S \times S$ . If the statement is not always true, provide a counterexample.

- (a) If  $R_1$  and  $R_2$  are symmetric, then  $R_1 \cap R_2$  is symmetric.
- (b) If  $R_1$  and  $R_2$  are antisymmetric, then  $R_1 \cup R_2$  is antisymmetric.

**Solution:**

- (a) We will show that if  $R_1$  and  $R_2$  are symmetric, then  $R_1 \cap R_2$  is symmetric. Suppose  $(a, b) \in R_1 \cap R_2$ . Then  $(a, b) \in R_1$  and  $(a, b) \in R_2$ . Because  $R_1$  is symmetric,  $(b, a) \in R_1$ . Because  $R_2$  is symmetric,  $(b, a) \in R_2$ . Therefore  $(b, a) \in R_1 \cap R_2$ . Therefore  $R_1 \cap R_2$  is symmetric.
- (b) This is not the case. Consider relations on  $\mathbb{N}$ :  $R_1 = \leq$  and  $R_2 = \geq$ . We have  $1 \leq 2$ , so  $(1, 2) \in R_1$  and  $(2, 1) \in R_2$ . Therefore,  $(1, 2), (2, 1) \in R_1 \cup R_2$  but  $1 \neq 2$ .