


Vibrational Motion

The Wave Equation

In one dimension wave motion can be described by

$$\frac{\partial^2 \psi}{\partial t^2} = v^2 \frac{\partial^2 \psi}{\partial x^2}$$

Where ψ is a function that describes the wave (the wave function) and v is the speed of the wave. This equation tells us about how a wave propagates in space and time.



The general solutions to this equation can be expressed as the superposition of two waves propagating in opposite directions.


$$\psi(x, t) = f_1(x - vt) + f_2(x + vt)$$

where f_1 and f_2 are arbitrary functions. The fluctuations in a wave can be perpendicular to the direction of motion (a **transverse** wave), or parallel to the direction of motion (**longitudinal** wave).

Sinusoidal Waves

We can verify that this differential equation really does describe a wave. Let's start with the simplest of waves, a sinusoid.

We can draw two equivalent sine waves, one describing oscillations in position (x), and another describing oscillations in time (t).



The equation for this wave can be written as

$$\psi(x, t) = A \sin(kx - \omega t + \phi)$$

We can vary either time or position in this equation, and we would see a sinusoidal variation in amplitude of the wave function.

Wavelength, λ

The distance between successive peaks.

Amplitude, A

The maximum value of the wave function

Frequency, ν

The number of cycles per second.

Phase, ϕ

The initial offset of the wave in x at time $t=0$.

Angular Wavenumber,^{1,2} k

The number of wavelengths in the distance 2π . $k = 2\pi / \lambda$.

Angular Frequency, ω

The number of cycles per second, measured in radians. $\omega = 2\pi\nu$.

Wave Velocity³, v

The speed of the wave. $v = \lambda\nu = \omega/k$.

¹ Thus far we have only considered 1D waves, so we don't have to resort to vectors. In the case of multidimensional waves, k is called the wave vector and also specifies the direction of propagation of the wave.

² Don't confuse **angular wavenumber** ($k=2\pi/\lambda$) used when talking about waves, and **wavenumber** used in spectroscopy. In spectroscopy, wavenumber is the number of wavelengths in one unit of length. It is the spatial analogue of frequency and usually expressed in units of cm^{-1} .

³ There are actually several kinds of wave velocity, here we are talking about the group velocity.

So is this sinusoid a solution to the general wave equation?

Differentiate the wave function with respect to t at fixed x , twice

$$\psi(x, t) = A \sin(kx - \omega t + \phi)$$

$$\frac{\partial^2 \psi}{\partial t^2} = -A\omega^2 \sin(kx - \omega t + \phi) = -\omega^2 \psi(x, t)$$

Repeat the (partial) differentiation of the wave function, but now with respect to x at fixed t :


$$\frac{\partial^2 \psi}{\partial x^2} = -Ak^2 \sin[kx - \omega t + \phi] = -k^2 \psi(x, t)$$

Combining these two equations, making use of the definition of the wave velocity, $v = \omega/k$, gets us back to the linear wave equation.

$$\frac{\partial^2 \psi}{\partial t^2} = v^2 \frac{\partial^2 \psi}{\partial x^2}$$

Simple harmonic motion

One easy example of a wave equation is from simple harmonic motion. SHM is present where there is a restoring force that is proportional to the displacement of the system (e.g a pendulum, springs, molecular vibrations, sound waves, etc).

An Example: Mass and Spring

Consider the forces on the mass when it is displaced from its resting position by a distance x . We know that $F=ma$. The spring also applies a force given by **Hooke's law** proportional to the displacement $F_{\text{restoring}} = -kx$. Here k is the spring constant.

$$m \frac{d^2 x}{dt^2} = -kx$$

Another second order differential equation! Let's tidy up:

$$\frac{d^2 x}{dt^2} + \frac{k}{m} x = 0$$

You will learn how to solve such equations in your mathematics course. We can quote the solution and check it really works:


$$x(t) = A \sin(\omega t + \phi)$$

Where ω is the angular frequency $\omega^2 = k/m$. If we substitute this into the differential equation, we can check it is a solution and gives us the result.

$$T = \frac{2\pi}{\omega} = 2\pi \sqrt{\frac{m}{k}}$$

Interestingly, the period is independent of the initial displacement.

An Example: Simple Pendulum



The equation of motion for a simple pendulum is given by

$$ma = -mg \sin \theta$$

We can approximate for small angles with

$$\sin \theta \approx \theta$$

So the equation of motion becomes

$$ma = -mg\theta$$

The easiest way to solve this is to convert a to an angular form ($a = aL$)

$$m\alpha L = -mg\theta$$

Giving us the second order differential equation

$$\frac{d^2\theta}{dt^2} + \frac{g}{L}\theta = 0$$

Just as in the previous example, we can substitute in a sinusoidal solution to show that this gives a period of oscillation of

$$T = 2\pi\sqrt{\frac{L}{g}}$$

Energy in Simple Harmonic Motion

Potential Energy

$$V(t) = \frac{1}{2}kx^2$$

$$x(t) = A \sin(\omega t + \phi)$$

$$V(t) = \frac{1}{2}kA^2 \sin^2(\omega t + \phi)$$

Kinetic Energy

$$K(t) = \frac{1}{2}mv^2$$

$$v(t) = A\omega \cos(\omega t + \phi)$$

$$K(t) = \frac{1}{2}mA^2\omega^2 \cos^2(\omega t + \phi)$$



$$K(t) = \frac{1}{2}kA^2 \cos^2(\omega t + \phi)$$

Total Energy


$$E = K + V$$

$$E = \frac{1}{2}kA^2 [\sin^2(\omega t + \phi) + \cos^2(\omega t + \phi)] = \frac{1}{2}kA^2$$

The vibrational frequency of a harmonic oscillator is independent of the total energy.




Vibration Motion of a Diatomic Molecule



In the absence of external forces, the motion of the centre of mass (CM) of a molecule can be treated separately from the relative motion of the two atoms. The relative motion describes the time-dependent changes in bond length of the molecule

In this case, we have an analogous system to a mass on a spring. The only difference here is that we must again use the relative or **reduced mass**, μ , of the system rather than the actual mass, as we are only considering the internal motion.

$$\mu = \frac{m_1 m_2}{(m_1 + m_2)} \quad \omega = \sqrt{\frac{k}{\mu}}$$


μ is the reduced mass of the oscillator and ω is the angular frequency.

An Example: Vibrational Frequency of HBr

Treating HBr as a simple harmonic oscillator with force constant 412 Nm^{-1} , calculate the expected vibrational wavelength of the molecule. [$3.76 \text{ }\mu\text{m}$] (The actual vibrational wavenumber is $2649.7 \text{ cm}^{-1} = 3.77 \text{ }\mu\text{m}$). Not bad for just SHM.