

DATASCI 406 FA 26, Quiz 2.2

Name	Uniqname	UMID

Each question is worth 1 point. Circle one answer for each.

Question 1 (1 point)

Which line draws n values from the 50/50 mixture of $N(-2, 1)$ and $N(2, 1)$?

1. `0.5 * rnorm(n, -2) + 0.5 * rnorm(n, 2)`
2. `rnorm(n, mean = sample(c(-2, 2), n, replace = TRUE))`
3. `rnorm(n, mean = 0.5 * (-2) + 0.5 * 2)`
4. `sample(c(rnorm(1, -2), rnorm(1, 2)), n, replace = TRUE)`

Question 2 (1 point)

X has density $f(x) = \frac{3}{2}x^2$ on $-1 \leq x \leq 1$. On $[0, 1]$, $X \mid X > 0$ has CDF x^3 . Which line draws n values of X ?

1. `runif(n, -1, 1)^(1/3)`
2. `sample(c(-1, 1), n, replace = TRUE) * runif(n)^(1/3)`
3. `sample(c(-1, 1), n, replace = TRUE) * runif(n)^3`
4. `2 * runif(n)^(1/3) - 1`

Question 3 (1 point)

X and Y are independent $U(0, 1)$ and $Z = X + Y$. Which is true?

1. $Z \sim U(0, 2)$, since adding two uniforms gives a uniform.
2. Z is a 50/50 mixture of $U(0, 1)$ and $U(1, 2)$.
3. Z has density z on $[0, 1]$ and $2 - z$ on $[1, 2]$.
4. $Z \sim N(1, 1/6)$ exactly, by the central limit theorem.

Question 4 (1 point)

Under $H_0 : \mu = 0$, data are drawn from the mixture $0.8 N(0, 1) + 0.2 N(0, 10)$ (variances), and the test rejects when $|\bar{X}| \geq 1.96/\sqrt{n}$. Why does it reject more than 5% of the time?

1. The contamination moves the mean of the data away from 0.
2. Each observation has variance $0.8 \cdot 1 + 0.2 \cdot 10 = 2.8$, so $\bar{X}$ is more spread out than the cutoff assumes.
3. The sample mean is a biased estimator under contamination.
4. n is too small; with $n = 1000$ the size would be 0.05.