

Rejection and Accept-Reject Sampling: Recap

Mark M. Fredrickson (mfredric@umich.edu)

Module 2: Generate, Session 3

Computational Methods in Statistics and Data Science (DATASCI 406)

Rejection sampling

The target is a candidate conditioned on an event: $Y = X \mid X \in A$.

- Draw X and **keep the draws that land in A** .
- The fraction kept is $P(X \in A)$. When A is rare, most draws are wasted.
- Examples: $X \mid X$ odd, a truncated normal, points in the unit disk.

Truncated normal, $Y = Z \mid a \leq Z \leq b$:


$$F_Y(y) = \frac{\Phi(y) - \Phi(a)}{\Phi(b) - \Phi(a)}, \quad Q_Y(p) = \Phi^{-1}(\Phi(a) + p[\Phi(b) - \Phi(a)]).$$

Check the kept draws with a QQ plot against Q_Y .

Accept-reject

Target density f , candidate density g , and a constant c with $c g(x) \geq f(x)$ wherever $f(x) > 0$.

1. Draw $Y \sim g$ and $U \sim U(0, 1)$.
2. **Accept** $X = Y$ if $U \leq f(Y)/(c g(Y))$; otherwise reject.



Beta(2, 2) under a uniform envelope with $c = 1.5$: keep the points below f .

Why it works

The output is a conditional distribution: $V = Y \mid U \leq f(Y)/(c g(Y))$.

$$P(V \leq x) = \frac{P(Y \leq x, U \leq f(Y)/(c g(Y)))}{P(U \leq f(Y)/(c g(Y)))} = \frac{\frac{1}{c}F(x)}{\frac{1}{c}} = F(x)$$

- The numerator step $\int_0^{f(y)/(cg(y))} du = f(y)/(c g(y))$ needs the ratio to be **at most 1**. That is why $c g \geq f$.
- The denominator is the **acceptance rate**, $1/c$.

Choosing g and c

- The best c is $\max_x f(x)/g(x)$. Find it by calculus, or evaluate the ratio on a grid.
- A c that is too large is still correct but wastes candidates.
- From m candidates about m/c are kept, so $\text{Var}(\bar{X}) \approx c \text{Var}(X) / m$.
- A g closer in shape to f gives a smaller c . For the truncated normal on $[0, 1]$:

envelope	c	rejected
$U(0, 1)$	1.17	14%
$\frac{2}{3}(2 - y)$	1.06	6%

The uniform envelope's estimate has about 10% more variance.

When accept-reject fails

Tails too light. For $f(x) = 0.25 \phi(x + 2) + 0.75 \phi(x - 1)$ and $g = \phi$, $f(x)/\phi(x) \rightarrow \infty$, so no c exists.

- g 's tails must be at least as heavy as f 's.
- The Cauchy works: its tails fall like $1/x^2$.
- A c found by evaluating on a grid only covers the grid. Check that the ratio is bounded outside it.