

Thread - 2026-03-03

<https://x.com/RiikonenMarko/status/2028646139558257107>

1/28 - 2026-03-03 01:38 UTC

2/3 March 2026, Rovaniemi. The daytime display was followed by lunar. I had to photograph on a road lined by trees, no choice, the swarm was limited to this spot.

I was really struck by the strong helic and Tape arcs. Interestingly, there is not much Parry to talk about. I have some frames to stack, maybe it materializes, but in any case it must be weak and I wonder if triangulars are to blame. There seemed to be quite a lot of them in the sample.


Noticed the flip-anthelic arc once the worst of panic had subsided. I wonder how many lunar cases there are? Certainly the first one for me.


It appears that I forgot to check the focus.





2/28 - 2026-03-03 21:43 UTC

2/3 March 2026, Rovaniemi. Camera was here just 20 meters from the snow gun. Two swarms, 13 x 30s and 10 x 30s. The first one was pure column. Notice the lunar 46° infralateral arc at the top of the image. I saw it after the second stage was over, it was the only halo in the sky. After a while I drove there to take a look and got the lunar display of the above post. Should have headed there right from the start and concentrate only on lunar, these lunar + spotlight displays are not that appealing, too messy.










3/28 - 2026-03-04 07:52 UTC

2/3 March 2026, Rovaniemi. Single frames from the 2nd wave. The action turned plate-only before tailing out completely. Possibly in a couple of frames dual-path 46° contact arc is visible. The minor intensity difference between counternadir arc and counter-Dufay indicates quite thick plates. Helic arc visible in a couple of frames.

I am reluctant starting stacking the lunar display. The movement of the display across the image frame makes is difficult to align the halos. We would need a software that not only stacks but reprojects every frame into a common angular coordinate system.



4/28 - 2026-03-04 08:51 UTC


2/3 March 2026, Rovaniemi. This is the first photo I took, at 1943, upon arrival. The swarm that was weak to begin with was already subsiding here, but moments before I was pretty sure I saw 9° halo in it. Car temp was -11 C on arrival, meaning it would rate as the new high temperature record for ice fog odd radii if the halo was indeed real. At 1940 railway station was at -9.9 C, Apukka -11.6 C.





5/28 - 2026-03-04 09:30 UTC

2/3 March 2026, Rovaniemi. After the big helic arc display subsided I came back to the gunning site, getting some Lowitz action before stratus took over and ended the fun. This is 3f stack, just wish it would have lasted longer. I wonder if that apparent circular Lowitz is what is seems, or Liljequist arc, or both. Probably both.

Anyway, we should re-think about the Lowitz naming, because what we now call as upper and lower Lowitz become lower and upper Lowitz, respectively, when the light is below horizon. My nomenclature article has a handful suggestions for fix <https://t.co/xT63PoeU4T>








6/28 - 2026-03-04 09:35 UTC

2/3 March 2026, Rovaniemi. The swarm shifted and I followed, getting just this one usable frame before it was all over.



7/28 - 2026-03-17 17:20 UTC


2/3 March 2026, Rovaniemi. It's difficult to stack this display. These are versions of the first 8 frames


stacked of the full 17 frame set. Probably won't try to expand. If somebody wants to have a go, raw files are available: <https://t.co/Q0aLkPcjCu>


One thing that stands out in comparison to the single image of the starting post is that Parry materialized out. In b-r it looks actually more like Lowitz.


Looks like there is a suggestion of the "prism-basal 46° contact arc" or, if you like, " 46° contact arc C" (these are among the name options in my nomenclature article). This particular 46° contact arc, which is marked with an arrow in the once-over simulation (elevation 29.3°), has been observed only once before. But it's not resolved enough too call it a catch.

I'll post later sim parameters. There is of course more to say about the display. Also crystal photos are waiting to be published.









8/28 - 2026-03-18 15:15 UTC

2/3 March 2026, Rovaniemi. Parameters of the previous post's simulation. This is one of those Parry displays that are caused by crystals grown along the triangular dimension. Everything about Parry orientation halos in the display calls for using such crystals, the mean shape of which is displayed in the parameter table.


At this light elevation triangularish Parry crystals open the door for discovery of two novel halos. They are shown in the simulation that has only Parry and column orientation for clarity (I have touched upon these also in my nomenclature article).


One is an arc in the Ounasvaara family, from raypath 146. It is marked in the simulation by the lower arrow and may be called a type of infralateral Ounasvaara arc. Possibly it would have been visible were it not for the restless forest background.

The other one is a type of "rotated counter-Parry arc", from raypath 346, denoted by the upper arrow.

Not even the slightest suggestion of it is seen. One might try, but I doubt stacking all 17 frames would coax it out.

Triangulars are needed for the usual Ounasvaara arc, but it gets weak at this high light elevation. It's located at the helic arc – supralatarec crossing and some intensity enhancement is visible in that location in b-r, so possibly it is present. Taking it into account in simulation would seem to call for using even more triangular Parry crystals, up to the point where there is practically no shallowcave Parry at all, in which case the arc must be mainly Lowitz arc.








9/28 - 2026-03-18 17:43 UTC


2/3 March 2026, Rovaniemi. Moved the camera on the road but of course soon a car came, limiting the stack to 5 frames. Anyway, the continuation in b-r from Tape arc downwards on 46° halo persists and I am almost convinced that the prism-basal 46° contact arc / 46° contact arc C is there.

The brightest top part of the helic arc tangent to counterhelic arc really stood out visually. It was the part I spotted first, my initial thought what what extra arc is there above the zenith arc.












10/28 - 2026-03-19 07:04 UTC

2/3 March 2026, Rovaniemi. Five dark Nikon frames stacked. Something that has been bothering me already in the Canon shots is that while there is strong Lowitz B there doesn't seem to be much of the Lowitz A. These are raypath retrogrades of each other so their relative contributions can't be tinkered with by basal face shape adjustments. In a way this should not be surprising given we have serious troubles to explain the displays with related arcs from basal face reflection.

What else? Parry arc is definitely there and I am not quite seeing the 46° contact arc C that is suggested by the Canon photos. In b-r, the helic arc is very weakly drawn out by two lines – that's the red rim.

The shots were taken at the same time as the first Canon stack in this thread (moon elev 29.3). Images are flared up with red hot pixels.






11/28 - 2026-03-19 07:28 UTC

2/3 March 2026, Rovaniemi. The very first set (5 frames) I took of this stage of the display at half past

nine, moon 27.9° . The Lowitz anomaly really stands out in b-r.






12/28 - 2026-03-19 08:29 UTC

2/3 March 2026, Rovaniemi. 7 frame stack at lunar elevation 29.0° . This actually seems to support the presence of 46° contact arc C. Horrid hot pixels. Maybe the camera was a bit too hot to start with. When it's not that cold outside, the car heater blow can heat it up to a point where it is almost too hot to touch.

Just happened to notice that in one of the above Nikon stacks anthelion is peeking out from between the trees. Also the parhelic circle blue edge is visible.






13/28 - 2026-03-19 12:18 UTC

2/3 March 2026, Rovaniemi. A selection of photos from 2158 to 2233, which was the crystal sampling period. The green dot marks the start of crystal photography on the side of the road at 2022. As there



was no change in the display composition, it just got weaker, the sample is representative.





14/28 - 2026-03-19 13:09 UTC

2/3 March 2026, Rovaniemi. The crystal sample agrees well with the simulation's Parry crystal shapes: there is a profusion of high triangulars. In general, I get the feeling that regular hexagons have an unusually low representation here.

If I had only this sample to look at, I would say there is Moilanen arc in the display for those elongated crystals. But nothing whatsoever of Moilanen is visible in the stacks.





15/28 - 2026-03-19 13:43 UTC


2/3 March 2026, Rovaniemi. However, if Moilanen prism is truncated, it can get pretty non-existent at

the relatively high lunar elevation of the display. The HaloPoint simulations show two scenarios for 28° light elevation, full Moila prism and 0.3 truncated.

I should have checked with spotlight near-horizon light situation, of course, no big operation needed, a couple of quick photos with phone would have sufficed. But as there is no such documentation we are left to do with the one lucky shot where the approaching car makes a display. It does not show Moila, but this hardly constitutes hard evidence, so I would say the issue of whether there was Moila or not is left open.


I wonder what's the furthest above-horizon elevation at which Moilanen arc has been documented?








16/28 - 2026-03-19 13:44 UTC

2/3 March 2026, Rovaniemi. So back to crystals, here is another set of photos.











17/28 - 2026-03-19 13:45 UTC

2/3 March 2026, Rovaniemi. More crystals.











18/28 - 2026-03-21 05:51 UTC



2/3 March 2026, Rovaniemi. More crystals. Notice the massive big ass full triangular at the bottom left (arrow).







19/28 - 2026-03-21 05:57 UTC

2/3 March 2026, Rovaniemi. Another set.









20/28 - 2026-03-21 06:12 UTC

2/3 March 2026, Rovaniemi. Until now we have been looking at the simulations, saying triangulars

made the Parry halos in the display. I wonder if this the first time we have also the crystals sampled?









21/28 - 2026-03-21 06:21 UTC

2/3 March 2026, Rovaniemi. More crystals.



Do columnar triangulars orient differently to plate shaped triangulars? There is this pure column orientation display by Luomanen, on 13 Dec 2007 in Pälkäne, Finland, that seems to have been through and through triangulars made: <https://t.co/IW3GWB3HJD>







22/28 - 2026-03-21 06:27 UTC



2/3 March 2026, Rovaniemi. A lone pyramid in the third photo (arrow).






23/28 - 2026-03-21 06:38 UTC

2/3 March 2026, Rovaniemi. And this is the last crystals.







24/28 - 2026-03-21 07:08 UTC

2/3 March 2026, Rovaniemi. Why nobody told me that? Contrary to what I wrote in one of the earlier posts, it is actually possible to adjust the relative brightness between the retrograde raypath Lowitz arc pair (A and B / upper and lower). Just have the orientation equilibrium at a tilted angle.

Everything kind falls in place with this. Not only is the display's retrograde Lowitz arc situation reproduced right by the simulation, but also the Lowitz C / circular that seems to be present only above parhelia. This suggests there is an aspect ratio range where the triangulars take Parry orientation and then some range where they are "indecisive", wobbling in Lowitz orientation.





HALOPOINT 2.0

☐ Save work-file
☐ Save light work-file
☒ Open in Analysis Window

Start Simulation

Add to the Simulation

General Parameters

Sun Elevation [°] 28
Sun Diameter True Diameter
Number of rays 1000000
Colour of Light:
☐ Single Wavelength [nm] 600
☒ Wavelength Range
From 400 nm to 700 nm
Number of hits allowed: 50

Simulation View

Lens Type Fisheye, equidistant
Focal Length [mm] 7.6
Sensor Size [mm] W: 24 H: 36
☐ Custom parameters a: 2 b: 0.5
Center of Image Sun
Azimuth: 0
Elevation: 0
☒ Portrait
☐ Landscape Rotation (CW) 0

Appearance

Dots Coloured
Background User Defined
Shade Levels 8
☐ Show Coordinate Grid
Major Step [°] 10
Minor Step [°] 2
Style Faint
☒ Horizon ☐ Hide Subhorizon

Ice Crystal Population Parameters

%	C - axis			Rotation			Prism Aspect Ratio			Upper Pyramid				Lower Pyramid				
	Mean	Std	Distribution	Mean	Std	Distribution	Mean	Std	Distribution	Apex	h	Std	Distribution	Apex	h	Std	Distribution	
☐ 1	1	0	1	Gaussian	90	360	Uniform	0	0	Gaussian	62	1	0	Gaussian	56,142	0	0	Gaussian
☐ 2	1	0	1	Gaussian	90	360	Uniform	0	0	Gaussian	63	1	0	Gaussian	56,142	0	0	Gaussian
☐ 3	1	0	20	Gaussian	90	0.5	Uniform	0.3	0.1	Gaussian	64	0	0	Gaussian	56,142	0	0	Gaussian
☐ 4	50	85	0.5	Gaussian	85	0.4	Gaussian	1	0.04	Uniform	65	0	0	Gaussian	41	0	0	Gaussian
☒ 5	16	-20	10	Gaussian	90	0.5	Gaussian	0.4	0.1	Uniform	66	0	0	Gaussian	56,142	0	0	Gaussian
☒ 6	40	90	360	Gaussian	90	360	Uniform	1	0.3	Uniform	67	0	0	Gaussian	56,142	0	0	Gaussian
☐ 7	1	0	1.5	Gaussian	90	360	Uniform	0	0	Uniform	34	1	0	Gaussian	56,14	0	0	Uniform
☒ 8	50	0	2	Gaussian	90	360	Uniform	0.4	0.1	Uniform	34	0	0	Gaussian	56,142	0	0	Gaussian
☐ 9	0.5	90	0.4	Gaussian	90	0.4	Gaussian	0.7	0.1	Uniform	56,142	0	0	Gaussian	56,142	0	0	Gaussian
☐ 10	3	0	15	Gaussian	90	1	Uniform	0.05	0	Gaussian	56,142	0	0	Gaussian	56,142	0	0	Gaussian

Ice Crystal Visualisation

View Angle Control
y
z
Reset

Line Width, Front 2
Line Width, Back 1
Dash Length (Back) 5
Dash Spacing (Back) 5
Back Line Colour Gray
Ice Crystal Population 5
Draw Selected Crystal



Prism Face Distances

Apply	Face Distance from c-axis								Std of Distance								Sync	Distribution
	3	4	5	6	7	8	3	4	5	6	7	8						
☒ 1	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	☐ Gaussian				
☒ 2	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	☐ Gaussian				
☒ 3	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	☐ Uniform				
☐ 4	1.3	1	1.3	1	1.3	1	0.3	0	0.3	0	0.3	0	☒ Uniform					
☒ 5	1	2	1	2	1	2	0	0	0	0	0	0	☐ Uniform					
☐ 6	1.6	1	1.6	1	1.6	1	0.3	0	0.3	0	0.3	0	☒ Uniform					
☒ 7	0.1	1	1	0.1	1	1	0	0	0	0	0	0	☐ Gaussian					
☐ 8	0.1	1	1	0.1	0	0	0	0	0	0	0	0	☐ Uniform					
☐ 9	1	2	1	2	1	2	0	0	0	0	0	0	☒ Uniform					
☐ 10	1	2	1	1	2	1	0	0	0	0	0	0	☐ Gaussian					

25/28 - 2026-03-21 08:07 UTC

2/3 March 2026, Rovaniemi. Tilted triangulars also appear to allow isolate both convex (B) and concave (A) Liljequist arcs. That's how these arcs appear in many displays; it's practically either or. The crystal is tilted tip-up and tip-down, respectively.

Of course there is a slew of issues here, not the least of which is the necessity to restrict the raypaths to three face hits.



HALOPOINT 2.0

Raypath Filtering Enabled

☐ Save work-file
☐ Save light work-file
☒ Open in Analysis Window

Start Simulation

Add to the Simulation

General Parameters

Sun Elevation [°]
Sun Diameter
Number of rays
Colour of Light:
☐ Single Wavelength [nm]
☒ Wavelength Range
From nm to nm
Number of hits allowed:

Simulation View

Lens Type
Focal Length [mm]
Sensor Size [mm] W: H:
☐ Custom parameters a: b:
Center of Image
Azimuth:
Elevation:
☒ Portrait ☐ Landscape
Rotation (CW)

Appearance

Dots
Background
Shade Levels
☐ Show Coordinate Grid
Major Step [°]
Minor Step [°]
Style
☒ Horizon ☐ Hide Subhorizon

Ice Crystal Population Parameters

	%	C - axis			Rotation			Prism Aspect Ratio			Upper Pyramid				Lower Pyramid			
		Mean	Std	Distribution	Mean	Std	Distribution	Mean	Std	Distribution	Apex	h	Std	Distribution	Apex	h	Std	Distribution
1	1	0	1	Gaussian	90	360	Uniform	0	0	Gaussian	62	1	0	Gaussian	56,142	0	0	Gaussian
2	1	0	1	Gaussian	90	360	Uniform	0	0	Gaussian	63	1	0	Gaussian	56,142	0	0	Gaussian
3	1	0	20	Gaussian	90	0.5	Uniform	0.3	0.1	Gaussian	64	0	0	Gaussian	56,142	0	0	Gaussian
4	50	85	0.5	Gaussian	85	0.4	Gaussian	1	0.04	Uniform	65	0	0	Gaussian	41	0	0	Gaussian
5	16	20	10	Gaussian	90	0.5	Gaussian	0.4	0.1	Uniform	66	0	0	Gaussian	56,142	0	0	Gaussian
6	40	90	360	Gaussian	90	360	Uniform	1	0.3	Uniform	67	0	0	Gaussian	56,142	0	0	Gaussian
7	1	0	1.5	Gaussian	90	360	Uniform	0	0	Uniform	34	1	0	Gaussian	56,14	0	0	Uniform
8	50	0	2	Gaussian	90	360	Uniform	0.4	0.1	Uniform	34	0	0	Gaussian	56,142	0	0	Gaussian
9	0.5	90	0.4	Gaussian	90	0.4	Gaussian	0.7	0.1	Uniform	56,142	0	0	Gaussian	56,142	0	0	Gaussian
10	3	0	15	Gaussian	90	1	Uniform	0.05	0	Gaussian	56,142	0	0	Gaussian	56,142	0	0	Gaussian

Ice Crystal Visualisation

View Angle Control
y
z
Reset
Line Width, Front
Line Width, Back
Dash Length (Back)
Dash Spacing (Back)
Back Line Colour
Ice Crystal Population
Draw Selected Crystal

Prism Face Distances

Apply	Face Distance from c-axis								Std of Distance								Sync	Distribution
	3	4	5	6	7	8	3	4	5	6	7	8						
1	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	0	Gaussian			
2	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	0	Gaussian			
3	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	0	Uniform			
4	1.3	1	1.3	1	1.3	1	0.3	0	0.3	0	0.3	0	0	0	Uniform			
5	1	2	1	2	1	2	0	0	0	0	0	0	0	0	Uniform			
6	1.6	1	1.6	1	1.6	1	0.3	0	0.3	0	0.3	0	0	0	Uniform			
7	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	0	Gaussian			
8	0.1	1	1	0.1	0	0	0	0	0	0	0	0	0	0	Uniform			
9	1	2	1	2	1	2	0	0	0	0	0	0	0	0	Uniform			
10	1	2	1	1	2	1	0	0	0	0	0	0	0	0	Gaussian			

HALOPOINT 2.0

Raypath Filtering Enabled

☐ Save work-file
☐ Save light work-file
☒ Open in Analysis Window

Start Simulation

Add to the Simulation

General Parameters

Sun Elevation [°]
Sun Diameter
Number of rays
Colour of Light:
☐ Single Wavelength [nm]
☒ Wavelength Range
From nm to nm
Number of hits allowed:

Simulation View

Lens Type
Focal Length [mm]
Sensor Size [mm] W: H:
☐ Custom parameters a: b:
Center of Image
Azimuth:
Elevation:
☒ Portrait ☐ Landscape
Rotation (CW)

Appearance

Dots
Background
Shade Levels
☐ Show Coordinate Grid
Major Step [°]
Minor Step [°]
Style
☒ Horizon ☐ Hide Subhorizon

Ice Crystal Population Parameters

	%	C - axis			Rotation			Prism Aspect Ratio			Upper Pyramid				Lower Pyramid			
		Mean	Std	Distribution	Mean	Std	Distribution	Mean	Std	Distribution	Apex	h	Std	Distribution	Apex	h	Std	Distribution
1	1	0	1	Gaussian	90	360	Uniform	0	0	Gaussian	62	1	0	Gaussian	56,142	0	0	Gaussian
2	1	0	1	Gaussian	90	360	Uniform	0	0	Gaussian	63	1	0	Gaussian	56,142	0	0	Gaussian
3	1	0	20	Gaussian	90	0.5	Uniform	0.3	0.1	Gaussian	64	0	0	Gaussian	56,142	0	0	Gaussian
4	50	85	0.5	Gaussian	85	0.4	Gaussian	1	0.04	Uniform	65	0	0	Gaussian	41	0	0	Gaussian
5	16	20	10	Gaussian	90	0.5	Gaussian	0.4	0.1	Uniform	66	0	0	Gaussian	56,142	0	0	Gaussian
6	40	90	360	Gaussian	90	360	Uniform	1	0.3	Uniform	67	0	0	Gaussian	56,142	0	0	Gaussian
7	1	0	1.5	Gaussian	90	360	Uniform	0	0	Uniform	34	1	0	Gaussian	56,14	0	0	Uniform
8	50	0	2	Gaussian	90	360	Uniform	0.4	0.1	Uniform	34	0	0	Gaussian	56,142	0	0	Gaussian
9	0.5	90	0.4	Gaussian	90	0.4	Gaussian	0.7	0.1	Uniform	56,142	0	0	Gaussian	56,142	0	0	Gaussian
10	3	0	15	Gaussian	90	1	Uniform	0.05	0	Gaussian	56,142	0	0	Gaussian	56,142	0	0	Gaussian

Ice Crystal Visualisation

View Angle Control
y
z
Reset
Line Width, Front
Line Width, Back
Dash Length (Back)
Dash Spacing (Back)
Back Line Colour
Ice Crystal Population
Draw Selected Crystal



Prism Face Distances

Apply	Face Distance from c-axis								Std of Distance								Sync	Distribution
	3	4	5	6	7	8	3	4	5	6	7	8						
1	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	0	Gaussian			
2	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	0	Gaussian			
3	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	0	Uniform			
4	1.3	1	1.3	1	1.3	1	0.3	0	0.3	0	0.3	0	0	0	Uniform			
5	1	2	1	2	1	2	0	0	0	0	0	0	0	0	Uniform			
6	1.6	1	1.6	1	1.6	1	0.3	0	0.3	0	0.3	0	0	0	Uniform			
7	0.1	1	1	0.1	1	1	0	0	0	0	0	0	0	0	Gaussian			
8	0.1	1	1	0.1	0	0	0	0	0	0	0	0	0	0	Uniform			
9	1	2	1	2	1	2	0	0	0	0	0	0	0	0	Uniform			
10	1	2	1	1	2	1	0	0	0	0	0	0	0	0	Gaussian			

26/28 - 2026-03-24 08:13 UTC

2/3 March 2026, Rovaniemi. This one I took while I was having also display in the beam. 3 x 30s.

Moon 22.7°. Tanarc in the building light at the bottom left.



27/28 - 2026-03-24 08:46 UTC

2/3 March 2026, Rovaniemi. When I left the snow making site to check the display at where the 46° infralateral arc was visible at the horizon, I took just 5 photos upon arrival at the location, resulting in the shown stack. I continued driving towards lake Sierijärvi in hope having the display in more open setting. But there was nothing there.

Coming back 10 minutes later to the same spot at the Sierijärventie 69 junction, the display had changed as shown by the side-by-side comparison. Stepping out of the car my first thought was that it was for the worse. Then I spotted the helic arc and Tape arcs. Interesting how the earlier stage has brighter Parry arc yet other related halos are much poorer.





28/28 - 2026-03-24 11:30 UTC

2/3 March 2026, Rovaniemi. Looks like I am thru all the material from this night. I may still try to attempt a full stack of the 17 frame Canon set, but for now I am back to the solar display.

