



Index

A

Acceptance, of hypothesis. *See* Hypothesis testing

Additive property, of chi-square random variables, 264

Algebra of events, 56–57, 56f, 57f

Analysis, of residuals, 384–386, 385–386f

Analysis of variance (ANOVA), 445–476

- introduction, 445–446
- one-way, 446, 448–459, 452t, 453f, 456t
 - multiple comparisons of sample means, 456–458
 - with unequal sample sizes, 458–459
- overview, 446–448
- two-way, 446
 - hypothesis testing for, 464–468, 467t, 468f
 - interaction, 446, 469–476, 475t, 476f
 - introduction and parameter estimation, 460–463

ANOVA. *See* Analysis of variance

Approximately normal, 31, 31f

Assessment, of models, 384–386, 385–386f

Assignable cause, 553

Association, causation v., 39

Associative law, 56–57

Averaging notation, in two-way analysis of variance, 461–462

B

Balanced case, in one-way analysis of variance, 459

Bar graph, 10, 10t, 11f

Bayes estimator, 236, 279–284

Bayes' formula, 68–75, 69f

Bayesian approach, 236, 604–606

Behrens-Fisher problem, 324–325

Bernoulli density function, 419

Bernoulli, Jacob, 5–6

Bernoulli, James, 141

Bernoulli parameter, maximum likelihood estimator of, 237–240

Bernoulli populations

- hypothesis testing in, 329–336
- testing equality of parameters in two of, 333–336

Bernoulli random variables, 141–148, 143f, 148f

- confidence interval for mean of, 268–272, 272t
- generation of, 639
- hypergeometric random variables and, 157–158
- sign test and, 522
- signed rank test and, 526–527

Beta distribution, 282

Between samples sum of squares, 451–452, 454, 456t, 459

Bias, of estimators, 274

Bimodal data set, 33, 34f

Binary output data, logistic regression models for, 418–421, 418f

Binomial distribution function, computation of, 147–148, 148f

Binomial distribution, hypothesis testing in, 329–336

Binomial random variables, 141–148, 143f, 148f

- binomial distribution function, 147–148, 148f
- central limit theorem and, 212, 215–216, 216f
- generation of, 639–640
- hypergeometric random variables and, 158–160, 225–226

Binomial random variables (*continued*)
 Poisson random variables and, 150–153
 sign test and, 522
 Bootstrap method, 623–630
 mean square error and, 627–628
 population mean and, 623–624, 628–630
 population variance and, 623–625
 probability and, 629–630
 random variable generation, 627
 Box plot, 27, 27f
 Box-Muller method, 642

C

Categorical data analysis, goodness of fit tests
 and, 489–514
 critical region determination by simulation,
 496–499, 498f
 dummy variables for, 416–418
 introduction, 489–490
 Kolmogorov–Smirnov goodness of fit test for
 continuous data, 510–514, 511f
 tests of independence in contingency tables,
 501–505
 tests of independence in contingency tables
 having fixed marginal totals, 505–510,
 506f
 tests when all parameters are specified,
 489–497, 493t, 498f
 tests when some parameters are unspecified,
 499–501
 Causation, association v., 39
 Central limit theorem, 210–219, 212–216f,
 approximate distribution of sample mean,
 216–218
 binomial random variables and, 212,
 215–216, 216f
 defined, 210
 independent random variables, 226–227
 Monte Carlo study and, 643
 probability mass function and, 212–215f
 sample size needed, 218–219, 219f
 Chance variation, 553
 Channel noise disturbance, 173
 Chebyshev's inequality, 27–30, 29t
 one-sided, 29–30
 random variables and, 128–131
 weak law of large numbers, 129
 Chi-square approximation, 497–498
 Chi-square density function, 188–193,
 190f, 192f
 Chi-square distribution, 188–193, 190f, 192f,
 327–328, 607
 gamma random variables and, 191–192, 192f
t-distribution and, 193–195, 193f, 194f
 Chi-square goodness of fit tests, history of, 6
 Chi-square random variables, 189, 190f
 additive property of, 264
 estimators of variance from, 447–448
F-distribution and, 195–196, 195f
 mean and variance of, 192
 sample mean and variance distribution with,
 222
 Choosing, of normal prior, 283–284
 Class boundaries, 15, 15f, 15t
 Class intervals, 14–15, 15f, 15t
 Classical approximation, for rank sum test,
 535–537
 Classical simulation, for rank sum test,
 537–539, 538f
 Coefficient of determination, sample correlation
 coefficient and, 382–384
 Coefficient of multiple determination, 411
 Column factors
 hypothesis testing for, 464–468, 467t, 468f
 in two-way analysis of variance, 460
 deviation from grand mean due to, 462
 row factor interaction with, 446, 469–476,
 475t, 476f
 Column sum of squares, 467t
 Combinations, permutations and, 63
 Combining, of unbiased estimators, 275–277
 Common density function, independent
 variables and, 102–103
 Commutative law, 56–57, 56f
 Comparison, of sample means, 456–458
 Complement
 in sample space, 55
 in Venn diagram, 56–57, 56f, 57f
 Composite hypothesis, 298
 Computational identity, of sum of squares of
 residuals, 366
 Conditional densities, 106–107
 Conditional distributions, 104–107,
 337–339
 random numbers and, 164–166, 166f
 Conditional probability, 65–68, 66f, 106
 independent events and, 75–79
 Poisson random variables and, 153–154
 Conditional probability density function,
 280–281

- Conditional probability mass function, 105–106
 - Confidence, 235
 - Confidence interval. *See also specific intervals*, e.g. 95 Percent confidence interval
 - for difference in means of two normal distributions, 260–268, 262–263f, 266–267f, 260t, 268t
 - for difference in population means, 456–458
 - for estimating unknown mean, 247–252
 - for exponential distribution in life testing, 594–596
 - for mean of Bernoulli random variable, 268–272, 272t
 - for mean of exponential distribution, 272–273
 - for normal mean with unknown variance, 252–258, 256f
 - one-sided lower, 249–250, 255–257, 256f, 260t, 261–265, 261–263f, 268t
 - one-sided upper, 249–250, 254–255, 260t, 261, 265–268, 266–267f
 - prediction interval v ., 381
 - for regression parameters
 - α , 376–377
 - β , 371–372
 - in sequential testing for exponential distribution in life testing, 599–601
 - two-sided, 248, 251, 255–257, 256f, 261–268, 262–263f, 266–267f
 - for unknown probability, 268–272, 272t
 - for variance of normal distribution, 259–260, 260t
 - Confidence interval estimators
 - of difference in means of two normal distributions, 260–265, 262–263f
 - for mean of exponential distribution, 272
 - of mean response, 377–379, 411–413
 - Contingency tables
 - with fixed marginal totals, tests of independence in, 505–510, 506f
 - tests of independence in, 501–505
 - Continuous data, Kolmogorov–Smirnov goodness of fit test for, 510–514, 511f
 - Continuous random variables, 93–94, 640–643
 - Control charts, 553
 - fraction defective, 565–567
 - mean, 554–562, 556f, 564f
 - case of unknown, 557–562, 559t
 - number of defect, 567–570, 569t
 - for population mean, 571–581
 - moving-average, 571–573, 573t, 574f
 - variance, 562–565, 564f
 - Control group, 164
 - Control limits
 - for fraction defective, 566–567
 - for mean control charts, 555–557, 556f
 - for moving-average, 572–573, 573t, 574f
 - for variance control charts, 563–564, 564f
 - Controls, 336
 - Correlation analysis, history of, 6
 - Counting
 - basic principles of, 60–63
 - notation and terminology, 63–65
 - Covariance, 121–122
 - Critical region, 298, 300
 - for goodness of fit tests when all parameters are specified, 490–491
 - one-sided, 307
 - simulation for determination of, 496–499, 498f
 - Critical value, for determining independence of characteristics of population member, 504
 - Cumulative distribution function, 91–93, 93f, 96
 - exponential random variables and, 177
 - probability density function and, 94–95, 94f
 - Cumulative sum control charts, 579–581
- D**
- Data collection, descriptive statistics and, 1–2
 - Data sets
 - describing, 9–17
 - frequency tables and graphs, 10, 10t, 11f, 12f
 - grouped data, histograms, ogives, and stem and leaf plots, 14–17, 14t, 15f, 15t, 16f, 18t
 - relative frequency tables and graphs, 10–14, 13–14f, 13t
 - normal and skewed, 31–33, 31f, 32f, 34f
 - summarizing, 17–27
 - sample mean, sample median, and sample mode, 17–22
 - sample percentiles and box plots, 24–27, 26t, 27f
 - sample variance and sample standard deviation, 22–24
 - Defects, probability of, 329–337
 - DeMorgan's laws, 57

- Density function. *See also* Probability density function
 Bernoulli, 419
 chi-square, 188–193, 190f, 192f
 common, 102–103
 conditional probability, 280–281
 F , 195–196, 195f
 gamma distribution and, 186–188
 joint, 236–237, 242, 244
 joint probability, 99–101
 of logistics distribution, 197
 normal, 168, 168f, 188, 189f
 posterior, 280, 284
 random variables and, 112
 Rayleigh, 591
 t , 193, 193f, 252, 253t
 Weibull, 608, 609f
- Density, mode of, 284
- Dependent events, 75–79, 78f
- Dependent variable. *See* Response variable
- DES. *See* Diethylstilbestrol
- Descriptive statistics
 Chebyshev's inequality, 27–30, 29t
 data collection and, 1–2
 describing data sets, 9–17
 frequency tables and graphs, 10, 10t, 11f, 12f
 grouped data, histograms, ogives, and stem and leaf plots, 14–17, 14t, 15f, 15t, 16f, 18t
 relative frequency tables and graphs, 10–14, 13–14f, 13t
 history of, 7
 normal data sets, 31–34, 32f, 34f
 paired data sets and the sample correlation coefficient, 34–41, 35t, 36f, 39f
 summarizing data sets, 17–27
 sample mean, sample median, and sample mode, 17–22
 sample percentiles and box plots, 24–27, 26t, 27f
 sample variance and sample standard deviation, 22–24
- Deviation from grand mean due to column j , 462
- Deviation from grand mean due to row i , 462
- Diethylstilbestrol (DES), 336
- Difference, in means of two normal distributions, 260–268, 262–263f, 266–267f, 268t
- Discrete inverse transform method, 639–640
- Discrete random variables, 91–92
 expectation and, 111, 111f
 generation of, 638–640
 probability mass function and, 92–93, 638–640
- Dispersion parameter, 197
- Distribution
 binomial, hypothesis testing in, 329–336
 chi-square, 188–193, 190f, 192f, 327–328, 607
 conditional, 337–339
 exponential, confidence interval for mean of, 272–273
 F , 328–329
 gamma, 595, 604–607
 hypothesis testing for determining equality of m population distributions, 508–509
 of least squares estimators, 361–367, 368f, 369f
 life, 245–247
 multivariate normal, 404
 normal
 confidence interval for variance of, 259–260, 260t
 estimation of difference in means of, 260–268, 262–263f, 266–267f, 268t
 Poisson
 goodness of fit tests for, 499–501
 hypothesis testing concerning mean of, 336–339
 variance in, 395–396
 prior, 279–284, 604–606
 probability, of estimator of mean response, 377–378
 rate of, 590
 of sample, goodness of fit tests for, 489–499, 493t, 498f
 uniform, estimating mean of, 244–245
- Distribution function. *See also* Cumulative distribution function; Probability distribution function
 binomial, 147–148, 148f
 of continuous random variable, 640–641
 empirical, 624–625
 moment generating function and, 128
 of normal random variables, 170–171
 Poisson
 computation of, 155–156
 number of defects and, 567–570, 569t
 probability and, 91–92
 random variables and, 91–92, 624

- of rank sum test, 531
- signed rank test for, 525, 526f
- two-sample problem and, 531
- Distribution results, summary of, 381–382
- Distributive law, 56–57, 57f
- Doll, R., 17
- Dot notation, in two-way analysis of variance, 461–462
- Double-blind test, 164
- Dummy variables, for categorical data, 416–418

E

- Effect of column j , 470
- Effect of row i , 470
- Empirical distribution function, 624–625
- Empirical rule, 33–34
- Entropy, 110
- Equal variance, testing equality of means of two normal populations with, 324–325, 325t
- Equality
 - of m population distributions, hypothesis testing for, 508–509
 - of means of two normal populations, 318–326, 319t, 321f, 323f, 325t
 - case of known variance, 318–320, 319t
 - case of unknown and equal variance, 324–325, 325t
 - case of unknown variance, 320–324, 321f, 323f
 - hypothesis testing of, 318–326, 319t, 321f, 323f, 325t
 - paired t -test, 325–326
 - of parameters in two Bernoulli populations, 333–336
 - of population means, hypothesis testing of, 446, 448–459, 452t, 453f, 456t
 - of variance, of two normal populations, 328–329
- Error
 - mean square error of point estimators, 273–279
 - type I, 298, 300
 - type II, 298, 302–305, 304f
- Error sum of squares, 465, 467t
- Estimated regression line, 360
- Estimates
 - defined, 236
 - interval, 235, 247–260
 - confidence interval for normal mean with unknown variance, 252–258, 256f

- confidence interval for variance of normal distribution, 259–260, 260t
- hypothesis testing v., 310
- prediction, 257–258
- for unknown mean, 247–252
- Estimation
 - of life distributions, 245–247
 - of mean of uniform distribution, 244–245
 - of mean response, 411–413
 - of parameters, 235–284
 - approximate confidence interval for mean of Bernoulli random variable, 268–272, 272t
 - Bayes estimator, 236, 279–284
 - confidence interval for mean of exponential distribution, 272–273
 - of difference in means of two normal distributions, 260–268, 262–263f, 266–267f, 268t
 - interval estimates, 235, 247–260, 310
 - introduction, 235–236
 - of life distributions, 245–247
 - maximum likelihood estimators, 235–247, 260, 284, 500
 - point estimator evaluation, 273–279
 - for two-way analysis of variance, 460–463
- Estimators
 - Bayes, 236, 279–284
 - bias of, 274
 - confidence interval
 - of difference in means of two normal distributions, 260–265, 262–263f
 - for mean of exponential distribution, 272
 - of mean response, 377–379, 411–413
 - defined, 236
 - of deviance from grand mean, 462–463
 - of grand mean, 462–463
 - least squares, 385
 - distribution of, 361–367, 368f, 369f
 - in multiple linear regressions, 400–408, 413–414
 - in polynomial regression, 397–399
 - of regression parameters, 359–361, 361f, 362f
 - for Weibull distribution in life testing, 610–612
 - maximum likelihood, 235–247, 279
 - of Bernoulli parameter, 237–240
 - of difference in means of two normal distributions, 260
 - evaluation of, 277–278

Estimators (*continued*)

- for exponential distribution in life testing, 593–594, 602–604
- least squares estimators as, 365–366
- for life distributions, 245–247
- in logistic regression models, 418
- for mean of exponential distribution, 272
- of normal population, 242–244
- of Poisson parameter, 240–241
- in sequential testing for exponential distribution in life testing, 599, 601
- for Weibull distribution in life testing, 608–610
- weighted least squares estimators as, 392
- point
 - evaluation of, 273–279
 - for hypothesis testing, 299–300
 - of mean response, 377, 411
- pooled, 261, 321
- unbiased, 274–279
- of variance, 447–448
 - for one-way analysis of variance, 446, 448–459, 452t, 453f, 456t
 - for two-way analysis of variance, 464–467
 - for two-way analysis of variance with interaction, 469–476, 475t, 476f
 - weighted least squares, 390–396, 395f
- Evaluation, of point estimator, 273–279
- Events, 54–55
 - algebra of, 56–57, 56f, 57f
 - independent, 75–79, 78f
 - odds of, 59
- Expectation, 107–111, 111f
 - properties of, 111–118
 - of a random variable function, 113–115
 - of sums of random variables, 115–118
- Expected value. *See* Expectation
- Exponential distribution
 - confidence interval for mean of, 272–273
 - gamma distribution and, 188
 - in life testing, 592–606, 598f
 - Bayesian approach, 604–606
 - sequential testing, 598–602, 598f
 - simulation testing with stopping at r th failure, 592–598
 - simulation testing with stopping by fixed time, 602–604
 - Poisson process and, 181–182
- Exponential random variables, 177–185, 181f
 - generation of, 641–642
 - memoryless, 178

moment generating functions and, 177

Poisson process, 181–183, 181f

probability and, 179–180

sample means for, 218–219, 219f

Exponentially weighted moving-average control charts, 573–578, 575f, 578f

F

Failure rate. *See* Hazard rate

F -density function, 195–196, 195f

F -distribution, 195–196, 195f, 328–329

Finite populations, sampling distributions from, 223–227

First quartile, 25–27

Fisher, Ronald A., 6–7

Fisher-Irwin test, 335

Fixed margins, contingency tables with, tests of independence in, 505–510, 506f

Fraction defective control charts, 565–567

Frequency interpretation

of expectations, 108

probability, 53

Frequency tables and graphs, 10, 10t, 11f, 12f

frequency histogram, 16

frequency polygon, 10, 10t, 12f

relative, 10–14, 13–14f, 13t

sample mean and, 19–20, 22

sample median and, 20–22

sample mode and, 22

F -statistic, in two-way analysis of variance with interaction, 474, 475t

Future response, prediction interval of, 379–381
in multiple linear regression, 411–416, 413t, 414f, 415f

G

Galton, Francis, 6, 372–373

Gamma density, 188, 189f

Gamma distribution, 595, 604–607

Gamma function, 186–187

Gamma random variables, 187–188, 189f

chi-square distribution and, 191–192, 192f

Gauss, Karl Friedrich, 5–6

Generation

of random numbers, 620–622

of random variables, 497, 627, 638–643

Goodness of fit tests, 489–514

critical region determination by simulation, 496–499, 498f

- introduction, 489–490
 - Kolmogorov–Smirnov goodness of fit test for continuous data, 510–514, 511f
 - tests of independence in contingency tables, 501–505
 - tests of independence in contingency tables having fixed marginal totals, 505–510, 506f
 - tests when all parameters are specified, 489–497, 493t, 498f
 - tests when some parameters are unspecified, 499–501
 - Gosset, W.S., 6
 - Grand mean, 462–463, 469
 - Graunt, John, 4–5, 4t, 5t
 - Grouped data, 14–17, 14t, 15f, 15t, 16f, 18t
- H**
- Halley, Edmund, 5
 - Hardy's lemma, 36
 - Hazard rate, 245, 589
 - Hazard rate functions, 589–592
 - Hill, A.B., 17
 - Histograms, 14t, 15f, 16–17, 18t
 - normal, 31–33, 31f, 32f, 34f
 - Hypergeometric random variables, 156–160
 - Bernoulli random variables and, 157–158
 - binomial random variables and, 158–160, 225–226
 - mean and variance of, 157–158
 - Hypothesis testing, 297–339
 - in Bernoulli populations, 329–336
 - of equality of m population distributions, 508–509
 - of equality of means of two normal populations, 318–326, 319t, 321f, 323f, 325t
 - case of known variance, 318–320, 319t
 - case of unknown and equal variance, 324–325, 325t
 - case of unknown variance, 320–324, 321f, 323f
 - paired t -test, 325–326
 - of equality of population means, 446, 448–459, 452t, 453f, 456t
 - of equality of variance of two normal populations, 328–329
 - of independence in contingency tables, 501–505
 - of independence in contingency tables having fixed marginal totals, 505–510, 506f
 - of independence of characteristics of population member, 501–505
 - interval estimates v., 310
 - introduction, 297
 - of mean of normal population, 299–317, 301f, 304f, 311t, 313f, 315f, 317t
 - case of known variance, 299–311, 301f, 304f, 311t
 - case of unknown variance, 311–317, 313f, 316f, 317t
 - for mean of Poisson distribution, 336–339
 - multiple linear regression and, 409–411, 410t
 - of multiple population means, 446–447
 - of probability distribution of sample, 489–499, 493t, 498f
 - of regression parameters
 - α , 376–377
 - β , 370–371
 - of regression to mean, 373–374
 - robustness of, 311
 - of row and column interaction, 469–476, 475t, 476f
 - significance levels, 298–299
 - for two-way analysis of variance, 464–468, 467t, 468f
 - for variance of normal population, 327–329

I

 - Independence, tests of
 - in contingency tables, 501–505
 - in contingency tables having fixed marginal totals, 505–510, 506f
 - Independent events, 75–79, 78f
 - Independent increment assumption, 181
 - Independent random variables, 101–104
 - central limit theorem for, 210–219, 226–227
 - moment generating functions of, 127–128
 - sample mean and variance distribution with, 222
 - sample mean distribution with, 2, 15
 - signed rank test and, 526–527
 - Independent variable. *See* Input variable
 - Indicator random variable, 90–91
 - covariance of, 125–126
 - expectation for, 109
 - variance of, 119–120
 - Individual moment generating functions, 127–128

Individual probability mass function, joint and, 96–99, 98t

Inferential statistics
 history of, 6–7
 probability models and, 2–3

Inheritance, regression to mean and, 372–375, 373f, 374f

Input variable, 357–358
 variation in response to, 382–384, 390–396, 395f

Interaction, two-way analysis of variance with, 446, 469–476, 475t, 476f

Intersection
 of sample space, 55
 in Venn diagram, 56–57, 56f, 57f

Interval estimates, 235, 247–260
 confidence interval for normal mean with unknown variance, 252–258, 256f
 confidence interval for variance of normal distribution, 259–260, 260t
 hypothesis testing v., 310
 prediction, 257–258
 for unknown mean, 247–252

Inverse transformation method, 639–641

i th order statistic, 594

J

Joint cumulative probability distribution function, 96, 103–104

Joint density
 conditional densities and, 107
 random numbers and, 166–168

Joint density function, 236–237, 242, 244

Joint probability density function, 99–101

Joint probability mass function
 conditional probability mass function and, 106
 individual and, 96–99, 98t, 99t

Jointly continuous, 99, 102–103

Jointly distributed random variables, 95–107, 98t, 99t
 conditional distributions, 104–107
 independent, 101–104

K

Kolmogorov's law of fragmentation, 243

Kolmogorov–Smirnov goodness of fit test, for continuous data, 510–514, 511f

Kolmogorov–Smirnov test statistic, 510–514, 511f

Kruskal–Wallis test, 540

L

Laplace, Pierre-Simon, 5–6

Least squares estimators, 385
 distribution of, 361–367, 368f, 369f
 in multiple linear regression, 400–408, 413–414
 in polynomial regression, 397–399
 of regression parameters, 359–361, 361f, 362f
 for Weibull distribution in life testing, 610–612
 weighted, 390–396, 395f

Left-end inclusion convention, 15

Level of significance. *See* Significance level

Levels, in two-way analysis of variance, 460

Life distributions, estimation of, 245–247

Life testing, 589–612
 exponential distribution in, 592–606, 598f
 Bayesian approach, 604–606
 sequential testing, 598–602, 604f
 simulation testing with stopping at r th failure, 592–598
 simulation testing with stopping by fixed time, 602–604
 hazard rate functions, 589–592
 introduction, 589
 two-sample problem, 606–608
 Weibull distribution in, 608–612, 609f

Likelihood function, 236–237

Line graph, 10, 10t, 11f

Linear regression equation, 357–358. *See also* Multiple linear regression
 assessment of, 384–386, 385–386f

Linearity, transforming to, 387–390, 388f, 389f, 389t, 390t

Logarithms, for transforming to linearity, 387–390, 388f, 389f, 389t, 390t

Logistic regression function, 416, 418f

Logistic regression models, for binary output data, 416–419, 418f

Logistics distribution, of random variables, 196–197

Logistics random variable, 197

Logit, 419

Lognormal distribution, 243

Lower confidence interval
 for difference in means of two normal distributions, 260–264, 261–262f, 267t
 for normal mean with unknown variance, 255–257, 256f
 for unknown mean, 249–251

- for unknown probability, 272t
- for variance of normal distribution, 260t
- Lower control limits
 - for exponentially weighted moving-average, 576–578, 578f
 - for fraction defective, 566–567
 - for mean control charts, 555–557, 556f
 - for moving-average, 572–573, 573t, 574f
 - for number of defects, 568–570
 - for variance control charts, 563–564, 564f
- M**
- Mann-Whitney test. *See* Rank sum test
- Marginal probability mass function, 98
- Markov's inequality, random variables and, 128–131
- Mass function. *See* Probability mass function
- Matrix notation
 - for multiple linear regression, 401–403
 - for polynomial regression, 400
- Maximum likelihood estimators, 235–247, 284, 500
 - of Bernoulli parameter, 237–240
 - of difference in means of two normal distributions, 260
 - evaluation of, 277–278
 - for exponential distribution in life testing, 593–594, 602–604
 - least squares estimators as, 365–366
 - for life distributions, 245–247
 - in logistic regression models, 418
 - for mean of exponential distribution, 272
 - of normal population, 242–244
 - of Poisson parameter, 240–241
 - in sequential testing for exponential distribution in life testing, 599, 601
 - for Weibull distribution in life testing, 608–610
 - weighted least squares estimators as, 392
- Mean. *See also* Population means; Sample mean
 - of Bernoulli random variable, confidence interval for, 268–272, 272t
 - of chi-square random variable, 192
 - confidence interval estimators of mean response, 377–379, 411–413
 - estimation of difference in means of two normal distributions, 260–268, 262–263f, 266–267f, 268t
 - of exponential distribution, confidence interval for, 272–273
 - for exponentially weighted moving-average, 575
 - grand, 462–463, 469
 - of hypergeometric random variables, 157–158
 - of least squares estimators, 362–364
 - for moving-average, 572
 - normal, confidence intervals for, 252–258, 256f
 - of normal population, hypothesis testing concerning, 299–317, 301f, 301f, 311t, 313f, 316f, 317t
 - case of known variance, 299–311, 301f, 304f, 311t
 - case of unknown variance, 311–317, 313f, 316f, 317t
 - of normal random variables, 169–170
 - permutation tests and, 634
 - of Poisson distribution, hypothesis testing for, 336–339
 - Poisson distribution with unknown value of, goodness of fit tests for, 499–501
 - population, 208–209, 209f
 - regression to, 372–376, 373f, 374f, 375t, 376f
 - testing equality of means of two normal populations, 318–326, 319t, 321f, 323f, 325t
 - case of known variance, 318–320, 319t
 - case of unknown and equal variance, 324–325, 325t
 - case of unknown variance, 320–324, 321f, 323f
 - paired *t*-test, 325–326
 - of uniform distribution, 244–245
 - of uniform random variables, 162
 - unknown
 - confidence intervals for normal mean with unknown variance, 252–258, 256f
 - control charts for, 557–562, 559t
 - estimates of, 247–252
- Mean control chart, 554–562, 556f, 564f
 - case of unknown, 557–562, 559t
- Mean life, maximum likelihood estimator of, 603–604
- Mean response
 - estimation of, 411–413
 - statistical inferences concerning, 377–379
- Mean square error
 - bootstrap method and, 627–628
 - of point estimators, 273–279

Median, sign test for, 523–524, 524f
 Memoryless, exponential random variables, 178
 Modal values, 22
 Mode, of density, 284
 Models, assessment of, 384–386, 385–386f
 Moment generating functions
 chi-square distribution, 188
 chi-square random variable, 188–190, 190f
 exponential random variables and, 177
 gamma distribution and, 186–188
 normal random variables and, 175
 of Poisson random variables, 149–150
 of random variables, 127–126
 Monte Carlo simulation, 255–257, 256f,
 622–623
 determining runs in, 643–644
 Moving-average control charts, 571–573, 573t,
 574f
 exponentially weighted, 573–578, 575f, 578f
 Multidimensional integrals, simulation of,
 255–257, 256f
 Multiple comparisons, of sample means,
 456–458
 Multiple linear regression, 400–416, 403t, 404f,
 405f, 406f, 410t, 413t, 414f, 415f
 Multiple probability distributions, equality
 testing of, 539–541
 Multiple regression equation, 358
 Multivariate normal distribution, 404
 Mutually exclusive, in sample space, 55

N

*Natural and Political Observations Made upon
 the Bills of Mortality*, 4–5, 4t, 5t
 Negatively correlated, 37
 Neyman, Jerzy, 7
 95 Percent confidence interval
 of difference in means of two normal
 distributions, 260–268, 252–253f
 for estimating unknown mean, 247–252
 for mean of exponential distribution,
 272–273
 of mean response, 378
 for normal mean with unknown variance,
 253–257, 254f
 for regression parameters, 372
 for unknown probability, 269–271
 95 Percent prediction interval, 416
 99 Percent confidence interval
 for estimating unknown mean, 250–251
 for unknown probability, 271
 90 Percent confidence interval
 of difference in means of two normal
 distributions, 265–268, 266–267f
 for variance of normal distribution, 259–260
 Nonparametric hypothesis tests, 521–545
 introduction to, 521
 runs test for randomness, 541–545, 543f
 sign test, 521–525, 524f
 signed rank test, 525–531, 526f
 two-sample problem, 531–535, 534f, 538f
 classical approximation and simulation,
 535–539
 equality of multiple probability
 distributions, 539–541
 Nonparametric inference, 207–208
 Nonrandom sample, 3
 Normal approximations, in permutation tests,
 633–637
 Normal data sets, 34, 31f, 32f, 34f
 Normal density function, 168, 168f, 188, 189f
 Normal distribution
 confidence interval for variance of, 259–260,
 260t
 estimation of difference in means of,
 260–268, 262–263f, 266–267f, 268t
 Normal equations
 in multiple linear regression, 401–403
 in polynomial regression, 397
 of regressions, 359–360
 Normal histograms, 33, 31f, 32f, 34f
 Normal mean, with unknown variance,
 confidence intervals for, 252–258, 256f
 Normal populations
 maximum likelihood estimator of, 242–244
 mean of
 hypothesis testing concerning, 299–317,
 301f, 304f, 311t, 313f, 316f, 317t
 testing equality of means of two normal
 populations, 318–326, 319t, 321f, 323f,
 325t
 sampling distributions from, 220–223
 joint distribution, 221–223
 sample mean distribution, 221
 variance of, hypothesis testing for, 327–329
 Normal prior, choosing of, 283–284
 Normal random variables, 168–177, 172f, 176f
 chi-square distribution, 188–193, 190f, 192f
 F-distribution, 195–196, 195f
 generation of, 642–643
 mean and variance of, 169–170
 normal density function, 168, 168f

- standard normal distribution and, 171–172, 172f, 175–177, 176f
- sums of, 175
- t -distribution, 193–195, 193f, 195f
- Notation
 - dot, in two-way analysis of variance, 461–462
 - for least squares estimators, 366
 - matrix
 - in multiple linear regression, 401–403
 - for polynomial regression, 400
- Null hypothesis, 298
 - permutation tests and, 631–633
- Number of defect control charts, 567–570, 569t
- O**
 - Observational study, 336
 - OC curve. *See* Operating characteristic curve
 - Odds for success, 419
 - Odds of event, 59
 - Ogives, 14t, 16–17, 16f, 18t
 - $100(1 - \alpha)$ Percent confidence interval of
 - difference in means of two normal distributions, 261–265, 268t
 - for estimating unknown mean, 250–251
 - for exponential distribution in life testing, 596
 - for mean of exponential distribution, 272
 - of mean response, 378
 - for normal mean with unknown variance, 252–258
 - for regression parameters
 - α , 377
 - β , 371–372
 - in sequential testing for exponential distribution in life testing, 599–600
 - for unknown probability, 269–271
 - for variance of normal distribution, 260t
 - $100(1 - \alpha)$ Percent confidence region, 263
 - $100(1 - \alpha)$ Percent prediction interval, 381, 416
 - One-sided Chebyshev's inequality, 29–30
 - One-sided critical region, 307
 - One-sided hypothesis tests
 - for mean of normal population, case of known variance, 306–310
 - for testing equality of means of two normal populations, 321
 - One-sided lower confidence interval
 - of difference in means of two normal distributions, 260–264, 261–262f, 267t
 - for normal mean with unknown variance, 255–257, 256f
 - for unknown mean, 249–251
 - for unknown probability, 272t
 - for variance of normal distribution, 260t
 - One-sided null hypothesis, sign test and, 524–525
 - One-sided t -tests, for mean of normal population with unknown variance, 314–317, 316f
 - One-sided upper confidence interval
 - for difference in means of two normal distributions, 261, 265–268, 266–267f
 - for normal mean with unknown variance, 254–255
 - for unknown mean, 249–251
 - for unknown probability, 272t
 - for variance of normal distribution, 260t
 - One-way analysis of variance, 446, 448–459, 452t, 453f, 456t
 - multiple comparisons of sample means, 456–458
 - with unequal sample sizes, 458–459
 - Operating characteristic (OC) curve, 303–304, 304f
 - for one-sided hypothesis testing for mean of normal population, 307–308
 - Out of control, 553, 555–557, 556f
 - Overlook probabilities, 74
- P**
 - Paired data sets, 34–36, 34t, 35f
 - sample correlation coefficient and, 37–40, 39f
 - Paired t -test, 325–326, 523
 - Parameter estimation, 235–284
 - approximate confidence interval for mean of Bernoulli random variable, 268–272, 272t
 - Bayes estimator, 236, 279–284
 - confidence interval for mean of exponential distribution, 272–273
 - of difference in means of two normal distributions, 260–268, 262–263f, 266–267f, 268t
 - interval estimates, 235, 247–260, 310
 - introduction, 235–236
 - of life distributions, 245–247
 - maximum likelihood estimators, 235–247, 260, 284
 - point estimator evaluation, 273–279

- Parameter estimation (*continued*)
 - for two-way analysis of variance, 460–463
 - for Weibull distribution in life testing, 610–612
- Parametric inference, 207–208
- Pearson, Egon, 7
 - The Pareto Distribution, 183–185
- Pearson, Karl, 6, 373, 496
- Permutation, 61–63
- Permutation tests, 630–638
 - implementation of, 631–632
 - normal approximations in, 633–637
 - null hypothesis and, 631–633
 - two sample, 637–638
- Pie chart, 12, 13–14f
- Point estimates, 235
- Point estimators
 - evaluation of, 273–279
 - for hypothesis testing, 299–300
 - of mean response, 377, 411
- Point prediction, 413
- Poisson distribution
 - hypothesis testing concerning mean of, 336–339
 - with unknown mean, goodness of fit tests for, 499–501
 - variance in, 395–396
- Poisson distribution function
 - computation of, 155–156
 - number of defects and, 567–570, 569t
- Poisson parameters
 - maximum likelihood estimator of, 240–241
 - testing of relationship between, 337–339
- Poisson probability mass function, 148–150, 149f, 154–155
- Poisson process, exponential random variables and, 181–183, 181f
- Poisson random variables, 148–156, 149f
 - binomial random variables and, 150–153
 - conditional probability and, 153–154
 - moment generating functions of, 149–150
 - probability mass function and, 148–150, 149f, 154–155
- Poisson, S.D., 148
- Polynomial regression, 397–400, 398f, 399f
- Pooled estimator, 266, 321
- Population distributions
 - empirical distribution and, 624
 - equality of, hypothesis testing for, 508–509
 - signed rank test for, 529–531
- Population means, 208–209, 209f
 - bootstrap method and, 623–624, 628–630
 - confidence interval for difference in, 456–458
 - control charts for, 571–581
 - cumulative sum, 579–581
 - exponentially weighted moving-average, 573–578, 575f, 578f
 - moving-average, 571–573, 573t, 574f
 - hypothesis testing of equality of, 446, 448–459, 452t, 453f, 456t
 - multiple, hypothesis testing of, 446–447
- Population median, sign test for, 523–524, 524f
- Population variance, 208–209
 - bootstrap method and, 623–625
- Populations
 - definition of, 207
 - samples and, 3
 - sampling distributions from
 - finite, 223–227
 - normal, 220–223
- Positively correlated, 37
- Posterior density function, 280, 284
- Power-function, of hypothesis test, 304
- Prediction interval
 - confidence interval v., 381
 - of future response, 379–381
 - of response at input level x_0 , 381
 - of response in multiple linear regression, 411–416, 413t, 414f, 415f
- Prior distributions, 279–284, 604–606
- Probability, 53–79
 - axioms of, 57–59, 59f
 - Bayes' formula, 68–75, 69f
 - Bernoulli random variables, 141–148
 - binomial random variables, 143–147
 - bootstrap method and, 629–630
 - central limit theorem, 210–219, 212–215f, 216f
 - chi-square distribution and, 187
 - conditional, 65–68, 66f, 106
 - continuous random variable and, 94
 - counting and, 60–65
 - of defects, 329–337
 - distribution function and, 91–92
 - events, 54–55
 - independent, 75–79, 78f
 - expectation, 107–111, 111f
 - exponential random variables and, 179–180
 - fraction defective, 565–567
 - introduction to, 53–54
 - overlook, 74

- Poisson random variables and, 148–156, 149f
 - of random variables, 89–90
 - rank sum tests, 532–533
 - sample space, 54–55
 - with equally likely outcomes, 59–65
 - signed rank test and, 527–529
 - of uniform random variables, 160–161, 161f
 - unknown, confidence interval for, 268–272, 272t
 - Venn diagram and algebra of events, 56–57, 56f, 57f
 - Probability density function, 93–94
 - cumulative distribution function and, 94–95, 94f
 - exponential random variables and, 177
 - joint, 99–101
 - of uniform random variables, 160–161, 161f
 - updated, 280
 - Probability distribution
 - of estimator of mean response, 377–378
 - of sample, goodness of fit tests for, 489–499, 493t, 498f
 - Probability distribution function
 - joint cumulative, 96, 103–104
 - Poisson, 148–156
 - of populations, 207
 - random variable and expectation, 111–113
 - signed rank test for, 529–531
 - Probability mass function, 92–93, 93f, 245–247
 - Bernoulli random variables, 142–144, 143f
 - binomial random variables, 142–144, 143f
 - central limit theorem and, 212–215f
 - conditional, 105–106
 - discrete random variables, 92–93, 638–640
 - expectation of, 107–108
 - hypergeometric random variables, 156–157
 - individual and joint, 96–99, 98t
 - marginal, 98
 - Poisson, 148–150, 149f, 154–155
 - Poisson random variables, 148–150, 149f, 154–155
 - Probability models, inferential statistics and, 2–3
 - Probability theory, statistics and, 5–6
 - Probit model, 419
 - Pseudo random numbers, 257, 620
 - p*-value
 - for determining independence of characteristics of population member, 504–505
 - for goodness of fit tests when all parameters are specified, 491, 494–499, 498f
 - for goodness of fit tests when some parameters are unspecified, 500–501
 - for hypothesis testing
 - in Bernoulli populations, 330–333, 335
 - of equality of population means, 452, 453f
 - of mean of normal population, 302, 307–310, 311t, 313, 315–317, 316f, 317t
 - of mean of Poisson distribution, 337–338
 - with multiple linear regression, 411
 - of regression parameters, β , 370
 - of regression to mean, 374
 - of variance of normal population, 327, 329
 - for Kolmogorov–Smirnov goodness of fit test, 513
 - for one-sided hypothesis testing for mean of normal population, 307–310
 - permutation tests for, 630–638
 - normal approximations in, 633–637
 - two sample, 637–638
 - rank sum test and, 533–535, 534f
 - in sequential testing for exponential distribution in life testing, 601–602
 - signed rank test for, 527–529
 - simulation for approximation of, 496–499, 498f
 - for testing equality of means of two normal populations, 321–324, 323f, 325t, 326
 - in two-way analysis of variance, 467, 467t, 468f, 473–474, 475t, 476f
- Q**
- Quality control, 553–581
 - fraction defective control charts, 565–567
 - introduction to, 553–554
 - mean control chart, 554–562, 556f, 564f
 - number of defect control charts, 567–570, 569t
 - population mean control charts, 571–581
 - cumulative sum, 579–581
 - exponentially weighted moving-average, 573–578, 575f, 578f
 - moving-average, 571–573, 573t, 574f
 - variance control chart, 562–565, 564f

R

- Random error, in response to input variable, 357–358, 361
- Random numbers, 620–623
 - definition of, 163, 163t
 - generation of, 620–622
 - Monte Carlo simulation approach, 622–623
 - pseudo, 620
 - use of, 164–166, 166f
- Random sample, 3, 207, 223
 - runs test for, 541–545, 543f
- Random variables, 89–92. *See also specific random variables*
 - Bernoulli and binomial, 141–148, 143f, 148f
 - central limit theorem, 210–219, 212–215f
 - Chebyshev's inequality, 128–131
 - continuous, 93–94, 640–643
 - density function and, 112
 - discrete, 91–92, 638–640
 - distribution function and, 91–92, 624
 - entropy of, 110
 - expectation of function of, corollary of, 114–115
 - expected value of sums of, 115–118
 - exponential, 177–182, 181f, 641–642
 - gamma distribution of, 186–188, 189f
 - generation of, 497, 627, 638–643
 - hypergeometric, 156–160
 - indicator, 90–91
 - jointly distributed, 95–107, 98t
 - conditional distributions, 104–107
 - independent, 101–104
 - logistics distribution, 196–197
 - Markov's inequality, 128–130
 - moment generating functions, 127–128
 - normal, 168–177, 172f, 176f, 642–643
 - chi-square distribution, 188–193, 190f, 192f
 - F -distribution, 195–196, 195f
 - t -distribution, 193–195, 193f, 194f
 - Poisson, 148–156, 149f
 - probability distribution function and expectation, 111–113
 - types of, 92–95, 93f, 94f
 - uniform, 160–168, 161f, 163t, 166f
 - variance of, 118–120, 162, 169–170, 192, 222, 447–448
 - variance of a sum of, 123–126
 - weak law of large numbers, 130
- Rank sum test, 521, 531–541, 534f, 538f
 - distribution function of, 531
 - Kruskal–Wallis* test, 540
 - probability and, 532–533
 - p -value and, 533–535, 534f
 - classical approximation and simulation, 535–539, 538f
- Rate of distribution, 590
- Rayleigh density function, 591
- Recursive formula, mean control chart and, 559–560, 559t
- Referents, 334
- Regression, 357–420
 - analysis of residuals and assessing models, 384–386, 385–386f
 - coefficient of determination and sample correlation coefficient, 382–384
 - distribution of least squares estimators, 361–367, 368f, 369f
 - history of, 6
 - introduction, 357–358, 358f
 - least squares estimators of regression parameters, 359–361, 361f, 362f
 - logistic regression models for binary output data, 416–420, 418f
 - to mean, 372–376, 373f, 374f, 375t, 376f
 - multiple linear, 400–416, 402t, 404f, 405f, 406f, 410t, 414f, 415f
 - predicting future responses, 411–416, 413t, 414f, 415f
 - polynomial, 397–400, 398f, 399f
 - statistical inferences about regression parameters, 367–382, 373f, 374f, 375t, 376f
 - α , 376–377
 - β , 368–376, 373f, 374f, 375t, 376f
 - mean response, 377–379
 - prediction interval of future response, 379–381
 - summary of distribution results, 381–382
 - transforming to linearity, 387–390, 388f, 389f, 389t, 390t
 - use of dummy variables, 416–418
 - weighted least squares, 390–396, 395f
- Regression coefficients, 358, 397
- Regression fallacy, 376
- Regression parameters
 - least squares estimators of, 359–361, 361f, 362f
 - statistical inferences about, 367–382, 373f, 374f, 375t, 376f
 - α , 376–377
 - β , 368–376, 373f, 374f, 375t, 376f
 - mean response, 377–379

prediction interval of future response, 379–381
 summary of distribution results, 381–382
 Rejection, of hypothesis. *See* Hypothesis testing
 Relative frequency tables and graphs, 10–14, 13–14f, 13t, 16
 Residuals, 364–366
 analysis of, 384–386, 385–386f
 in multiple linear regression, 408–409, 411
 standardized, 385–386, 385–386f
 Response variable, 357–358
 prediction interval of future response, 379–381
 in multiple linear regression, 411–416, 413t, 414f, 415f
 variation in, 382–384
 with input variable, 390–396, 395f
 Robustness, of hypothesis test, 311
 Row factors
 hypothesis testing for, 464–468, 467t, 468f
 in two-way analysis of variance, 460
 column factor interaction with, 446, 469–476, 475t, 476f
 deviation from grand mean due to, 462
 Row sum of squares, 466, 467t
 Run, 541
 Runs test for randomness, 521, 541–542, 543f

S

Sample
 definition of, 207
 populations and, 3
 Sample 100 p percentile, 24–25
 Sample correlation coefficient, 37–41, 39f
 association v. causation, 40–41
 coefficient of determination and, 382–384
 properties of, 37–38, 41
 Sample mean, 17–20, 22
 central limit theorem for, 216–218
 distribution of, with chi-square random variables, 222
 for exponential random variables, 218–219, 219f
 for independent random variables, 2, 15
 multiple comparisons of, 456–458
 of normal data set, 31
 for normal population, 220–221
 population, 208–209, 209f
 sample variance distribution with, 221–223
 Sample median, 20–22, 31
 Sample mode, 22
 Sample percentiles, 24–25
 Sample quartiles, 25–27, 27f
 Sample size, one-way analysis of variance with unequal sample sizes, 458–459
 Sample spaces, 54–55
 having equally likely outcomes, 59–65
 Sample standard deviation, 24, 219–220
 Sample variance, 22–24, 219–220
 for normal population, 220
 sample mean distribution with, 221–223
 Sampling, 207
 Sampling distributions
 form finite populations, 223–227
 form normal populations, 220–223
 joint distribution, 221–223
 sample mean distribution, 221
 Scatter diagram, 34–36, 36f, 358, 358f, 373, 385–386, 385–386f, 397
 Second quartile, 25–27
 Selection, of normal prior, 283–284
 Sequence of interarrival times, 182
 Sequential testing, for exponential distribution in life testing, 598–602, 598f
 Sign test, 521–525, 524f
 Bernoulli random variables, 522
 binomial random variables, 522
 one-sided null hypothesis and, 524–525
 paired t -test v., 523
 for population median, 523–524, 524f
 Signed rank test, 521, 525–531, 526f
 for distribution function, 525, 526f
 for probability distribution function, 529–531
 for p -value, 527–529
 Significance level, 298–299
 Significance level α test
 for determining independence of characteristics of population member, 503–505
 for goodness of fit tests when all parameters are specified, 496
 for hypothesis testing
 in Bernoulli populations, 329–333
 of equality of population means, 452, 459
 of mean of normal population, 300–302, 307–309, 311t, 312, 314–316, 317t
 of mean of Poisson distribution, 336–337
 of regression to mean, 374
 of variance of normal population, 329
 for Kolmogorov–Smirnov goodness of fit test, 513–514

Significance level α test (*continued*)
 for testing equality of means of two normal populations, 318–321, 324–325, 325t
 in two-way analysis of variance, 467, 467t, 472–474, 475t

Simple hypothesis, 298

Simple regression equation, 358, 362f, 369f
 assessment of, 384–386, 385–386f

Simulation
 for determination of critical region, 496–499, 498f
 of single and multidimensional integrals, 251–253, 256f

Simulation run, 623
 in Monte Carlo study, 643–644

Simulation testing, for exponential distribution
 in life testing, 592–598, 602–604

Single integrals, simulation of, 255–257, 256f

Skewed data set, 31, 32f

Skewed random variables, 142, 143f

Standard deviation
 definition of, 121
 mean control chart and, 560–561
 variance control chart, 562–563

Standard logistic, 197

Standard normal distribution, 171–172, 172f, 175–177, 176f
 central limit theorem and, 217
 of mean control chart, 555
 t -distribution and, 192–195, 193f, 194f

Standard normal random variable, 221
 central limit theorem and, 210, 216–217

Standardized residuals, 385–386, 385–386f

Stationary increment assumption, 181–182

Statistical analysis, 1

Statistical inferences, about regression
 parameters, 367–382, 373f, 374f, 375t, 376f
 α , 376–377
 β , 368–376, 373f, 374f, 375t, 376f
 mean response, 377–379
 prediction interval of future response, 379–381
 summary of distribution results, 381–382

Statistical theory, 1

Statistics
 application of, 6–7
 definition of, 1, 6–7, 6t, 207–208
 descriptive, 1–2
 history of, 3–7, 4t, 5t, 6t

inferential, 2–3
 introduction to, 1–7

Stem and leaf plots, 16–17, 18t
 of normal data set, 33
 sample mean and, 21
 sample median and, 21

Subjective interpretation, probability, 53

Success, odds for, 419

Sum of squares
 column, 467t
 error, 465, 467t
 row, 466, 467t
 between samples, 451–452, 454, 456t, 459
 within samples, 450, 454, 456t, 459
 in two-way analysis of variance with
 interaction, 471–474, 475t

Sum of squares identity, 453–454

Sum of squares of residuals, 364–366, 408–409, 411

Survival rate, 245–246

T

t -density function, 193, 193f, 252, 253f

t -distribution, 193–195, 193f, 194f, 223

Test statistic
 for determining independence of
 characteristics of population member, 503–505
 for goodness of fit tests when all parameters
 are specified, 490–492, 494–496
 for goodness of fit tests when some
 parameters are unspecified, 500–501
 for hypothesis testing
 in Bernoulli populations, 330
 of equality of population means, 451–452, 455, 456t
 of mean of normal population, 302, 307, 309, 315t, 312–315, 317t
 of regression parameters, 370–371
 of regression to mean, 374
 of variance of normal population, 327, 329

Kolmogorov–Smirnov, 510–514, 511f

for one-sided hypothesis testing for mean of
 normal population, 307, 309

for testing equality of means of two normal
 populations, 320–322, 324, 325t, 326

for testing independence in contingency
 tables, 506–507, 506f

in two-way analysis of variance, 467, 467t

Testing. *See* Goodness of fit tests; Hypothesis testing; Life testing

- Tests of independence
 - in contingency tables, 501–505
 - in contingency tables having fixed marginal totals, 505–510, 506f
- Third quartile, 25–27
- Threshold model, 420
- Ties
 - rank sum test and, 531
 - signed rank test and, 530–531
- T*-method, 456–458
- Total-time-on-test statistic, 594–595, 604
- T*-random variable, 264
- Transformation, to linearity, 387–390, 388f, 389f, 389t, 390t
- Treatment group, 164
- Tree diagram, random numbers and, 166, 166f
- t*-tests, 311–317, 313f, 316f, 317t
 - one-sided, 314–317, 316f
 - paired, 325–326
 - p*-value of two-sample, 323f
 - two-sided, 311–313, 313f
- Two sample permutation tests, 637–638
- Two-sample problem, 521, 531–539, 534f, 538f
 - distribution function of, 531
 - in life testing, 606–608
 - probability and, 532–533
 - p*-value and, 533–535, 534f
 - classical approximation and simulation, 535–539, 538f
- Two-sided confidence interval, 248, 251
 - of difference in means of two normal distributions, 261–268, 262–263f, 266–267f
 - for normal mean with unknown variance, 255–257, 256f
 - for unknown probability, 272t
- Two-sided *t*-tests, for mean of normal population with unknown variance, 311–313, 313f
- Two-way analysis of variance, 446
 - hypothesis testing for, 464–468, 467t, 468f
 - with interaction, 446, 469–476, 475t, 476f
 - introduction and parameter estimation, 460–463
- Type I errors, 298, 300
- Type II errors, 298, 302–305, 304f
- U**
- Unbalanced case, in one-way analysis of variance, 459
- Unbiased estimators, 274–279
- Uniform distribution, estimating mean of, 244–245
- Uniform random variables, 160–168, 161f, 163t, 166f
 - mean and variance of, 162
 - probability density function of, 160–161, 161f
 - random numbers, 166–168
- Union
 - of sample space, 55
 - in Venn diagram, 56–57, 56f, 57f
- Unit normal distribution. *See* Standard normal distribution
- Unknown mean
 - confidence intervals for normal mean with unknown variance, 252–257, 256f
 - estimates of, 247–252
- Unknown parameters. *See* Parameter estimation
- Unknown probability, confidence interval for, 268–272, 272t
- Unknown variance
 - confidence intervals for normal mean with, 252–257, 256f
 - hypothesis testing for mean of normal population with, 311–317, 313f, 316f, 317t
 - testing equality of means of two normal populations with, 320–325, 321f, 323f, 325t
- Updated probability density function, 280
- Upper confidence interval
 - for difference in means of two normal distributions, 255, 265–266, 266–267f
 - for normal mean with unknown variance, 254–255
 - for unknown mean, 249–251
 - for unknown probability, 272t
 - for variance of normal distribution, 260t
- Upper control limits
 - for exponentially weighted moving-average, 576–578, 578f
 - for fraction defective, 566–567
 - for mean control charts, 555–557, 556f
 - for moving-average, 572–573, 573t, 574f
 - for number of defects, 568–570
 - for variance control charts, 563–564, 564f

V

Variance, 118–121. *See also* Analysis of variance;
 Population variance; Sample variance
 of chi-square random variable, 192, 222,
 447–448
 covariance, 121–122
 definition of, 119
 distribution of, with chi-square random
 variables, 222
 estimators of, 447–448
 for one-way analysis of variance, 446,
 448–459, 452t, 453f, 456t
 in two-way analysis of variance, 464–467
 in two-way analysis of variance with
 interaction, 469–476, 475t, 476f
 for exponentially weighted moving-average,
 575–576
 of hypergeometric random variables,
 157–158
 of independent random variables, 222
 of indicator random variable, 119–121
 known
 equality of means of two normal
 populations with, 318–320, 319t
 hypothesis testing for mean of normal
 population with, 299–311, 301f, 304f,
 311t
 of least squares estimators, 362–364,
 405–407
 for moving-average, 572
 of normal distribution, confidence interval
 for, 259–260, 259t
 of normal population, hypothesis testing for,
 327–329
 of normal random variables, 169–170

permutation tests and, 634–635
 population, 208–209
 of random variables, 118–121, 123–126,
 162, 169–170, 193, 222, 447–448
 in response to input variable, 382–384,
 390–396, 395f
 sample, 22–24, 219–220
 of a sum of random variables, 124–126
 of uniform random variables, 162
 unknown
 confidence intervals for normal mean with,
 252–257, 256f
 equality of means of two normal
 populations with, 320–324, 321f, 323f
 hypothesis testing for mean of normal
 population with, 311–317, 313f, 316f,
 317t
 unknown and equal, testing equality of
 means of two normal populations with,
 324–325, 325t
 Variance control chart, 562–565, 564f
 Venn diagram, 56–57, 56f, 57f
 probability axioms and, 58–59, 59f

W

Weak law of large numbers, 130
 Weibull density function, 608, 609f
 Weibull distribution, in life testing, 608–612,
 609f
 Weighted average, 19–20
 Weighted least squares estimators, 390–396,
 395f
 Wilcoxon test. *See* Rank sum test
 Within samples sum of squares, 450, 454, 456t,
 459