

Measure Theory

Mark Ibrahim

*based on Principles of Real Analysis by Aliprantis and Burkinshaw

Contents

1 Preliminaries	2
2 Algebras and Measures	2
2.1 Semirings and Sigma-algebras of Sets (section 12)	2
2.1.1 semirings	2
2.1.2 algebras	2
2.2 Measures on Semirings (section 13)	3
2.2.1 Examples of Measures on S	3
2.3 Outer Measures (section 14)	3
2.4 Outer Measures generated by a measure (section 15)	4
2.4.1 Cantor Set	4
2.5 Lebesgue Measure (section 18)	4
2.5.1 What are the Borel sets in the reals?	4
2.5.2 Regular Borel Measure	5
3 Integration: functions	5
3.1 Measurable Functions (section 16)	5
3.1.1 Sequences of Functions and Measurability	5
3.1.2 Ergov's Theorem (16.7)	5
3.2 Simple and step functions (section 17)	6
3.2.1 sigma-finite	6
4 Lebesgue Integral	6
4.1 Upper Functions (section 21)	6
4.2 Integrable Functions (section 22)	7
4.2.1 How does integrable relate to other properties?	7
4.2.2 When is f integrable?	7
4.2.3 Big: Levi, Fatou, and Lebesgue Dominated Convergence	7
4.3 Riemann Integrals (section 23)	8
4.4 Product Measures and Iterated Integrals (section 26—only a sketch)	8
5 Function Spaces (Chapter 5)	9
5.1 norms on vector spaces (section 27)	9
5.2 Linear Operators (section 28)	9
5.2.1 Dual Spaces	9
5.3 Lp Spaces (section 31)	10
5.3.1 Essentially Bounded Functions	10
5.3.2 What is in Lp?	10
5.3.3 Dense Functions in Lp	11

5.3.4	Counting Measure and lp	11
6	Hilbert Spaces (Chapter 6)	11
6.1	Inner Product Spaces (section 32)	11
6.1.1	Useful Inequalities	12
6.2	Hilbert Spaces (section 33)	12
6.2.1	Orthogonal Complement and Spans	12
6.3	Orthonormal Bases	12
6.4	Fourier Analysis	13
6.4.1	Best Approximation Theorem	13
6.4.2	Fourier coefficients	13
6.4.3	Riesz-Fischer	13
6.4.4	Fourier on L2	13
6.5	Risz-Representation Theorem	14
7	Miscellaneous Examples	14
8	Questions	14

1 Preliminaries

a function $f : A \rightarrow B$ is **continuous**
 $\iff f^{-1}(\text{open set})$ is an open set.

a bounded sequence a_n has a lim sup
defined as $\lim_{N \rightarrow \infty} \sup\{a_N, a_{N+1}, \dots\}$
"largest tail"

a_n converges if $\limsup = \liminf$.

A **Hausdorff** topological space (T2 space) is a topological space where any two points can be separated by open sets.

$$\max\{a, b\} = \frac{a+b}{2} + \frac{|a-b|}{2}.$$

union of countably sets is countable.

2 Algebras and Measures

2.1 Semirings and Sigma-algebras of Sets (section 12)

2.1.1 semirings

a collection S of subsets of a set X is called a **semiring** if

1. $\emptyset \in S$,
2. $A \cap B \in S$, and
3. $A - B = C_1 \cup \dots \cup C_n$ for $C_1, \dots, C_n \in S$.

Any countable union in S can be written as a countable **disjoint union**.

e.g., $S = \{[a, b) \mid a \leq b \in \mathbb{R}\}$ is a semiring, not an algebra.

* note $[a, a) = \emptyset$.

2.1.2 algebras

a nonempty collection S of subsets of a set X is an **algebra** if

1. $A \cap B \in S$
2. and $A^c \in S$.

Nice properties of algebras are:

- $\emptyset, X \in S$
- S is closed under finite unions and finite intersections as well as subtraction

¹(X, T) is a topological space with a set X and subsets T if $\emptyset, X \in T$, and T is closed under unions (even uncountable), finite intersections.

a σ -algebra is an algebra that is closed under countable unions.

Borel sets of a topological space (X, T) is a σ -algebra generated by the open sets.

2.2 Measures on Semirings (section 13)

A function μ from a semiring S to $[0, \infty]$ is a **measure on S** if

1. $\mu(\emptyset) = 0$
2. countably additive: $\mu(\cup_{n=1}^{\infty} A_n) = \sum_{n=1}^{\infty} \mu(A_n)$.

* $\cup_{n=1}^{\infty} A_n$ must be in S and each is disjoint.

* don't need to check if S is a σ -algebra!

• If $A \subseteq B$, ($A, B \in S$), then $\mu(A) \leq \mu(B)$.

Alternatively, can show μ is a measure if and only if "squeeze"

1. $\mu(\emptyset) = 0$
2. $\sum_{i=1}^n \mu(A_i) \leq \mu(A)$ if $\cup_{i=1}^n A_i \subseteq A$ and A_i are disjoint.
3. $\mu(B) \leq \sum_{n=1}^{\infty} \mu(B_n)$, "subadditive" if $B \subseteq \cup_{n=1}^{\infty} B_n$.

2.2.1 Examples of Measures on S

- **Counting Measure** $\mu(A) = |A|$
- **Dirac Measure** Fix $a \in X$, $\mu_a(A) = 0$ if $a \notin A$, else 1.
- **Lebesgue Stieltjes** For $f : \mathbb{R} \rightarrow \mathbb{R}$, increasing, left continuous and $S = \{[a, b) | a \leq b \in \mathbb{R}\}$, $\mu([a, b)) = f(b) - f(a)$.
- **Lebesgue Measure on S** , denoted λ is defined by $\lambda([a, b)) = b - a$.

2.3 Outer Measures (section 14)

an **outer measure** is a function $\bar{\mu} : P(X) \rightarrow [0, \infty]$ such that

1. $\bar{\mu}(\emptyset) = 0$
2. if $A \subseteq B$, $\bar{\mu}(A) \leq \bar{\mu}(B)$
3. countably subadditive: $\bar{\mu}(\cup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} \bar{\mu}(A_n)$

*an outer measure is not always a measure!

A subset E of X is **measurable** if for all $A \subseteq X$,

$$\bar{\mu}(A) = \bar{\mu}(A \cap E) + \bar{\mu}(A \cap E^c)$$

A nicer equivalent way to show E is measurable is by considering all A in S with $\mu^*(A) < \infty$ and showing

$$\mu(A) \geq \mu^*(A \cap E) + \mu^*(A \cap E^c)$$

Nice Properties

- every A in S is μ^* -measurable
- if $\bar{\mu}(E) = 0$, E is measurable
- for E_i measurable and any $A \subseteq X$,
$$\bar{\mu}(\cup_{i=1}^n A \cap E_i) = \sum_{i=1}^n \bar{\mu}(A \cap E_i)$$

the collection of measurable subsets is denoted by Λ . This collection is a σ -algebra!

Remarkably, the outer measure $\bar{\mu}$ restricted to Λ is a measure!

2.4 Outer Measures generated by a measure (section 15)

The outer measure μ^* generated by a measure μ is defined for any subset A of X ,

$$\mu^*(A) = \inf \left\{ \sum_{n=1}^{\infty} \mu(A_n) : A \subseteq \bigcup_{n=1}^{\infty} A_n \text{ for } A_n \in S \right\}$$

μ^* is called the Cathéodory extension of μ . By convention $\mu^*(A) = \infty$ if no cover exists in S .

On semiring S , $\mu^* = \mu$.

For E_n measurable, if $E_n \uparrow E$, then $\mu^*(E_n) \uparrow \mu^*(E)$. * $E_n \uparrow E$ means:

- 1) $E_1 \subseteq E_2 \subseteq \dots$
- 2) $\bigcup_{n=1}^{\infty} E_n = E$

* note E must be measurable since it's the union of measurable sets

For B_n measurable with $\mu^*(B_n) < \infty$, if $B_n \downarrow B$, then $\mu^*(B_i) \downarrow \mu^*(B)$.

a measure space is **finite** if $\mu^*(X) < \infty$.

For X a **finite measure** space E is measurable, if and only if

$$\mu^*(E) + \mu^*(E^c) = \mu^*(X)$$

For all $A \subseteq X$, there is a measurable set E such that $A \subseteq E$ and $\mu^*(A) = \mu^*(E)$.

2.4.1 Cantor Set

Cantor set $C = \bigcap_{n=1}^{\infty} c_n$, where

$$c_1 = [0, 1] - (1/3, 2/3)$$

$$c_2 = c_1 - ((1/9, 2/9) \cup (7/9, 8/9))$$

each c_n is closed, because it's a closed set minus open sets.

- C has measure 0
- $|C| = |\mathbb{R}|$

- every point of C is an accumulation point of C

Vitali set is an example of a **non-measurable** subset of \mathbb{R} .

2.5 Lebesgue Measure (section 18)

Outer Lebesgue measure λ^* is defined as $\lambda^*(A) = \inf \left\{ \sum_{i=1}^{\infty} \lambda(a_i, b_i) : A \subseteq \bigcup_{i=1}^{\infty} (a_i, b_i) \right\}$

* note $\lambda(a, b) = b - a$.

* often, we say Lebesgue measure instead of outer Lebesgue measure.

By result about $E_n \uparrow E$ from section 15, we can show (a, b) , $[a, b]$, and $(a, b]$ are all measurable with same measure.

$E \subseteq \mathbb{R}$ is **Lebesgue measurable** \iff there is open $O \subseteq \mathbb{R}$ for each ϵ such that $E \subseteq O$ and $\lambda(O - E) < \epsilon$.

Every Borel set in \mathbb{R} is λ -measurable

2.5.1 What are the Borel sets in the reals?

By definition, it's the σ -algebra generated by open sets in \mathbb{R} . (Borel σ -algebra is generated by intervals of the form $(-\infty, a]$, for $a \in \mathbb{Q}$).

Borel sets contain:

- all closed sets
- union of all open sets or closed sets
- intersection of all open/closed sets

* we can write any open set in \mathbb{R} as disjoint countable union of open intervals!

2.5.2 Regular Borel Measure

For X , a Hausdorff topological space and B the borel sets in X , a measure μ on B is called a **regular borel measure** if

1. $\mu(K) < \infty$ if K is compact
2. for B a borel set, $\mu(B) = \inf\{\mu(O) | O \text{ is open } B \subseteq O\}$
3. for O open, $\mu(O) = \sup\{\mu(K) | K \text{ is compact and } K \subseteq O\}$
1. λ is a regular borel measure
2. Dirac measure is a regular borel measures
3. Counting measure is not for example $[0, 1]$ is compact, but has infinite measure
4. any **translation invariant** regular borel measure on \mathbb{R} is $c\lambda$ for some $c \in \mathbb{R}^+$

3 Integration: functions

3.1 Measurable Functions (section 16)

a relation holds **almost everywhere** if set where it fails has measure 0.

$f : X \rightarrow \mathbb{R}$ is a **measurable function** if

- $f^{-1}(O)$ is measurable, for all open sets O
- $f^{-1}(a, \infty)$ is measurable, for all a in \mathbb{R}

If $f, g : X \rightarrow \mathbb{R}$, $f = g$ **almost everywhere** and f is measurable, then g is measurable too!

"= a.e. means measurability carries over"

² $f^+ = f(x)$ if $f(x) \geq 0$ or 0 otherwise.

If $f, g : X \rightarrow \mathbb{R}$ are **measurable** then $\{x \in X | f(x) > g(x)\}$ is measurable.

Sum, product, constant multiple, $\|$, max, and f^{+2} of measurable functions is also measurable!

3.1.1 Sequences of Functions and Measurability

recall (from analysis): $f_n \rightarrow f$ **uniformly** means $|f_n(x) - f(x)| < \epsilon$ for all x if you go out far enough in the sequence.

Key Theorem: If $f_n \rightarrow f$ **uniformly** and f_n are continuous, then f is continuous.

We can define \limsup (\liminf) for any **bounded** sequence.

For a sequence of measurable functions $\{f_n\}_{n=1}^{\infty}$

- If $f_n \rightarrow f$ a.e., then f is measurable func.
- If $\{f_n\}_{n=1}^{\infty}$ is bounded, then \limsup is a measurable function (so is \liminf)

A sequence of functions, $\{f_n\}_{n=1}^{\infty}$ ($f_n : X \rightarrow \mathbb{R}$) converges **almost uniformly** on X if for any ϵ , there exists a measurable set F where $\mu(F) < \epsilon$ and $\{f_n\} \rightarrow f$ **uniformly** on $X - F$.

If $f_n \rightarrow f$ **almost uniformly** on X and $\mu(X) < \infty$ then, $|f_n(x) - f(x)| < \epsilon$ for all $n > \text{some } N \in \mathbb{N}$, and all x in a set J where $\mu(J^c) < \delta$.

3.1.2 Ergov's Theorem (16.7)

If $f_n \rightarrow f$ **almost uniformly** on X , then $f_n \rightarrow f$ pointwise **almost everywhere** on X .

Also, if $\mu(X) < \infty$ and $f_n \rightarrow f$ pointwise on X , then $f_n \rightarrow f$ uniformly on X .

counter example: if $\mu(X)$ is not finite, consider $X = \mathbb{R}$, $\mu = \lambda$ and $f_n = \chi_{[n, n+1]}$. Then, $f_n \rightarrow 0$, but not almost uniformly

3.2 Simple and step functions (section 17)

nice properties of χ_A

- $A \subseteq B \iff \chi_A \leq \chi_B$
- $\chi_{A \cap B} = \chi_A \chi_B$ (equivalently $\min\{\chi_A, \chi_B\}$)
- $\chi_{A \cup B} = \chi_A + \chi_B - \chi_{A \cap B} (= \max\{\chi_A, \chi_B\})$

a measurable function $f : X \rightarrow \mathbb{R}$ is a **simple function** if it takes on finitely many values.

the standard representation of a simple function is

$$\sum_i^n a_i \chi_{A_i}$$

where a is are distinct nonzero outputs and A inputs

If each A_i has finite measure, then f is called a **step function**.

The **integral** of a step function ϕ is

$$\int \phi du = \sum_i^n a_i \mu^*(A_i)$$

*it turns out any representation, even when A_i are not disjoint (or a_i distinct) yield the same integral value

addition and scalar multiplication can be split over integrals as expected.

If $\phi \geq \psi$ a.e., then $\int \phi \geq \int \psi$

*holds if $\psi = 0$ or \geq is =

If ϕ_n is a **sequence of step functions**

with $\phi_n \downarrow 0$ a.e., then $\int \phi_n \downarrow 0$.

(similarly if $\phi_n \uparrow \psi$ a.e.)

*careful, $\uparrow \psi$, but $\downarrow 0$

*also $\phi_n \rightarrow \phi$ isn't good enough!

If $\phi_n \uparrow f$ a.e. and $\psi_n \uparrow f$ a.e., then

$$\int \phi_n = \int \psi_n \text{ as } n \rightarrow \infty$$

We can show A is **measurable** if we can find step functions $\phi_n \uparrow \chi_A$. In this case, $\mu^*(A) = \lim \int \phi_n$

For any measurable $f \geq 0$, there exists **simple** ψ_n such that

$$0 \leq \psi_n \uparrow f$$

3.2.1 sigma-finite

X is a σ -**finite measure space** if there exists E_i such that $\cup_{i=1}^\infty E_i = X$, $\mu(E_i) < \infty$, and $E_1 \subseteq E_2 \subseteq \dots$

Who cares? Well if X is σ -finite then for a **measurable** $f \geq 0$ a.e., then there exists **step** $\phi_n \uparrow f$ a.e.

4 Lebesgue Integral

4.1 Upper Functions (section 21)

$f : X \rightarrow \mathbb{R}$ is an **upper function** if there exist step ϕ_n such that

- $\phi_n \uparrow f$ a.e.
- $\lim \int \phi_n du < \infty$

ϕ_n is called a **generating sequence** for f .

* all step functions are upper functions
* f upper does **not** imply $-f$ is upper

The integral of f an **upper function** is defined as

$$\int f du = \lim \int \phi_n du$$

* the value is independent of our choice of ϕ_n because if any other $\psi_n \uparrow f$ too, then $\int \phi_n = \int \psi_n$ as $n \rightarrow \infty$

sums, scalar multiples, maxes of upper functions are upper functions.

If $f \geq g$ a.e. (both upper) then $\int f \geq \int g$ (same for $g = 0$)

If a **sequence of upper functions** $f_n \uparrow f$ a.e. and $\lim \int f_n < \infty$ then f is upper and $\int f = \lim \int f_n$ (similarly if $f_n \downarrow 0$)

4.2 Integrable Functions (section 22)

a function f is **integrable** if $f = u - v$, both upper functions.

We define $\int f$ as $\int u - \int v$

* well-defined no matter the representation of f

4.2.1 How does integrable relate to other properties?

- **upper** functions are integrable
- **step** functions are integrable (b/c step are upper)
- integrable implies **measurable**
 - measurable does **not** imply integrable
 - e.g., constant functions are measurable, but only integrable when $\mu(X) < \infty$.

Canonical way to write integrable

$$f = f^+ - f^-$$

b/c: both f^+ and f^- are upper if f is integrable

4.2.2 When is f integrable?

If integrable $f = g$ a.e., then g is integrable (and integrals are equal).

sums, scalar multiples, max, || of integrable are integrable.

* $|f|$ integrable does **not** imply f is integrable.

If f is measurable and $h \leq f \leq g$ a.e. for h, g integrable, then f is **integrable**.

“measurable sandwiched between integrable is integrable”

nice properties of f integrable:

- if $f \geq 0$ a.e. then f is **upper**
- $A = \{x | |f(x)| \geq \epsilon\}$ has **finite measure** (A is also measurable)
b/c: $|f|$ is measurable so $|f|^{-1}(\epsilon, \infty)$

For f, g integrable,

1. $\int |f| = 0 \iff f = 0$ a.e.
2. If $f \geq g$ a.e., then $\int f \geq \int g$
3. $\int |f| \geq \left| \int f \right|$

If E is **measurable**, f is **integrable**, then

$$\int_X f = \int_E f + \int_{X-E} f$$

4.2.3 Big: Levi, Fatou, and Lebesgue Dominated Convergence

Levi's Theorem

For f_n a sequence of **integrable** functions such that $f_n \leq f_{n+1}$ a.e. for all n and $\lim \int f_n < \infty$, then there exists f integrable such that $f_n \uparrow f$ a.e.

(and $\lim \int f_n = \int f$)

“an integrable function waits at the top of an increasing sequence”

* f is defined a.e. on X

nice consequence: If integrable $f_n > 0$ a.e., with $\sum_{n=1}^{\infty} \int f_n < \infty$, then $\sum f_n$ defined an integrable function and

$$\int \sum_{n=1}^{\infty} f_n = \sum_{n=1}^{\infty} \int f_n$$

*not true in Riemann land!

* trick: when $f_1 \leq f_2 \leq \dots$ can make

a positive sequence by considering $f_1 - f_1, f_2 - f_1, \dots$

Fatou's Lemma

For integrable $f_n \geq 0$ a.e. for all n and $\liminf \int f_n < \infty$, then

$$\int \liminf f_n \leq \liminf \int f_n$$

where $\liminf f_n$ defines an integrable function a.e. on X .

"lim inf of integrable is integrable and less than integral of parts"

Lebesgue Dominated Convergence

1. **measurable** $f_n \rightarrow f$ a.e.
2. $|f_n| \leq g$ a.e. for g **integrable**

then

$$\int f = \lim_{n \rightarrow \infty} \int f_n$$

where f_n and f are integrable (for all n)

"interchange lim and \int for measurable functions bounded by an integrable function"

4.3 Riemann Integrals (section 23)

a partition P is just a collection of points inside an interval.

A second partition Q **refines** P if $P \subseteq Q$.

The **Upper** Riemann sum is

$$U(f, P) = \sum_i^n M_i(x_i - x_{i-1})$$

where M_i is the sup of f on $[x_{i-1}, x_i]$. (similarly lower sum is defined with m_i , the inf on the interval)

If Q refines P , then $U(f, Q) \leq U(f, P)$.
($L(f, Q) \geq L(f, P)$).

f is **Riemann integrable** if

$$\begin{aligned} \lim_{||P_i|| \rightarrow 0} U(f, P_i) &= \lim_{||P_i|| \rightarrow 0} L(f, P_i) \\ &= \int f(x) dx \end{aligned}$$

where $||P_i||$ the length of the largest subinterval.

* The Riemann integral of f can also be defined when $\sup L = \inf S$; this value is said to be the integral of f .

* **Riemann's Criterion** f is integrable if if L and U can be made arbitrarily close by selecting a sufficiently fine partition.

Every **Riemann** integrable function is **Lebesgue** integrable.

A **bounded function** $f : [a, b] \rightarrow \mathbb{R}$ is **Riemann** integrable

$$\iff$$

f is **continuous** a.e.

4.4 Product Measures and Iterated Integrals (section 26—only a sketch)

If S and T are semirings, then their cross-product: $S \times T$ is also a semiring. (similarly, for measures μ and ν)

a function $f : X \times Y$ is $\mu \times \nu$ integrable by computing cross-sections:

$$\int_{X \times Y} f d(\mu \times \nu) = \int_X \int_Y f d\mu d\nu$$

"Fubini" says the order of $\int_X \int_Y$ doesn't matter!

Tonelli's Theorem f is $\mu \times \nu$ -measurable and $\int_X \int_Y |f| d\nu d\mu$ exists (or other order) then $\int \int f$ exists.

5 Function Spaces (Chapter 5)

5.1 norms on vector spaces (section 27)

A real valued function $\|\cdot\|$ on a vector space V is a **norm** if for all v in V ,

1. $\|v\| > 0$ and $\|v\| = 0 \iff v = 0$
2. $\|\alpha v\| = |\alpha| \|v\|$, for all $\alpha \in \mathbb{R}$
3. $\|v + w\| \leq \|v\| + \|w\|$ (triangle)

* a norm space \implies a metric space (but not the converse)

Can show $|\|v\| - \|w\|| \leq \|v - w\|$.
by triangle

Examples of norms in different spaces:

- "Euclidean norm": $\sqrt{v_1^2 + v_2^2 + \dots}$
- "sup norm": $\|f\|_{sup} = \sup |f(x)|$ over all x .
(only valid in space of bounded, real-valued functions)
- " L^p " norm: $\|f\|_p = \left(\int |f|^p \right)^{1/p}$
only valid in $L^p(X)$ space = $\{f | f \text{ is measurable and } |f|^p \text{ is integrable}\}$
* note on \mathbb{R}^n , the L^p norm is:
 $\|(a_1, \dots, a_n)\|_p = (|a_1|^p + \dots + |a_n|^p)^{1/p}$

A **bounded normed** space is one where $\|v\| \leq M$ for some constant M .

A normed space (a vector space with a norm) is a **Banach** space if every Cauchy sequence converges (aka **complete**).

Two norms are equivalent if there are $K, M > 0$: $K\|x\|_1 \leq \|x\|_2 \leq M\|x\|_1$ for all x .

*In a finite dimensional vector space, all norms are equivalent

5.2 Linear Operators (section 28)

A **Linear Operator** (or transformation) is a map T between two vector spaces V and W such that:

$$T(aV + bW) = aT(V) + bT(W)$$

The **Operator norm** of T , $\|T\|$ is

$$\sup\{\|T(v)\| : \|v\| = 1\}$$

we say T is bounded if $\|T\|$ is finite.

What's equivalent to T being **bounded**?

1. $\|T(v)\| \leq M\|v\|$ for all v ($M \geq 0$)
2. T is continuous at zero
3. T is continuous
4. $\ker(T)$ is a closed set

The **collection** of **bounded** linear operators from V to W is denoted $L(V, W)$

Nice Properties:

- $L(V, W)$ is a normed linear space using $\|\cdot\|_{operator}$
- If $\dim V < \infty$ then every operator is a bounded operator.
- If W is complete (w.r.s to $\|\cdot\|_w$), then $L(V, W)$ is complete

$L(V, \mathbb{R})$ is always a **complete normed** linear space ("Banach space").

5.2.1 Dual Spaces

For any normed linear space V , the **dual space** of V is $L(V, \mathbb{R}) = V^*$

"bounded linear transformations from V to \mathbb{R} "

For any $1/p + 1/q = 1$ ($p, q > 1$), $(L^p)^* = L^q$.

"dual of L^p is equivalent to L^q "

* we get L^2 is self-dual

Also note the double dual $((L^p)^*)^* \xrightarrow{L^p}$, isometrically isomorphic.

5.3 Lp Spaces (section 31)

L^p is the collection functions f such that

- 1) f is measurable
- 2) $|f|^p$ is integrable

This collection, L^p forms a space.

We can define a norm on L^p $|||_P$ by

$$||f||_p = (\int |f|^p)^{1/p}$$

Proof of triangle inequality is called **Minkowski's Inequality** only holds for finite $p > 1$,

$$||f + g||_p \leq ||f||_p + ||g||_p$$

for $f, g \in L^p$.

Holder's Inequality says if $1/p + 1/q = 1$ (called "conjugate exponents") and $f \in L^p, g \in L^q$ then

$$\int |fg| \leq ||f||_p ||g||_q$$

*implies $fg \in L^1$

Riesz-Fischer L^p is complete (every Cauchy seq converges) for all $p \geq 1$ (with respect to L^p -norm)

*Careful: $||f - f_n||_p \rightarrow 0$ doesn't imply $f_n \rightarrow f$

we don't get pointwise convergence

* we do get that some subsequence converges pointwise

$$\lim ||f - f_n||_p = 0 \not\Rightarrow f_n \rightarrow f$$

However we do get:

If $\lim ||f - f_n||_p = 0$ (in $L^p, p \geq 1$), then there exists a subsequence of $f_n, f_k \rightarrow f$ a.e.

* cauchy \iff some subsequence converges pointwise

e.g., $X = [0, 1]$ and $\mu = \lambda$,

$$A_1 = [0, 1]$$

$$A_2 = [0, 1/2], [1/2, 1]$$

$$A_3 = [0, 1/3], [1/3, 2/3], [2/3, 1]$$

...

$$\text{Let } f_1 = \chi_{A_1}, f_2 = \chi_{A_2} \dots$$

Then for all n and any $p \geq 1$,

$$f_n^p = f_n$$

but $||f_n||^p \rightarrow 0$.

(there is a subsequence of f_n that converges pointwise to 0 a.e.)

another example is

$$X = \mathbb{R}, f_n = \chi_{[n, n+1]}, f_n^p = f_n \rightarrow 0 = f,$$

but

$f_n \not\rightarrow f$ in L^p , because $||f_n||_p = 1$ for any n .

If $f_n \rightarrow f$ a.e. and $\lim ||f_n||_p = ||f||_p$, then $\lim ||f - f_n||_p = 0$ "pointwise and norms approaching implies convergence in L^p "

5.3.1 Essentially Bounded Functions

If $|fg| \leq h$, some integrable function, then $fg \in L^1$.

a function f is **essentially bounded** if

$$|f(x)| \leq M$$

for almost all x .

The **essential supremum**, denoted by $||f||_\infty$ is

$$||f||_\infty = \inf M \mid |f(x)| \leq M \text{ for almost all } x$$

e.g., all constant functions are in L^∞ .

Also note:

* $L^\infty \not\subseteq L^p$ for any $p < \infty$ when

$$\mu(X) = \infty$$

* $L^\infty \subseteq L^p$ for any p if $\mu(X) < \infty$.

If X is σ -finite and $F \in (L^1)^*$ then there exists $g \in L^\infty$ such that for any $f \in L^1$

$$F(f) = \int fg$$

5.3.2 What is in L^p ?

$C_b(K)$ is the set of continuous and bounded functions on K .

e.g., If K is compact, then every continuous function is bounded (Extreme Value Theorem).

*can show $C_b(K)$ for K compact is a complete n.l.s using $|||_{sup}$.

Any **continuous** function with **compact support** is in $L^p(\mathbb{R})$ ($p \geq 1$)

*a function $f : \mathbb{R} \rightarrow \mathbb{R}$ has **compact support** if the closure of $f(\{x|f(x) \neq 0\})$ is compact.

*recall: uniform limit of continuous functions is continuous

On $[1, \infty)$,

$$1/x^a \in L^p \iff ap > 1$$

On $(0, 1]$

$$1/x^a \in L^p \iff ap < 1$$

In a finite measure space, $L^q \subseteq L^p$ if $1 \leq p \leq q$.

5.3.3 Dense Functions in L^p

A is **dense** in B if for any $b \in B$, there is an a such that $d(a, b) < \epsilon$

nice fact: For $A \subseteq B \subseteq M$, if A is dense in B and B is dense in M , then A is dense in M .

* The collection of step functions is **dense** in L^p (for $1 < p < \infty$).

Also linear combinations of $\chi_{(a,b)}$ are dense in L^p ($p \geq 1$).

For μ a **regular Borel measure** on a Hausdorff locally compact topological space X , the collection of continuous functions with compact support is **dense** in L^p (for $1 \leq p < \infty$).

* Hausdorff space is one where two points can be separated by open sets.

* locally compact means every point lives in a compact neighborhood.

* remember a regular borel measure has additional requirements on compact and borel sets see: 2.5.2.

notation: $C_c(X)$ is the set of continuous real-valued functions on X with compact support

What's more? The collection of Riemann integrable functions on $[a, b]$ is dense in $L^1([a, b])$

5.3.4 Counting Measure and l^p

$$l^p = L(\mathbb{N}, \text{counting})$$

For a sequence $f_n \in l^p$, $f \in l^p$

$$\iff$$

$$\int |f|^p \text{ converges}$$

meaning $\sum |f_n(\mathbb{N})|^p$ converges.

For $p \leq q$, if $f \in l^p$ then $f \in l^q$.

why? $f \in l^p \implies \sum |a_n|^p$ converges,

so $|a_n|^p < 1$

hence $|a_n|^q \leq |a_n|^p$

6 Hilbert Spaces (Chapter 6)

6.1 Inner Product Spaces (section 32)

an **inner product** on a vector space V is a function from $V \times V$ to \mathbb{R} such that

- "linear" $(ax + by, z) = a(x, z) + b(y, z)$ (for $x, y, z \in V$ and $a, b \in \mathbb{R}$)
- "symmetric" $(x, y) = (y, x)$
- "positive definite" $(x, x) \geq 0$ and $(x, x) = 0 \iff x = 0$.

In an **inner product space** (space with an inner product), the **norm** (induced by the inner product) of a vector $v \in V$ is

$$\|v\| = \sqrt{(v, v)}$$

Two vectors x, y are **orthogonal** ($x \perp y$) if $(x, y) = 0$. In a real vector space, we also have a notion of angles between vectors:

$$\cos(\theta) = \frac{(x, y)}{\|x\| \|y\|}$$

6.1.1 Useful Inequalities

- Cauchy-Schwarz: $|(x, y)| \leq \|x\| \|y\|$
- Parallelogram Law: $\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2$
 - a norm $\| \cdot \|$ is induced by an inner product \iff Parallelogram holds
 - in a real vector space we then have: $(x, y) = \frac{1}{4}(\|x + y\|^2 - \|x - y\|^2)$
- Bessel's Inequality: If $\{x_i\}$ is a collection of orthonormal vectors in an inner product space, $\sum_i |(x, x_i)|^2 \leq \|x\|^2$.
- Pythagorean Theorem: for x_1, x_2, \dots, x_n pairwise orthogonal, $\|x_1 + \dots + x_n\|^2 = \|x_1\|^2 + \dots + \|x_n\|^2$

6.2 Hilbert Spaces (section 33)

A **Hilbert space** is a complete **inner product space**.

* the norm is induced by the inner product (i.e. $\|x\| = \sqrt{(x, x)}$)

Examples:

- l^2 under the inner product: $(x, y) = \text{"dot product"}$
- $L^2(u)$ is another example

6.2.1 Orthogonal Complement and Spans

The **orthogonal complement** of a subset A of an inner product space X is

$$A^\perp = \{x \in X : x \perp y \text{ for all } y \in A\}$$

"set of vector orthogonal to all vectors in A "

* A^\perp is a vector subspace

*careful: $(A^\perp)^\perp \subseteq A$, but not necessarily equal to A

When X is a **Hilbert space** and A is a closed subspace, then A and A^\perp span the entire Hilbert space.

in this case, we have $(A^\perp)^\perp = A$

In a Hilbert space, a subspace of M is **dense** \iff only zero vector is orthogonal to M .

The elements of a subset of a Hilbert space are **linearly independent** if 0 is not in the set and the elements are pairwise \perp .

6.3 Orthonormal Bases

In a Hilbert space, for a family of orthonormal vectors $\{e_i\}$ the following are equivalent

- 0 is not in $\{e_i\}$ and $\{e_i\}$ spans a dense subset of H
- If $x \perp e_i$ for each i , then $x = 0$
- Parseval's Identity: for each vector x , $\|x\|^2 = \sum_i |(x, e_i)|^2$
- for each vector, $(x, e_i) \neq 0$ at most countably many times and $x = \sum_i (x, e_i) e_i$ converges

For an **orthonormal basis** $\{e_i\}$ in a Hilbert space, the family of scalars $\{(x, e_i)\}$ are the **Fourier coefficients**.

*Fourier coefficients are always with respect to a basis

Every Hilbert space H is **linearly isometric** (a linear, norm preserving map exists) to a Hilbert space of the form $l_2(\mathbb{Q})$.

Specifically, $L : H \rightarrow l_2(I)$ defined by $L(x) = \{(x, e_i)\}_i$ is such a map.

An infinite dimensional Hilbert space is **separable** \iff it's **linearly isometric** to l^2 .

In an infinite dimensional Hilbert space H ,

H is **separable**

\iff

it has a countable orthonormal basis

*in this case, every orthonormal basis is countable
"if one then all"

6.4 Fourier Analysis

6.4.1 Best Approximation Theorem

Let e_1, \dots be an orthonormal set (not necessarily a basis)

Define $H_N = \text{span}\{e_1, \dots, e_N\}$

The map $\pi_N : H \rightarrow H_N$ defined by

$$\pi_N(v) = \sum_i (v, e_i) e_i$$

is the **orthogonal projection of v into H_N** .

it's also a linear transformation and $(v - \pi_N(v)) \perp H_N$.

"Best Approximation Theorem": If $v \in H, w \in H_N$, then

$$\|v - w\| \geq \|v - \pi_N(v)\|$$

" $\pi_N(v)$ is the unique closest point to v in H_N "

If $H_1 \subseteq H_2 \subseteq \dots \subseteq H$ (finite dimensional) and $v \in H$, then

$$\|v - \pi_1(v)\| \geq \|v - \pi_2(v)\| \geq \dots$$

and $\pi_{N+1}(v) = \pi_N(v) + (v, e_{N+1})e_{N+1}$.

Stone-Weierstrauss

tells you about the collection continuous functions and density of vector subspace

Applying Stone-Weierstrauss to $L^2([0, 2\pi])$ we get:

for any continuous $f : [0, 2\pi] \rightarrow \mathbb{R}$, there is a polynomial $p(x)$ such that

$$|f(x) - p(x)| < \epsilon \text{ for any } x$$

6.4.2 Fourier coefficients

Let e_1, \dots be an orthonormal set in a separable Hilbert space H .

Define the i th Fourier coefficient as

$$\hat{e}_i(v) = (v, e_i)$$

* we can think about Best approx, Bessel, and Parseval's in terms of \hat{e}_i

If e_1, \dots is an orthonormal basis then for any v there exists $\alpha_1, \dots, \alpha_N$ in \mathbb{R} such that

$$\|v - \sum_{i=1}^N \alpha_i e_i\| < \epsilon.$$

Using the above with Best approximation Theorem we get

$$\lim_{N \rightarrow \infty} \pi_N(v) = v$$

(implying $\lim_{N \rightarrow \infty} \|\pi_N(v)\|^2 = \|v\|^2$) *in H -norm

6.4.3 Riesz-Fischer

Riesz-Fischer: Let e_1, \dots be an orthonormal basis of H , then the map

$$v \rightarrow \{\hat{e}_i(v)\}_i^\infty$$

is an **isometric isomorphism** of H with l^2 .

*isometric: is norm preserving

* up to isometric isomorphism, l^2 is the only separable countably finite dimensional Hilbert space.

6.4.4 Fourier on L^2

For $H = L^2(I)$ (I is an interval $[a, b]$),

$$\int_I |f|^2 = \sum_{i=1}^{\infty} \left| \int f e_i \right|^2$$

for all $f \in L^2(I)$.

* note: $\int f e_i = (f, e_i)$ in L^2

Every $f \in L^2(I)$ has a Fourier series that converges to f (not pointwise!)

Classical Fourier basis for L^2 are for $n = 1, 2, \dots$:

On $[0, 1]$: $1, \sqrt{2} \sin(2\pi nx), \sqrt{2} \cos(2\pi nx)$
 On $[0, 2\pi]$ or $[-\pi, \pi]$: $\frac{1}{\sqrt{2\pi}}, \frac{\sin(xn)}{\sqrt{\pi}}, \frac{\cos(xn)}{\sqrt{\pi}}$

For pointwise convergence, we restrict our attention to nicer functions:

If $f, f' \in C([0, 1])$ and $f(0) = f(1)$, then the classical Fourier series **converges absolutely and uniformly** to f .

* C^1 functions are dense in $L^p(I)$ (for example $L^2([0, 1])$)

Jordan's Theorem for f **sectionally continuous** on $[0, 1]$, then Fourier series converges to

$$\frac{1}{2}f(x_0^+) + \frac{1}{2}f(x_0^-)$$

at x_0 if left/right hand derivatives exist.

Specifically, Fourier series converges to $f(x_0)$ if f is continuous at x_0 .

Riemann-Lebesgue Lemma: If $f(x) \in L^p([a, b])$ then

$$\lim_{m \rightarrow \infty} \int_a^b f(t) \sin(mt) = 0$$

(same for cos).

*this implies for any $f \in L^p$, the Fourier coefficients $\rightarrow 0$, as $m \rightarrow \infty$.

6.5 Riesz-Representation Theorem

f is a **positive** function if $f(x) \geq 0$ for all x .

A continuous **linear functional** F (map from $C_c(X) \rightarrow \mathbb{R}$) is **positive** if $F(f) \geq 0$ for all positive f

*note F is in the dual space of $C_c(X)$

Riesz-Representation Theorem: If F is any positive linear functional on $C_c(X)$ then there exists a regular Borel measure μ such that $F_u(f) = \int f = F(f)$.

*can show map is linear: $F(af + bg) = aF(f) + bF(g)$.

7 Miscellaneous Examples

- In $X = \mathbb{R}, \mu = \lambda, f(x) = 1$ isn't integrable, but if $f(x) = 1$ on $[a, b]$ and zero elsewhere it is!

- $L^p(I)$ is a **separable** metric space

- $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$

why? compute fourier for $f(x) = x$ on $[-\pi, \pi]$

- For $l^2 = L^2(\mathbb{N})$ with counting measure, $\{\chi_{\{i\}}\}_{i \in \mathbb{N}}$ is an orthonormal basis of l^2 .

- $\sum \frac{a_n}{n}$ (fourier coefficients) converge absolutely for $f \in L^2([0, 1])$.

8 Questions

1.