

# Quantum Field Theory on a Highly Symmetric Lattice

Marco Aliberti

Università degli Studi di Torino

July 6, 2023

# Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated.


# Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated. There are two possible approaches:

Perturbative




# Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form


$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated. There are two possible approaches:

Perturbative




Non-Perturbative




# Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small  $g \Leftrightarrow$  Feynman diagrams with  $n$  loops

# Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small  $g \Leftrightarrow$  Feynman diagrams with  $n$  loops
- UV divergencies need to be eliminated

# Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small  $g \Leftrightarrow$  Feynman diagrams with  $n$  loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as  $g \rightarrow 0$

# Perturbative vs Non-Perturbative

Perturbative



Non-Perturbative




- Straightforward series expansion in powers of small  $g \Leftrightarrow$  Feynman diagrams with  $n$  loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as  $g \rightarrow 0$
- No straightforward approach




# Perturbative vs Non-Perturbative

## Perturbative



- Straightforward series expansion in powers of small  $g \Leftrightarrow$  Feynman diagrams with  $n$  loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as  $g \rightarrow 0$


## Non-Perturbative



- No straightforward approach
- Can have a natural cut-off for high momenta  $\Rightarrow$  No UV divergencies


# Perturbative vs Non-Perturbative

## Perturbative



- Straightforward series expansion in powers of small  $g \Leftrightarrow$  Feynman diagrams with  $n$  loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as  $g \rightarrow 0$

## Non-Perturbative



- No straightforward approach
- Can have a natural cut-off for high momenta  $\Rightarrow$  No UV divergencies
- Can predict quantities with essential singularities as  $g \rightarrow 0$

# Lattice Field Theory

## Definition: Lattice $\Lambda$

$$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z}, \{e_i\} \text{ basis of } \mathbb{R}^n \}$$

Hypercubic lattice:  $\{e_i\}$  is the canonical basis of  $\mathbb{R}^n$ ,  $a$  is called *lattice spacing*.




Figure: Example of a cubic lattice in  $\mathbb{R}^3$ .

## Definition: Lattice $\Lambda$

$$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z}, \{e_i\} \text{ basis of } \mathbb{R}^n \}$$

Hypercubic lattice:  $\{e_i\}$  is the canonical basis of  $\mathbb{R}^n$ ,  $a$  is called *lattice spacing*.

Basic idea: fields can take values only in given parts of the lattice,  $x \rightarrow n \in \Lambda$ .




Figure: Example of a cubic lattice in  $\mathbb{R}^3$ .

# Lattice Field Theory

## Definition: Lattice $\Lambda$

$$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z}, \{e_i\} \text{ basis of } \mathbb{R}^n \}$$

Hypercubic lattice:  $\{e_i\}$  is the canonical basis of  $\mathbb{R}^n$ ,  $a$  is called *lattice spacing*.

Basic idea: fields can take values only in given parts of the lattice,  $x \rightarrow n \in \Lambda$ .

Examples:

- Scalar fields  $\Phi(x) \rightarrow \Phi(n)$  on sites




Figure: Example of a cubic lattice in  $\mathbb{R}^3$ .

# Lattice Field Theory

## Definition: Lattice $\Lambda$

$$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z}, \{e_i\} \text{ basis of } \mathbb{R}^n \}$$

Hypercubic lattice:  $\{e_i\}$  is the canonical basis of  $\mathbb{R}^n$ ,  $a$  is called *lattice spacing*.

Basic idea: fields can take values only in given parts of the lattice,  $x \rightarrow n \in \Lambda$ .

Examples:

- Scalar fields  $\Phi(x) \rightarrow \Phi(n)$  on sites
- Vector fields  $A_\mu(x) \rightarrow A_\mu(n)$  on links




Figure: Example of a cubic lattice in  $\mathbb{R}^3$ .

# Lattice Field Theory

## Definition: Lattice $\Lambda$

$$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z}, \{e_i\} \text{ basis of } \mathbb{R}^n \}$$

Hypercubic lattice:  $\{e_i\}$  is the canonical basis of  $\mathbb{R}^n$ ,  $a$  is called *lattice spacing*.

Basic idea: fields can take values only in given parts of the lattice,  $x \rightarrow n \in \Lambda$ .

Examples:

- Scalar fields  $\Phi(x) \rightarrow \Phi(n)$  on sites
- Vector fields  $A_\mu(x) \rightarrow A_\mu(n)$  on links
- Object with  $k$  indices on  $k$ -simplexes




Figure: Example of a cubic lattice in  $\mathbb{R}^3$ .

# Lattice Field Theory

## Definition: Lattice $\Lambda$

$$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z}, \{e_i\} \text{ basis of } \mathbb{R}^n \}$$

Hypercubic lattice:  $\{e_i\}$  is the canonical basis of  $\mathbb{R}^n$ ,  $a$  is called *lattice spacing*.

Basic idea: fields can take values only in given parts of the lattice,  $x \rightarrow n \in \Lambda$ .

Examples:

- Scalar fields  $\Phi(x) \rightarrow \Phi(n)$  on sites
- Vector fields  $A_\mu(x) \rightarrow A_\mu(n)$  on links
- Object with  $k$  indices on  $k$ -simplexes

## Attention!

Spinorial fields are trickier.




Figure: Example of a cubic lattice in  $\mathbb{R}^3$ .