Quantum Field Theory on a Highly Symmetric Lattice

Marco Aliberti

Università degli Studi di Torino

July 5, 2023

Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i
angle = \int_{\phi_i}^{\phi_f} \mathcal{D}\left[\phi
ight] \mathrm{e}^{-S\left[\phi
ight]}$$

need to be evaluated.

Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i
angle = \int_{\phi_i}^{\phi_f} \mathcal{D}\left[\phi
ight] \mathrm{e}^{-S\left[\phi
ight]}$$

need to be evaluated. There are two possible approaches:

Perturbative



Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i
angle = \int_{\phi_i}^{\phi_f} \mathcal{D}\left[\phi
ight] \mathrm{e}^{-S\left[\phi
ight]}$$

need to be evaluated. There are two possible approaches:

Perturbative

Non-Perturbative



Perturbative



• Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops



Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated



Perturbative



- Straightforward series expansion in powers of small g ⇔
 Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \to 0$



Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \to 0$

Non-Perturbative



No straightforward approach

Perturbative



- Straightforward series expansion in powers of small g ⇔
 Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as g o 0



- No straightforward approach
- Can have a natural cut-off for high momenta ⇒ No UV divergencies

Perturbative



- Straightforward series expansion in powers of small g ⇔
 Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$



- No straightforward approach
- Can have a natural cut-off for high momenta ⇒ No UV divergencies
- ullet Can predict quantities with essential singularities as g o 0