

Quantum Field Theory on a Highly Symmetric Lattice

Marco Aliberti

Università degli Studi di Torino

July 6, 2023

Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated.


Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated. There are two possible approaches:

Perturbative




Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form


$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated. There are two possible approaches:

Perturbative




Non-Perturbative




Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops

Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated

Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$

Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$
- No straightforward approach


Perturbative vs Non-Perturbative

Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$


Non-Perturbative



- No straightforward approach
- Can have a natural cut-off for high momenta \Rightarrow No UV divergencies


Perturbative vs Non-Perturbative

Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$

Non-Perturbative



- No straightforward approach
- Can have a natural cut-off for high momenta \Rightarrow No UV divergencies
- Can predict quantities with essential singularities as $g \rightarrow 0$

What is a Lattice?

Definition: Lattice Λ

$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z} \}$, with $\{e_i\}$ any basis of \mathbb{R}^n




Figure: A bidimensional lattice.

What is a Lattice?

Definition: Lattice Λ

$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z} \}$, with $\{e_i\}$ any basis of \mathbb{R}^n




Figure: A bidimensional lattice.

Hypercubic lattice

$\{e_i\}$ is the canonical basis of \mathbb{R}^n
 a is called *lattice spacing*.




Figure: A square lattice.

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites
- **Vector fields** $A_\mu(x) \rightarrow A_\mu(n)$ on links




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites
- **Vector fields** $A_\mu(x) \rightarrow A_\mu(n)$ on links
- Object with k indices on k -simplexes




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites
- **Vector fields** $A_\mu(x) \rightarrow A_\mu(n)$ on links
- Object with k indices on k -simplexes




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Attention!

Spinorial fields are trickier to be discretized.