

Quantum Field Theory on a Highly Symmetric Lattice

Marco Aliberti

Università degli Studi di Torino

14th October, 2023

Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated.


Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated. There are two possible approaches:

Perturbative




Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form


$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated. There are two possible approaches:

Perturbative




Non-Perturbative




Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops

Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated

Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$

Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$
- No straightforward approach


Perturbative vs Non-Perturbative

Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$


Non-Perturbative



- No straightforward approach
- Can have a natural cut-off for high momenta \Rightarrow No UV divergencies


Perturbative vs Non-Perturbative

Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$

Non-Perturbative



- No straightforward approach
- Can have a natural cut-off for high momenta \Rightarrow No UV divergencies
- Can predict quantities with essential singularities as $g \rightarrow 0$

What is a Lattice?

Definition: Lattice Λ

$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z} \}$, with $\{e_i\}$ any basis of \mathbb{R}^n




Figure: A bidimensional lattice.

What is a Lattice?

Definition: Lattice Λ

$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z} \}$, with $\{e_i\}$ any basis of \mathbb{R}^n




Figure: A bidimensional lattice.

Hypercubic lattice

$\{e_i\}$ is the canonical basis of \mathbb{R}^n
 a is called *lattice spacing*.




Figure: A square lattice.

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites
- **Vector fields** $U_\mu(x) \rightarrow U_\mu(n)$ on links

Parallel Transporter

$$U_\mu(x) = \exp(igaA_\mu(x))$$




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites
- **Vector fields** $U_\mu(x) \rightarrow U_\mu(n)$ on links
- Object with k indices on k -simplexes

Parallel Transporter

$$U_\mu(x) = \exp(igaA_\mu(x))$$




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites
- **Vector fields** $U_\mu(x) \rightarrow U_\mu(n)$ on links
- Object with k indices on k -simplexes

Beware!

Spinorial fields are trickier to be discretized.

Parallel Transporter

$$U_\mu(x) = \exp(igaA_\mu(x))$$




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Gauge-Invariant Observables and Wilson Action

The Yang-Mills continuum action is
$$S_E = \frac{1}{4} \int d^4x F^{a\mu\nu}(x) F_{\mu\nu}^a(x).$$

On the lattice, every closed path is gauge-invariant.




Figure: Gauge-invariant paths on a bidimensional lattice. [Sigdel, 2016]

Gauge-Invariant Observables and Wilson Action

The Yang-Mills continuum action is
$$S_E = \frac{1}{4} \int d^4x F^{a\mu\nu}(x) F_{\mu\nu}^a(x).$$

On the lattice, every closed path is gauge-invariant.

Definition: Plaquette $U_{\mu\nu}(n)$

$$U_\mu(n) U_\nu(n + \mu) U_\mu^\dagger(n + \nu) U_\nu^\dagger(n)$$




Figure: Gauge-invariant paths on a bidimensional lattice. [Sigdel, 2016]

Gauge-Invariant Observables and Wilson Action

The Yang-Mills continuum action is
$$S_E = \frac{1}{4} \int d^4x F^{a\mu\nu}(x) F_{\mu\nu}^a(x).$$

Definition: Plaquette $U_{\mu\nu}(n)$

$$U_\mu(n) U_\nu(n + \mu) U_\mu^\dagger(n + \nu) U_\nu^\dagger(n)$$

On the lattice, every closed path is gauge-invariant.

Wilson's Idea

$$S = \frac{\beta}{2N} \sum_{n,\mu,\nu} \Re \text{Tr} (1 - U_{\mu\nu}(n))$$




Figure: Gauge-invariant paths on a bidimensional lattice. [Sigdel, 2016]

Polyakov Loops and Potential

If the time coordinate is taken to be periodic, more closed paths arise.

Polyakov Loop

$$P(n) = \text{Tr} \prod_{t=0}^{T-1} U_t(n)$$




Figure: Gauge-invariant paths on a bidimensional lattice. [Sigdel, 2016]

Polyakov Loops and Potential

If the time coordinate is taken to be periodic, more closed paths arise.

The expectation value of two Polyakov loops is the potential.

Polyakov Loop

$$P(n) = \text{Tr} \prod_{t=0}^{T-1} U_t(n)$$

Potential

$$V(R) = -\frac{1}{T} \log \langle P(0) P^\dagger(R) \rangle$$




Figure: Gauge-invariant paths on a bidimensional lattice. [Sigdel, 2016]

Poincaré Group can be divided in:

Translations

Rotations

Poincaré Group can be divided in:

Translations

$$x^\mu \rightarrow x^\mu + \varepsilon^\mu$$

\Downarrow

$$n \rightarrow n + a\hat{\mu}$$

Rotations

Poincaré Group can be divided in:

Translations

$$x^\mu \rightarrow x^\mu + \varepsilon^\mu$$

$$\Downarrow$$

$$n \rightarrow n + a\hat{\mu}$$

Rotations

$$x^\mu \rightarrow R^\mu_\nu x^\nu \quad R \in SO(4)$$

$$\Downarrow$$

$$n \rightarrow \Gamma n \quad \Gamma \in T$$

T : group of rotations of multiples of 90° around any axis.

Poincaré Group can be divided in:

Translations

$$x^\mu \rightarrow x^\mu + \varepsilon^\mu$$

\Downarrow

$$n \rightarrow n + a\hat{\mu}$$

$$a\hat{\mu} \rightarrow \varepsilon^\mu \text{ for } a \rightarrow 0$$

Rotations

$$x^\mu \rightarrow R^\mu_\nu x^\nu \quad R \in SO(4)$$

\Downarrow

$$n \rightarrow \Gamma n \quad \Gamma \in T$$

T : group of rotations of multiples of 90° around any axis.

Lattice Symmetries

Poincaré Group can be divided in:

Translations

$$x^\mu \rightarrow x^\mu + \varepsilon^\mu$$

\Downarrow

$$n \rightarrow n + a\hat{\mu}$$

$$a\hat{\mu} \rightarrow \varepsilon^\mu \text{ for } a \rightarrow 0$$

Rotations

$$x^\mu \rightarrow R^\mu_\nu x^\nu \quad R \in SO(4)$$

\Downarrow

$$n \rightarrow \Gamma n \quad \Gamma \in T$$

T : group of rotations of multiples of 90° around any axis.

$$\Gamma \rightarrow R \text{ for } a \rightarrow 0$$

Lattice Symmetries

Poincaré Group can be divided in:

Translations

$$x^\mu \rightarrow x^\mu + \varepsilon^\mu$$



$$n \rightarrow n + a\hat{\mu}$$

$$a\hat{\mu} \rightarrow \varepsilon^\mu \text{ for } a \rightarrow 0$$

Rotations

$$x^\mu \rightarrow R^\mu_\nu x^\nu \quad R \in SO(4)$$



$$n \rightarrow \Gamma n \quad \Gamma \in T$$

T : group of rotations of multiples of 90° around any axis.

$$\Gamma \not\rightarrow R \text{ for } a \rightarrow 0$$

Important:

Rotational invariance seems to be broken.

Rotational Invariance Restoration - Lang and Rebbi

Equipotential surfaces become spheres as the continuum limit is approached.




Figure: Restoration of rotational invariance from (a) $\beta = 2$, $n_s = 8$, $n_t = 4$ to (b) $\beta = 2.25$, $n_s = 16$, $n_t = 6$; the curves represent equipotential curves.
[Lang and Rebbi, 1982]

Rotational Invariance Restoration

Values of β are slightly different from Lang and Rebbi's because $a(\beta) \approx \Lambda e^{-b_0 \beta}$, with $\Lambda, b_0 > 0$ ¹.




Figure: Potential from $\beta = 2.20$, $n_s = 8$, $n_t = 4$.




Figure: Potential from $\beta = 2.35$, $n_s = 16$, $n_t = 6$.

¹The simulation code is based on the code presented in refs. [Panero, 2009; Mykkänen, Panero, and Rummukainen, 2012].

Higher Symmetry Lattices

Other, more rotational-symmetric, lattices have been used:

Body Centered Tesseract

[Celmaster, 1982]

- 24 nearest neighbours
- 1152-element symmetry group




Figure: Two-dimensional projection of a BCT. [Celmaster, 1982]

The SH lattice has 8 nearest neighbours and a 288-element symmetry group

Higher Symmetry Lattices

Other, more rotational-symmetric, lattices have been used:

Body Centered Tesseract

[Celmaster, 1982]

- 24 nearest neighbours
- 1152-element symmetry group




Figure: Two-dimensional projection of a BCT. [Celmaster, 1982]

F_4 coroots lattice [Neuberger, 1987]

- 48 nearest neighbours
- 2304-element symmetry group




Figure: Two-dimensional projection of a F_4 coroots lattice. [Wikipedia, 2023]

F_4 Coroots Lattice

- Obtained from the roots lattice of the exceptional Lie algebra F_4 and its dual;




Figure:

F_4 Coroots Lattice

- Obtained from the roots lattice of the exceptional Lie algebra F_4 and its dual;
- Has 48 nearest neighbours:
 - The 24 roots are all possible permutations of coordinate positions of $(\pm 1, \pm 1, 0, 0)$




Figure: The 24 roots (red) of the F_4 lattice, projected on a bidimensional plane.

F_4 Coroots Lattice

- Obtained from the roots lattice of the exceptional Lie algebra F_4 and its dual;
- Has 48 nearest neighbours:
 - The 24 roots are all possible permutations of coordinate positions of $(\pm 1, \pm 1, 0, 0)$
 - The 24 dual roots (coroots) are:
 - The 8 possible permutations of $(\pm 1, 0, 0, 0)$
 - The 16 vectors of the form $(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2})$




Figure: The 24 roots (red) and the 24 coroots (yellow) of the F_4 lattice, projected on a bidimensional plane.

F_4 Coroots Lattice

- Obtained from the roots lattice of the exceptional Lie algebra F_4 and its dual;
- Has 48 nearest neighbours:
 - The 24 roots are all possible permutations of coordinate positions of $(\pm 1, \pm 1, 0, 0)$
 - The 24 dual roots (coroots) are:
 - The 8 possible permutations of $(\pm 1, 0, 0, 0)$
 - The 16 vectors of the form $(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2})$
- Exists only in 4 dimensions;




Figure: The 24 roots (red) and the 24 coroots (yellow) of the F_4 lattice, projected on a bidimensional plane.

F_4 Coroots Lattice

- Obtained from the roots lattice of the exceptional Lie algebra F_4 and its dual;
- Has 48 nearest neighbours:
 - The 24 roots are all possible permutations of coordinate positions of $(\pm 1, \pm 1, 0, 0)$
 - The 24 dual roots (coroots) are:
 - The 8 possible permutations of $(\pm 1, 0, 0, 0)$
 - The 16 vectors of the form $(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2})$
- Exists only in 4 dimensions;
- Contains the Simple Hypercubic lattice and the BCT;




Figure: The 24 roots (red) and the 24 coroots (yellow) of the F_4 lattice, projected on a bidimensional plane.

F_4 Coroots Lattice

- Obtained from the roots lattice of the exceptional Lie algebra F_4 and its dual;
- Has 48 nearest neighbours:
 - The 24 roots are all possible permutations of coordinate positions of $(\pm 1, \pm 1, 0, 0)$
 - The 24 dual roots (coroots) are:
 - The 8 possible permutations of $(\pm 1, 0, 0, 0)$
 - The 16 vectors of the form $(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2})$
- Exists only in 4 dimensions;
- Contains the Simple Hypercubic lattice and the BCT;
- Has been used only to simulate scalar fields, in [Neuberger, 1987].







Figure: The 24 roots (red) and the 24 coroots (yellow) of the F_4 lattice, projected on a bidimensional plane.

- Implement the F_4 lattice in the simulation program and make efficiency studies;



- Implement the F_4 lattice in the simulation program and make efficiency studies;
- Make a rotational invariance study on the new lattice, hoping to get better results than the Simple Hypercubic lattice.



Thank you for your attention



Sigdel, Dibakar (2016). “Two Dimensional Lattice Gauge Theory with and without Fermion Content”. In: *FIU Electronic Theses and Dissertations* 3224. DOI: 10.25148/etd.FIDC001748. URL: https://digitalcommons.fiu.edu/etd/3224?utm_source=digitalcommons.fiu.edu%2Fetd%2F3224&utm_medium=PDF&utm_campaign=PDFCoverPages.



Lang, C. B. and C. Rebbi (1982). “Potential and Restoration of Rotational Symmetry in $SU(2)$ Lattice Gauge Theory”. In: *Phys. Lett.* B115. [, 322 (1982)], p. 137. DOI: 10.1016/0370-2693(82)90813-9.



Panero, Marco (2009). “Thermodynamics of the QCD plasma and the large- N limit”. In: *Phys. Rev. Lett.* 103, p. 232001. DOI: 10.1103/PhysRevLett.103.232001. arXiv: 0907.3719 [hep-lat].



Mykkänen, Anne, Marco Panero, and Kari Rummukainen (2012). “Casimir scaling and renormalization of Polyakov loops in large- N gauge theories”. In: *JHEP* 1205, p. 069. DOI: [10.1007/JHEP05\(2012\)069](https://doi.org/10.1007/JHEP05(2012)069). arXiv: [1202.2762](https://arxiv.org/abs/1202.2762) [hep-lat].



Celmaster, William (1982). “Gauge Theories on the Body - Centered Hypercubic Lattice”. In: *Phys. Rev. D* 26, p. 2955. DOI: [10.1103/PhysRevD.26.2955](https://doi.org/10.1103/PhysRevD.26.2955).



Neuberger, Herbert (1987). “SPINLESS FIELDS ON F(4) LATTICES”. In: *Phys. Lett. B* 199, pp. 536–540. DOI: [10.1016/0370-2693\(87\)91623-6](https://doi.org/10.1016/0370-2693(87)91623-6).



Wikipedia (2023). URL: https://en.wikipedia.org/wiki/F4_%28mathematics%29 (visited on 08/29/2023).